




Contents lists available at ScienceDirect

Journal of Symbolic Computation

journal homepage: www.elsevier.com/locate/jsc

Computation of 2-groups of narrow logarithmic divisor classes of number fields

Jean-François Jaulent^{a,1}, Sebastian Pauli^b, Michael E. Pohst^c,
Florence Soriano-Gafiuk^d

^a Université de Bordeaux, Institut de Mathématiques de Bordeaux, 351, Cours de la Libération, 33405 Talence Cedex, France

^b University of North Carolina at Greensboro, Department of Mathematics and Statistics, Greensboro, NC 27402, USA

^c Technische Universität Berlin, Institut für Mathematik MA 8-1, Straße des 17. Juni 136, 10623 Berlin, Germany

^d Université Paul Verlaine de Metz, LMAM, Ile du Saulcy, 57000 Metz, France

ARTICLE INFO

Article history:

Received 29 November 2007

Accepted 15 April 2008

Available online 30 September 2008

Keywords:

Logarithmic class group

Wild kernel

Narrow logarithmic classes

ABSTRACT

We present an algorithm for computing the 2-group $\widetilde{\mathcal{C}}_F^{\text{res}}$ of narrow logarithmic divisor classes of degree 0 for number fields F . As an application, we compute in some cases the 2-rank of the wild kernel $WK_2(F)$ and the 2-rank of its subgroup $K_2^\infty(F) := \bigcap_{n \geq 1} K_2^n(F)$ of infinite height elements in $K_2(F)$.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

Jaulent (1994a) pointed out that the wild kernel $WK_2(F)$ of a number field F can also be studied via logarithmic class groups, the arithmetic of which he therefore developed (Jaulent, 1994b).

More precisely, if F contains a primitive 2ℓ th root of unity, the ℓ -rank of $WK_2(F)$ coincides with the ℓ -rank of the logarithmic class group $\mathcal{C}\ell_F$. So an algorithm for computing $\mathcal{C}\ell_F$ for Galois extensions F was developed first by Diaz y Diaz and Soriano (1999) and later generalized and improved for an arbitrary number field by Diaz y Diaz et al. (2005).

In case the prime ℓ is odd and the field F does not contain a primitive ℓ th root of unity one considers the cyclotomic extension $F' := F(\zeta_\ell)$, uses the isomorphism

$$\mu_\ell \otimes \widetilde{\mathcal{C}}\ell_{F'} \simeq WK_2(F')/WK_2(F')^\ell,$$

E-mail addresses: Jean-Francois.Jaulent@math.u-bordeaux1.fr (J.-F. Jaulent), s_pauli@uncg.edu (S. Pauli), pohst@math.tu-berlin.de (M.E. Pohst), soriano@univ-metz.fr (F. Soriano-Gafiuk).

¹ Tel.: +33 5 40 00 61 33; fax: +33 5 40 00 69 55.

and gets back to F via the so-called transfer (Jaulent and Soriano-Gafiuk, 2001; Soriano-Gafiuk, 2000; Jaulent and Michel, 2006). The algorithmic aspect of this is treated by Pauli and Soriano-Gafiuk (2004).

In case $\ell = 2$, whenever the condition $\zeta_{2\ell} \in F$ is not fulfilled, the relationship between logarithmic classes and exotic symbols is more intricate. For instance, when the number field F has a real embedding, Soriano (1997) observed that one may then define a narrow version of the logarithmic class group by analogy with the classical ideal class groups; and she used this for approximating the wild kernel more closely (Soriano-Gafiuk, 2000). But, unexpectedly, the 2-rank of this restricted logarithmic class group $\mathcal{C}\ell_F^{\text{res}}$ sometimes differs from the 2-rank of the group $WK_2(F)$. Moreover, in this case the wild kernel $WK_2(F)$ may differ from its subgroup $K_2^\infty(F) := \bigcap_{n \geq 1} K_2^n(F)$ of infinite height elements in $K_2(F)$. This was observed by J. Tate and then made more explicit by Hutchinson (2001, 2004).

That last difficulty was finally solved by Jaulent and Soriano-Gafiuk (2004); the authors constructed a positive class group *ad hoc* $\mathcal{C}\ell_F^{\text{pos}}$ which has the same 2-rank as the wild kernel $WK_2(F)$. Nevertheless, in case the set PE_F of dyadic exceptional primes of the number field F is empty, that group $\mathcal{C}\ell_F^{\text{pos}}$ appears as a factor of the full narrow logarithmic class group $\mathcal{C}\ell_F^{\text{res}}$ (without any assumption on the degree), so one may still use narrow logarithmic classes in order to compute the 2-rank of the wild kernel.

In the present paper we use the results by Diaz Diaz et al. (2005) on (ordinary) logarithmic class groups and develop an algorithm for computing the narrow groups $\mathcal{C}\ell_F^{\text{res}}$ in arbitrary number fields F . As a consequence, this algorithm calculates the 2-rank of the wild kernel $WK_2(F)$ whenever the field F has no dyadic exceptional places.

The computation of the 2-rank of $WK_2(F)$ in the remaining case ($PE_F \neq \emptyset$) will be solved in a forthcoming article where we compute the finite positive class group $\mathcal{C}\ell_F^{\text{pos}}$ and its subgroup $\widetilde{\mathcal{C}\ell}_F^{\text{pos}}$ of positive classes of degree 0.

2. The group of narrow logarithmic classes $\widetilde{\mathcal{C}\ell}_F^{\text{res}}$

In this preliminary section we recall the definition and the main properties of the arithmetic of restricted (or narrow) logarithmic classes. Jaulent (1994b) and Soriano-Gafiuk (2000) give a more detailed account.

Throughout this paper the prime number ℓ equals 2 and F is a number field of degree $n = r + 2c$ with r real places, c complex places and d dyadic places.

According to Jaulent (1994a), for every finite place \mathfrak{p} of F there exists a 2-adic \mathfrak{p} -valuation $\tilde{v}_{\mathfrak{p}}$ which is related to the wild \mathfrak{p} -symbol in case the cyclotomic \mathbb{Z}_2 -extension of $F_{\mathfrak{p}}$ contains i . The degree $\deg_{F,\mathfrak{p}}$ of the place \mathfrak{p} is a 2-adic integer such that the image of $\mathcal{R}_F := \mathbb{Z}_2 \otimes_{\mathbb{Z}} F^\times$ under the map $\text{Log} \mid_{\mathfrak{p}}$ is the \mathbb{Z}_2 -module $\deg_{F,\mathfrak{p}} \mathbb{Z}_2$ (Jaulent, 1994b), where Log denotes the usual 2-adic logarithm and $\mid_{\mathfrak{p}}$ is the 2-adic absolute value at the place \mathfrak{p} . The construction of the 2-adic logarithmic valuations $\tilde{v}_{\mathfrak{p}}$ yields:

$$\forall \alpha \in \mathcal{R}_F := \mathbb{Z}_2 \otimes_{\mathbb{Z}} F^\times : \sum_{\mathfrak{p} \in \text{Pl}_F^0} \tilde{v}_{\mathfrak{p}}(\alpha) \deg_{F,\mathfrak{p}} = 0,$$

where Pl_F^0 is the set of finite places of the number field F . Setting

$$\widetilde{\text{div}}_F(\alpha) := \sum_{\mathfrak{p} \in \text{Pl}_F^0} \tilde{v}_{\mathfrak{p}}(\alpha) \mathfrak{p}$$

with values in $\mathcal{D}\ell_F := \bigoplus_{\mathfrak{p} \in \text{Pl}_F^0} \mathbb{Z}_2 \mathfrak{p}$, we obtain by \mathbb{Z}_2 -linearity:

$$\deg_F(\widetilde{\text{div}}_F(\alpha)) = 0. \tag{1}$$

We then define the subgroup of logarithmic divisors of degree 0 by:

$$\widetilde{\mathcal{D}}\ell_F := \left\{ \alpha = \sum_{\mathfrak{p} \in \text{Pl}_F^0} a_{\mathfrak{p}} \mathfrak{p} \in \mathcal{D}\ell_F \mid \deg_F \alpha := \sum_{\mathfrak{p} \in \text{Pl}_F^0} a_{\mathfrak{p}} \deg_{F,\mathfrak{p}} = 0 \right\}$$

and the group of principal logarithmic divisors as the image of \mathcal{R}_F by $\widetilde{\text{div}}_F$:

$$\widetilde{\mathcal{P}}\ell_F := \{ \widetilde{\text{div}}_F(\alpha) \mid \alpha \in \mathcal{R}_F \}.$$

Due to (1), $\widetilde{\mathcal{P}}\ell_F$ is a subgroup of $\widetilde{\mathcal{D}}\ell_F$. Also, by the so-called extended Gross conjecture the factor group

$$\widetilde{\mathcal{C}}\ell_F := \widetilde{\mathcal{D}}\ell_F / \widetilde{\mathcal{P}}\ell_F$$

is a finite group, the 2-group of logarithmic divisor classes (of degree 0) of the field F introduced by Jaulent (1994b).

Now let $\mathcal{P}\mathcal{R}_F := \{p_1, \dots, p_r\}$ be the (non-empty) set of real places of the field F and F^+ be the subgroup of all totally positive elements in F^\times , i.e. the kernel of the sign-map

$$\text{sign}_F^\infty : F^\times \rightarrow \{\pm 1\}^r$$

which maps $x \in F$ onto the vector of the signs of the real conjugates of x . For

$$\widetilde{\mathcal{P}}\ell_F^+ := \{ \widetilde{\text{div}}_F(\alpha) \mid \alpha \in \mathcal{R}_F^+ := \mathbb{Z}_2 \otimes_{\mathbb{Z}} F^+ \}$$

Soriano (1997) introduced the factor group

$$\widetilde{\mathcal{C}}\ell_F^{\text{res}} := \widetilde{\mathcal{D}}\ell_F / \widetilde{\mathcal{P}}\ell_F^+,$$

which is called 2-group of narrow (or restricted) logarithmic divisor classes (of degree 0). Under the extended Gross conjecture $\widetilde{\mathcal{C}}\ell_F^{\text{res}}$ is finite.

In order to make it more suitable for actual computations, we may define it in a slightly different way by introducing real signed divisors of degree 0:

Definition 2.1. With the notations above, the 2-group of real signed logarithmic divisors (of degree 0) is the direct sum:

$$\widetilde{\mathcal{D}}\ell_F^{\text{res}} := \widetilde{\mathcal{D}}\ell_F \oplus \{\pm 1\}^r;$$

and the subgroup of principal real signed logarithmic divisors is the image:

$$\widetilde{\mathcal{P}}\ell_F^{\text{res}} := \{ (\widetilde{\text{div}}_F(\alpha), \text{sign}_F^\infty(\alpha)) \mid \alpha \in \mathcal{R}_F \}$$

of $\mathcal{R}_F := \mathbb{Z}_2 \otimes_{\mathbb{Z}} F^\times$ under the $(\widetilde{\text{div}}_F, \text{sign}_F^\infty)$ map. The factor group:

$$\widetilde{\mathcal{C}}\ell_F^{\text{res}} := \widetilde{\mathcal{D}}\ell_F^{\text{res}} / \widetilde{\mathcal{P}}\ell_F^{\text{res}}$$

is the 2-group of narrow logarithmic divisor classes (of degree 0).

Due to the weak approximation theorem, every class in $\widetilde{\mathcal{C}}\ell_F^{\text{res}}$ can be represented by a pair $(\alpha, \mathbf{1})$ where the vector $\mathbf{1}$ has all entries 1. So the canonical map $\alpha \mapsto (\alpha, \mathbf{1})$ induces a morphism from $\widetilde{\mathcal{D}}\ell_F$ onto $\widetilde{\mathcal{C}}\ell_F^{\text{res}}$, the kernel of which is $\widetilde{\mathcal{P}}\ell_F^+$. We conclude as expected:

$$\widetilde{\mathcal{C}}\ell_F^{\text{res}} = \widetilde{\mathcal{D}}\ell_F^{\text{res}} / \widetilde{\mathcal{P}}\ell_F^{\text{res}} \simeq \widetilde{\mathcal{D}}\ell_F / \widetilde{\mathcal{P}}\ell_F^+.$$

We are now in a situation to present an algorithm for computing narrow logarithmic classes. It uses our previous results by Diaz y Diaz et al. (2005) on (ordinary) logarithmic classes and mimics the classical feature concerning narrow and ordinary ideal classes. We note that this algorithm is a bit

If we can find generators of the 2-group of logarithmic units (see Section 4) and therefore relations (2), the columns of the following matrix $R \in \mathbb{Z}_2^{(v+r) \times (v+2r+c+1)}$ generate all relations for the $(a_j, 1)$ and the $(0, e_j)$:

$$R = \left(\begin{array}{cccc|cccc} 2^{n_1} & & & & 0 & \cdots & 0 & & 0 & \cdots & 0 & 0 \\ & \ddots & & & \vdots & & \vdots & & \vdots & & \vdots & \vdots \\ & & 2^{n_v} & & 0 & \cdots & 0 & & 0 & \cdots & 0 & 0 \\ \hline & & & & 2 & & & & b_{1,1} & \cdots & b_{1,r+c} & 1 \\ & c_{i,j} & & & & \ddots & & & \vdots & & \vdots & \vdots \\ & & & & & & 2 & & b_{v,1} & \cdots & b_{v,r+c} & 1 \end{array} \right).$$

4. Generators for the 2-group of logarithmic units

In the following we describe how generators of the 2-group of logarithmic units can be computed. The 2-group of logarithmic units

$$\tilde{\mathcal{E}}_F = \{x \in \mathcal{R}_F \mid \forall p : \tilde{v}_p(x) = 0\} = \{x \in \mathcal{E}'_F \mid \forall p \mid 2 : \tilde{v}_p(x) = 0\}$$

is a subgroup of the 2-group of 2-units \mathcal{E}'_F . If we assume that there are exactly d places p_1, \dots, p_d containing 2 in F , we have, say,

$$\mathcal{E}'_F = \{\pm 1\} \times \langle \varepsilon_1, \dots, \varepsilon_{r+c+d-1} \rangle.$$

We fix a precision e for our 2-adic approximations by requiring for elements ε of \mathcal{E}'_F the relation

$$\tilde{v}_{p_i}(\varepsilon) \equiv 0 \pmod{2^e} \quad (1 \leq i \leq d).$$

We obtain a system of generators of $\tilde{\mathcal{E}}_F$ by computing the nullspace of the matrix

$$M = \left(\begin{array}{ccc|ccc} & & & 2^e & \cdots & 0 \\ \tilde{v}_{p_i}(\varepsilon_j) & & & \vdots & \ddots & \vdots \\ & & & 0 & \cdots & 2^e \end{array} \right)$$

with $r + c + 2d - 1$ columns and d rows. We assume that the nullspace is generated by the columns of the matrix

$$M' = \left(\begin{array}{cc} C & \\ - & D \end{array} \right)$$

where C has $r + c + d - 1$ and D exactly d rows. It suffices to consider C . Each column $(n_1, \dots, n_{r+c+d-1})^{tr}$ of the matrix C corresponds to a unit

$$\prod_{i=1}^{r+c+d-1} \varepsilon_i^{n_i} \in \tilde{\mathcal{E}}_F \mathcal{E}'_F 2^e$$

so that we can choose

$$\tilde{\varepsilon} := \prod_{i=1}^{r+c+d-1} \varepsilon_i^{n_i}$$

as an approximation for a logarithmic unit. This procedure yields $k \geq r + c$ logarithmic units. If the integer k which we get in our calculations is not much larger than $r + c$ then we will proceed with the k generating elements of $\tilde{\mathcal{E}}_F$ obtained. Otherwise, we reduce the number of generators by computing a basis of the submodule of $\mathbb{Z}^{r+c+d-1}$ which is the span of the columns of C . Of course, by the generalized Gross conjecture we would have exactly $r + c$ such units. This allows us to assert that the precision e , chosen earlier in this section, was sufficient.

5. Applications in K-theory

We adopt the notations and definitions in this section from Jaulent and Soriano-Gafiuk (2004). In particular i denotes a primitive fourth root of unity; and we say that the number field F is *exceptional* when i is not contained in the cyclotomic \mathbb{Z}_2 -extension F^c of F , i.e. whenever the cyclotomic extension $F^c[i]/F$ is not procyclic.

We say that a non-complex place \mathfrak{p} of a number field F is *signed* whenever the local field $F_{\mathfrak{p}}$ does not contain the fourth root of unity i . These are the places which do not decompose in the extension $F[i]/F$. For such a place \mathfrak{p} , there exists a non-trivial sign-map

$$\text{sign}_{\mathfrak{p}} : F_{\mathfrak{p}}^{\times} \rightarrow \{\pm 1\},$$

given by the Artin reciprocity map $F_{\mathfrak{p}}^{\times} \rightarrow \text{Gal}(F_{\mathfrak{p}}[i]/F_{\mathfrak{p}})$ of class field theory.

We say that a non-complex place \mathfrak{p} of F is *logarithmically signed* if and only if one has $i \notin F_{\mathfrak{p}}^c$. These are the places which do not decompose in $F^c[i]/F^c$. So the finite set PS_F of logarithmic signed places of the field F only contains:

- (i) the subset PR_F of infinite real places and
- (ii) the subset PE_F of exceptional dyadic places, i.e. the set of logarithmic signed places above the prime 2.

We say that a non-complex place \mathfrak{p} of F is *logarithmically primitive* if and only if \mathfrak{p} does not decompose in the first step E/F of the cyclotomic \mathbb{Z}_2 -extension F^c/F . Finally we say that an exceptional number field F is *primitive* whenever there exists an exceptional dyadic place which is logarithmically primitive.

Naturally, the task arises to determine logarithmically signed places, i.e. those non-complex places of F for which i is not contained in $F_{\mathfrak{p}}^c$:

Proposition 5.1. *Let $E_{\mathfrak{p}}$ be the first quadratic extension of $F_{\mathfrak{p}}$ in the tower of field extension from $F_{\mathfrak{p}}$ to $F_{\mathfrak{p}}^c$. Then $i \in F_{\mathfrak{p}}^c$ holds precisely for $i \in E_{\mathfrak{p}}$.*

Proof. Since $F_{\mathfrak{p}}^c/F_{\mathfrak{p}}$ is a \mathbb{Z}_2 -extension, it contains exactly one quadratic extension $E_{\mathfrak{p}}$ of $F_{\mathfrak{p}}$. So we immediately obtain: $i \in F_{\mathfrak{p}}^c \Leftrightarrow F_{\mathfrak{p}}(i) \subseteq E_{\mathfrak{p}} \Leftrightarrow i \in E_{\mathfrak{p}}$. \square

Remark 5.2. The extension $E_{\mathfrak{p}}$ is $F_{\mathfrak{p}}(\alpha_k)$ where k is the smallest integer such that α_k does not belong to $F_{\mathfrak{p}}$ with $\alpha_0 = 0$ and $\alpha_{k+1} = \sqrt{2} + \alpha_k$.

We assume in the following that the number field F has no exceptional dyadic place. Let us introduce the group $\mathcal{C}\ell_F^{\text{res}}$ of narrow logarithmic classes without any assumption of degree:

$$\mathcal{C}\ell_F^{\text{res}} = \mathcal{D}\ell_F^{\text{res}} / \widetilde{\mathcal{P}}\ell_F^{\text{res}}.$$

Via the degree map, we obtain the direct decomposition:

$$\mathcal{C}\ell_F^{\text{res}} \simeq \mathbb{Z}_2 \oplus \widetilde{\mathcal{C}}\ell_F^{\text{res}},$$

where the torsion subgroup $\widetilde{\mathcal{C}}\ell_F^{\text{res}}$ was already computed in the previous section. So the quotient of exponent 2

$${}^2\mathcal{C}\ell_F^{\text{res}} := \mathcal{C}\ell_F^{\text{res}} / (\mathcal{C}\ell_F^{\text{res}})^2$$

contains both as hyperplanes the two quotients, ${}^2\widetilde{\mathcal{C}}\ell_F^{\text{res}}$ relative to $\widetilde{\mathcal{C}}\ell_F^{\text{res}}$ and ${}^2\mathcal{C}\ell_F^{\text{pos}}$ relative to the positive class group $\mathcal{C}\ell_F^{\text{pos}}$ introduced by Jaulent and Soriano-Gafiuk (2004).

Since, according to Jaulent and Soriano-Gafiuk (2004), this gives the 2-rank of the wild kernel $\text{WK}_2(F)$, we can extend the results of Hutchinson (2001, 2004) as follows:

Theorem 5.3. Let F be a number field which has no exceptional dyadic places.

- (i) If F is not exceptional (i.e. in case $i \in F^c$) the wild kernel $WK_2(F)$ coincides with the subgroup $K_2^\infty(F) = \bigcap_{n \geq 1} K_2^n(F)$ of infinite height elements in $K_2(F)$; the group $\widetilde{\mathcal{C}}\ell_F^{\text{res}}$ of narrow logarithmic classes coincides with the group $\widetilde{\mathcal{C}}\ell_F$ of (ordinary) logarithmic classes; and one has immediately:

$$\text{rk}_2 WK_2(F) = \text{rk}_2 K_2^\infty(F) = \text{rk}_2 \widetilde{\mathcal{C}}\ell_F^{\text{res}} = \text{rk}_2 \widetilde{\mathcal{C}}\ell_F.$$

- (ii) If F is exceptional (i.e. in case $i \notin F^c$) the subgroup $K_2^\infty(F)$ has index 2 in the wild kernel $WK_2(F)$ and one still has:

$$\text{rk}_2 WK_2(F) = \text{rk}_2 \widetilde{\mathcal{C}}\ell_F^{\text{res}} \geq 1.$$

- (ii,a) In case $WK_2(F)$ and $K_2^\infty(F)$ have the same 2-rank, this gives:

$$\text{rk}_2 K_2^\infty(F) = \text{rk}_2 \widetilde{\mathcal{C}}\ell_F^{\text{res}} \geq 1.$$

- (ii,b) Also, in case $K_2^\infty(F)$ is a direct summand in $WK_2(F)$, one has:

$$\text{rk}_2 K_2^\infty(F) = \text{rk}_2 \widetilde{\mathcal{C}}\ell_F^{\text{res}} - 1.$$

Proof. In the non-exceptional case, the number field F is not locally exceptional, i.e. has no logarithmic signed places: $PE_F = PR_F = \emptyset$. In particular, narrow logarithmic classes coincide with ordinary logarithmic classes and the result follows from (Jaulent and Soriano-Gafiuk, 2004).

In the exceptional case, the number field F may have real places, so the narrow logarithmic class group $\widetilde{\mathcal{C}}\ell_F^{\text{res}}$ may differ from the ordinary logarithmic class group. Moreover, because of the assumption $PE_F = \emptyset$ and the results of Hutchinson (Hutchinson, 2001, 2004), the subgroup $K_2^\infty(F)$ has index 2 in the wild kernel $WK_2(F)$. \square

Remark 5.4. It remains to determine whether a number field F is not exceptional, i.e. whether the cyclotomic \mathbb{Z}_2 -extension F^c contains the fourth root of unity i . Of course if $i \in F^c$ then i is contained in the quadratic subfield E/F in F^c . Now the finite subfields of \mathbb{Q}^c are the real cyclic fields $\mathbb{Q}^{(s)} = \mathbb{Q}[\zeta_{2^{s+2}} + \zeta_{2^{s+2}}^{-1}]$ and the finite extensions of F^c are of the form $F\mathbb{Q}^{(s)}$. So we only need to check whether i is contained in $F[\zeta_{2^{s+2}} + \zeta_{2^{s+2}}^{-1}]$ where s is minimal with $\zeta_{2^{s+2}} + \zeta_{2^{s+2}}^{-1} \notin F$.

6. Examples

The methods described here were implemented in the computer algebra system Magma (Cannon et al., 2006). Many of the fields used in the examples were results of queries to the QaoS number field database (Daberkow and Weber, 1996; Freundt et al., 2006).

The wild kernel $WK_2(F)$ of a number field F is contained in the tame kernel $K_2(\mathcal{O}_F)$. Let $\mu(F)$ be the order of the torsion subgroup of F^\times and for a prime \mathfrak{p} of F over p denote by $\mu^1(F_{\mathfrak{p}})$ the p -Sylow subgroup of the torsion subgroup of $F_{\mathfrak{p}}^\times$. By coupling Moore’s exact sequence and the localization sequence (Gras, 1986, section 1) one obtains the index formula (Belabas and Gangl, 2004, equation (6)):

$$(K_2(\mathcal{O}_F) : WK_2(F)) = \frac{2^r}{|\mu(F)|} \prod_{\mathfrak{p}} |\mu^1(F_{\mathfrak{p}})|,$$

where \mathfrak{p} runs through all finite places and r is the number of real places of F . We apply this in the determination of the structure of $WK_2(F)$ in the cases where the structure of $K_2(\mathcal{O}_F)$ is known.

In the tables Abelian groups are given as a list of the orders of their cyclic factors. Furthermore we use the following notation:

- f is an irreducible polynomial over the integers;
- F denotes the number field generated by a root of f ;
- r is the number of real places of F ;
- d_F denotes the discriminant for a number field F ;

- Gal denotes the Galois group of f ;
- $K_2(O_F)$ denotes the structure of the tame kernel of F ;
- $[\cdot]$ denotes the index $(K_2(O_F) : WK_2(F))$;
- $\mathcal{C}\ell_F$ denotes the class group, P the set of dyadic places;
- $\mathcal{C}\ell'_F$ denotes the 2-part of $Cl/\langle P \rangle$;
- $\widetilde{\mathcal{C}\ell}_F$ denotes the logarithmic class group;
- $\mathcal{C}\ell_F^{res}$ denotes the group of narrow logarithmic classes;
- rk_2 denotes the 2-rank of the wild kernel WK_2 ;
- WK_2 denotes the wild kernel in $K_2(F)$;
- K_2^∞ denotes the subgroup of infinite height elements in $K_2(F)$.

Imaginary quadratic fields. Belabas and Gangl (2004) have developed an algorithm for the computation of the tame kernel $K_2\mathcal{O}_F$. The following table contains the structure of $K_2\mathcal{O}_F$ as computed by Belabas and Gangl and the 2-rank of the wild kernel $WK_2(F)$ calculated with our methods. We also give the structure of the wild kernel if it can be deduced from the structure of $K_2\mathcal{O}_F$ and of the rank of the wild kernel computed here or in (Pauli and Soriano-Gafiuk, 2004). The structure of the tame kernel $K_2(\mathcal{O}_F)$ of all fields except for the starred entries has been proven by Belabas and Gangl. The table gives the structure of the wild kernel of all imaginary quadratic fields F with no exceptional places and discriminant $|d_F| < 1000$.

d_F	$\mathcal{C}\ell_F$	$K_2(O_F)$	$[\cdot]$	$\mathcal{C}\ell'_F$	$\widetilde{\mathcal{C}\ell}_F$	$\widetilde{\mathcal{C}\ell}_F^{res}$	rk_2	WK_2	K_2^∞
-68	[4]	[8]	2	[2]	[2]	[2]	1	[4]	[2]
-132	[2,2]	[4]	2	[2]	[2]	[2]	1	[2]	[]
-136	[4]	[4]	1	[2]	[2]	[2]	1	[4]	[2]
-164	[8]	[4]	2	[4]	[4]	[4]	1	[2]	[]
-228	[2,2]	[12]	6	[2]	[2]	[2]	1	[2]	[]
-260	[2,4]	[4]	2	[4]	[4]	[4]	1	[2]	[]
-264	[2,4]	[6]	3	[4]	[4]	[4]	1	[2]	[]
-292	[4]	[4]	2	[2]	[2]	[2]	1	[2]	[]
-328	[4]	[2]	1	[2]	[2]	[2]	1	[2]	[]
-356	[12]	[4]	2	[2]	[2]	[2]	1	[2]	[]
-388	[4]	[8]	2	[2]	[2]	[2]	1	[4]	[2]
-420	[2,2,2]	[2,4]	2	[2,2]	[2,2]	[2,2]	2	[2,2]	[2]
-452	[8]	[8]	2	[4]	[4]	[4]	1	[4]	[2]
-456	[2,4]	[2]	1	[4]	[4]	[4]	1	[2]	[]
-516	[2,6]	[12]	6	[2]	[2]	[2]	1	[2]	[]
-520	[2,2]	[2]	1	[2]	[2]	[2]	1	[2]	[]
-548	[8]	[4]	2	[4]	[4]	[4]	1	[2]	[]
-580	[2,4]	[4]	2	[4]	[4]	[4]	1	[2]	[]
-584	[16]	[2]	1	[8]	[8]	[8]	1	[2]	[]
-644	[2,8]	[2,16]	2	[2,4]	[2,4]	[2,4]	2	[2,8]	[?]
-708	[2,2]	[4]	2	[2]	[2]	[2]	1	[2]	[]
-712	[8]	[2]	1	[4]	[4]	[4]	1	[2]	[]
-740	[2,8]	[4]	2	[8]	[8]	[8]	1	[2]	[]
-772	[4]	[8]	2	[2]	[2]	[2]	1	[4]	[2]
-776	[20]	[4]	1	[2]	[2]	[2]	1	[4]	[2]
*-804	[2,6]	[36]	6	[2]	[2]	[2]	1	[6]	[3]
-836	[2,10]	[4]	2	[2]	[2]	[2]	1	[2]	[]
-840	[2,2,2]	[2,6]	3	[2,2]	[2,2]	[2,2]	2	[2,2]	[2]
-868	[2,4]	[2,4]	2	[2,2]	[2,2]	[2,2]	2	[2,2]	[2]
-904	[8]	[4]	1	[4]	[4]	[4]	2	[4]	[2]
-964	[12]	[8]	2	[2]	[2]	[2]	1	[4]	[2]
-996	[2,6]	[4]	2	[2]	[2]	[2]	1	[2]	[]

Real quadratic fields. The table contains all real quadratic fields F with no exceptional places and discriminant $|d_F| < 1000$. All these fields are exceptional.

d_F	$\mathcal{C}l_F$	$[\cdot]$	$ P $	$ PE $	$\mathcal{C}l'_F$	$\tilde{\mathcal{C}}l_F$	$\tilde{\mathcal{C}}l_F^{res}$	rk_2
28	[]	8	1	0	[]	[]	[2]	1
56	[]	4	1	0	[]	[]	[2]	1
60	[2]	24	1	0	[]	[]	[2]	1
92	[]	8	1	0	[]	[]	[2]	1
120	[2]	4	1	0	[]	[]	[2]	1
124	[]	8	1	0	[]	[]	[2]	1
156	[2]	8	1	0	[]	[]	[2]	1
184	[]	4	1	0	[]	[]	[2]	1
188	[]	8	1	0	[]	[]	[2]	1
220	[2]	8	1	0	[]	[]	[2]	1
248	[]	4	1	0	[]	[]	[2]	1
284	[]	8	1	0	[]	[]	[2]	1
312	[2]	12	1	0	[]	[]	[2]	1
316	[3]	8	1	0	[]	[]	[2]	1
348	[2]	24	1	0	[]	[]	[2]	1
376	[]	4	1	0	[]	[]	[2]	1
380	[2]	8	1	0	[]	[]	[2]	1
412	[]	8	1	0	[]	[]	[2]	1
440	[2]	4	1	0	[]	[]	[2]	1
444	[2]	8	1	0	[]	[]	[2]	1
476	[2]	8	1	0	[2]	[2]	[2, 2]	2
604	[]	8	1	0	[]	[]	[2]	1
632	[]	4	1	0	[]	[]	[2]	1
636	[2]	24	1	0	[]	[]	[2]	1
668	[]	8	1	0	[]	[]	[2]	1
696	[2]	4	1	0	[]	[]	[2]	1
732	[2]	8	1	0	[]	[]	[2]	1
760	[2]	4	1	0	[]	[]	[2]	1
764	[]	8	1	0	[]	[]	[2]	1
796	[]	8	1	0	[]	[]	[2]	1
824	[]	4	1	0	[]	[]	[2]	1
860	[2]	8	1	0	[]	[]	[2]	1
888	[2]	12	1	0	[]	[]	[2]	1
892	[3]	8	1	0	[]	[]	[2]	1
924	[2, 2]	24	1	0	[2]	[2]	[2, 2]	2
952	[2]	4	1	0	[2]	[2]	[2, 2]	2
956	[]	8	1	0	[]	[]	[2]	1
988	[2]	8	1	0	[]	[]	[2]	1

Real quadratic fields with class numbers 32 and 64. The table contains extensions with class number 32 up to discriminant 222 780 and extensions with class number 64 up to discriminant 805 596.

d_F	$\mathcal{C}l_F$	$[\cdot]$	$ P $	$ PE $	$\mathcal{C}l'_F$	$\widetilde{\mathcal{C}l}_F$	$\widetilde{\mathcal{C}l}_F^{res}$	rk_2
112 924	[2,16]	8	1	0	[16]	[16]	[2,16]	2
120 796	[2,16]	8	1	0	[16]	[16]	[2,16]	2
136 120	[2,16]	4	1	0	[16]	[16]	[2,16]	2
153 660	[2,2,8]	8	1	0	[2,8]	[2,8]	[2,2,8]	3
158 844	[2,2,8]	8	1	0	[2,8]	[2,8]	[2,2,8]	3
163 576	[2,16]	4	1	0	[2,8]	[2,8]	[2,2,8]	3
170 872	[2,16]	4	1	0	[16]	[16]	[2,16]	2
176 316	[2,16]	24	1	0	[16]	[16]	[2,16]	2
176 440	[2,16]	4	1	0	[16]	[16]	[2,16]	2
196 540	[2,16]	8	1	0	[16]	[16]	[2,16]	2
202 524	[2,16]	24	1	0	[16]	[16]	[2,16]	2
207 480	[2,2,2,4]	4	1	0	[2,2,4]	[2,2,4]	[2,2,2,4]	4
213 180	[2,2,2,4]	24	1	0	[2,2,4]	[2,2,4]	[2,2,2,4]	4
221 276	[2,16]	8	1	0	[16]	[16]	[2,16]	2
222 780	[2,2,8]	8	1	0	[2,8]	[2,8]	[2,2,8]	3
374 136	[2,2,16]	12	1	0	[2,16]	[2,16]	[2,2,16]	3
382 204	[2,32]	8	1	0	[32]	[32]	[2,32]	2
449 436	[2,2,16]	8	1	0	[2,16]	[2,16]	[2,2,16]	3
484 764	[2,2,16]	24	1	0	[2,16]	[2,16]	[2,2,16]	3
506 940	[2,2,2,8]	24	1	0	[2,2,8]	[2,2,8]	[2,2,2,8]	4
805 596	[2,2,16]	24	1	0	[2,16]	[2,16]	[2,2,16]	3

Biquadratic extensions. The table contains quadratic and biquadratic number fields. The biquadratic fields are the compositum of the first quadratic extensions and one of the other quadratic extensions. All fields are exceptional.

F	d_F	r	$\mathcal{C}l_F$	$[\cdot]$	$ P $	$\mathcal{C}l'_F$	$\widetilde{\mathcal{C}l}_F$	$\widetilde{\mathcal{C}l}_F^{res}$	rk_2
K	9 660	2	[2,2,2]	8	1	[2,2]	[2,2]	[2,2,2]	3
L_1	9 340	2	[10]	8	1	[]	[]	[2]	1
KL_1		4	[2,2,2,10]	128	2	[2,2,2]	[2,2,2]	[2,2,2,2,2]	5
L_2	13 020	2	[2,2,2]	24	1	[2,2]	[2,2]	[2,2,2]	3
KL_2		4	[2,2,4]	384	2	[2,2]	[2,2,2]	[2,2,2,2,2,2]	6
L_3	15 708	2	[2,2,2]	8	1	[2,2]	[2,2]	[2,2,2]	3
KL_3		4	[2,2,4,28]	128	2	[2,2,4]	[2,2,4]	[2,2,2,2,2,4]	6

References

- Belabas, K., Gangl, H., 2004. Generators and relations for $K_2 O_F$. *K-Theory* 31, 195–231.
- Cannon, J.J., et al. 2006. The computer algebra system Magma, The University of Sydney.
- Daberkow, M., Weber, A., 1996. A database for number fields. In: Jacques, Calmet, Carla, Limongelli (Eds.), *Design and Implementation of Symbolic Computation Systems – DISCO'96*. In: *Lecture Notes in Computer Science*, vol. 1128. Springer-Verlag, pp. 320–330.
- Diaz y Diaz, F., Soriano, F., 1999. Approche algorithmique du groupe des classes logarithmiques. *J. Number Theory* 76, 1–15.
- Diaz y Diaz, F., Jaulent, J.-F., Pauli, S., Pohst, M.E., Soriano, F., 2005. A new algorithm for the computation of logarithmic class groups of number fields. *Experiment. Math.* 14, 67–76.
- Freundt, S., Karve, A., Krahmhann, A., Pauli, S., 2006. KASH: Recent developments. In: *Mathematical Software - ICMS 2006, Second International Congress on Mathematical Software*. In: LNCS, vol. 4151. Springer, Berlin.
- Gras, G., 1986. Remarks on K_2 of number fields. *J. Number Theory* 23 (3), 322–335.
- Hutchinson, K., 2001. The 2-Sylow subgroup of the wild kernel of exceptional number fields. *J. Number Theory* 87, 222–238.
- Hutchinson, K., 2004. On tame and wild kernels of special number fields. *J. Number Theory* 107, 222–238.
- Jaulent, J.-F., 1994a. Sur le noyau sauvage des corps de nombres. *Acta Arith.* 67, 335–348.
- Jaulent, J.-F., 1994b. Classes logarithmiques des corps de nombres. *J. Théor. Nombres Bordeaux* 6, 301–325.
- Jaulent, J.-F., Soriano-Gafiuk, F., 2001. Sur le noyau sauvage des corps de nombres et le groupe des classes logarithmiques. *Math. Z.* 238, 335–354.
- Jaulent, J.-F., Soriano-Gafiuk, F., 2004. 2-groupe des classes positives d'un corps de nombres et noyau sauvage de la K-théorie. *J. Number Theory* 108, 187–208.
- Jaulent, J.-F., Michel, A., 2006. Approche algorithmique des noyaux étales sauvages des corps de nombres. *J. Number Theory* 120, 72–91.
- Pauli, S., Soriano-Gafiuk, F., 2004. The discrete logarithm in logarithmic ℓ -class groups and its applications in K-Theory. In: *Algorithmic Number Theory*. In: *Lecture Notes in Comput. Sci.*, vol. 3076. Springer, pp. 367–378.
- Soriano, F., 1997. Classes logarithmiques au sens restreint. *Manuscripta Math.* 93, 409–420.
- Soriano-Gafiuk, F., 2000. Sur le noyau hilbertien d'un corps de nombres. *C. R. Acad. Sci. Paris* 330 (Série I), 863–866.