

Polo-like Kinase-1 Controls Proteasome-Dependent Degradation of Claspin during Checkpoint Recovery

Ivan Mamely,^{1,5} Marcel ATM van Vugt,^{2,5,7} Veronique AJ Smits,³ Jennifer I. Semple,¹ Bennie Lemmens,² Anastassis Perrakis,⁴ René H. Medema,^{2,6,*} and Raimundo Freire^{1,6}

¹Unidad de Investigación
Hospital Universitario de Canarias
Ofra s/n, La Cuesta
38320 Tenerife
Spain

²Department of Medical Oncology
University Medical Centre Utrecht
Heidelberglaan 100
3584CG Utrecht
The Netherlands

³Department of Cell Biology and Genetics
Erasmus MC
P.O. Box 1738
3000 DR Rotterdam
The Netherlands

⁴Division of Molecular Carcinogenesis
Netherlands Cancer Institute
Plesmanlaan 121
1066 CX Amsterdam
The Netherlands

Summary

DNA-damage checkpoints maintain genomic integrity by mediating a cell-cycle delay in response to genotoxic stress or stalled replication forks. In response to damage, the checkpoint kinase ATR phosphorylates and activates its effector kinase Chk1 in a process that critically depends on Claspin [1]. However, it is not known how exactly this kinase cascade is silenced. Here we demonstrate that the abundance of Claspin is regulated through proteasomal degradation. In response to DNA damage, Claspin is transiently stabilized, and its expression depends on Chk1 kinase activity. In addition, we show that Claspin is degraded upon mitotic entry, a process that depends on the β -TrCP-SCF ubiquitin ligase and Polo-like kinase-1 (Plk1). We demonstrate that Claspin interacts with both β -TrCP and Plk1 and that inactivation of these components or the β -TrCP recognition motif in Claspin prevents its mitotic degradation. Interestingly, expression of a nondegradable Claspin mutant inhibits recovery from a DNA-damage-induced checkpoint arrest. Thus, we conclude that Claspin levels are tightly regulated, both during unperturbed cell cycles and after

DNA damage. Moreover, our data demonstrate that the degradation of Claspin at the onset of mitosis is an essential step for the recovery of a cell from a DNA-damage-induced cell-cycle arrest.

Results and Discussion

Claspin Turnover Is Regulated during the Cell Cycle and in Response to DNA Damage

Activation of Chk1 by the DNA-damage-responsive checkpoint kinase ATR critically depends on the presence of Claspin, which acts as an adaptor protein to link Chk1 and ATR [1, 2]. Interestingly, Claspin levels were shown to oscillate during the cell cycle, suggesting that the checkpoint response is cell-cycle dependent [2]. To investigate this dependence, we analyzed Claspin levels at different stages during the cell cycle (Figure 1A). Claspin levels are high during S and early G2 phase, whereas Claspin levels sharply decrease as cells accumulate in the G2 and M phases (Figure 1A). Indeed, we could show by immunofluorescence that Claspin is virtually absent in mitotic cells (Figure 1B). We further confirmed downregulation of Claspin in mitotic cells by using the microtubule-destabilizing drug nocodazole, which blocks cell-cycle progression in prometaphase of mitosis (Figure 1C). Moreover, Claspin levels rapidly increased upon release from the nocodazole block, at time points when cells progressively entered G1 and S phase (Figure S1A in the Supplemental Data available online). Finally, cells arrested in G2 after treatment with the DNA-damaging agent camptothecin displayed high levels of Claspin (Figure S1B). These results indicate that Claspin is stabilized in cells that are prevented from entering mitosis and suggest that Claspin is degraded at mitotic onset, possibly precluding further activation of Chk1 by ATR.

We also analyzed the levels of Claspin after different genotoxic insults. We observed an increase in Claspin levels shortly after UV treatment and a decrease at later time points (Figure 1D). These fluctuations in Claspin levels were also observed with other types of DNA-damaging agents, including hydroxyurea (causing stalled replication forks), etoposide, and camptothecin (the last two causing both single- and double-strand DNA breaks) (Figure 1D and data not shown). Thus, in addition to the observed decrease of Claspin at mitotic entry, we also found upregulation of Claspin levels in response to genotoxic stress.

The rapid increase in Claspin levels after DNA damage suggested that this could not be due to transcriptional regulation, and we tested whether altered proteasomal degradation is involved. Indeed, treating U2OS cells with the proteasome inhibitor MG132 resulted in an increase of Claspin levels, up to the maximal level of expression that is seen in response to DNA damage (Figures 2A and 2B). Interestingly, addition of MG132 to damaged cells could not elicit a further increase in Claspin expression (Figure 2B) but did prevent the

*Correspondence: r.h.medema@med.uu.nl

⁵These authors contributed equally to this work.

⁶These authors contributed equally to this work.

⁷Present Address: Center for Cancer Research, Massachusetts Institute of Technology, 77 Massachusetts Ave., Cambridge, Massachusetts 021329.




Figure 1. Claspin Regulation during the Cell Cycle and after DNA Damage

(A) U2OS cells were synchronized at the G1/S transition by double thymidine block, and cells were collected at different times after release for flow cytometry and western blot analysis.
 (B) U2OS cells were analyzed by immunofluorescence using DAPI, anti-Claspin and anti-phospho-HistoneH3 as a mitotic marker. Arrowhead points at a mitotic cell.
 (C) U2OS cells were left untreated or arrested with nocodazole for 18h and whole cell extracts (WCE) were prepared for analysis of Claspin expression. Cyclin E was used as an S-phase marker.
 (D) U2OS cells were treated with UV light (40 J/m²) or with 0.5 μM camptothecin (CPT) for 1 hr. Cells were collected at the indicated time points for analysis of Claspin expression.

reduction normally seen in Claspin levels at the later stages of the damage response (data not shown). Moreover, we found that the reduction seen for Claspin expression in mitotic cells was blocked by MG132, indicating that this reduction is due to enhanced processing of Claspin by the proteasome (Figure 2C). Altogether, these data demonstrate that Claspin is continuously turned over by the proteasome in undamaged cells, that it becomes transiently stabilized in response to DNA damage, and that its turnover is even further enhanced at mitotic entry. Consistent with this, we could detect a smear of higher molecular weight than Claspin in cells cotransfected with HA-tagged Claspin and His-tagged ubiquitin, indicating that Claspin is indeed poly-ubiquitinated (Figure 2D). Furthermore, we could demonstrate poly-ubiquitination of endogenous Claspin in cells treated with proteasome inhibitors (Figure 2E). Notably, Claspin ubiquitination was significantly enhanced in mitotic cells (Figure 2E), whereas overall levels of HA-Claspin were considerably lower in mitotic cells than in the asynchronous cultures, consistent with the notion that Claspin turnover is increased upon entry into mitosis.

Chk1 and Polo-like Kinase-1 Exert antagonistic effects on Claspin turnover

Several examples of regulated proteasomal degradation by means of substrate phosphorylation exist in the

literature, and because both Chk1 and Plx1 are known to interact with Claspin [1, 3], we tested whether Chk1 and Polo-like kinase-1 (Plk1) can affect Claspin turnover. Indeed, whereas expression of wild-type Chk1 did not destabilize Claspin, overexpression of a kinase-dead version of Chk1 resulted in destabilization of Claspin protein levels (Figure 3A). Consistent with these observations, downregulation of Chk1 by siRNA also resulted in a decrease in Claspin levels (Figure 3B). These data confirm recent findings by Chini et al. [4] but furthermore show that this Chk1-mediated stabilization of Claspin requires Chk1 kinase activity.

We next set out to determine whether Plk1, the human Plx1 homolog, was also able to modulate Claspin levels. Plk1 is expressed in G2 and mitosis but only becomes activated as cells enter mitosis, which is also when Claspin levels drop. Indeed, we could show that overexpression of Plk1 resulted in a downregulation of Claspin (Figure 3C). This effect depends on the catalytic activity of Plk1; expression of a kinase-defective mutant of Plk1 could not cause downregulation of Claspin (Figure 3C). In addition, by using GST fusions comprising different regions of Claspin [2], we were able to show that Plk1 can interact with the C-terminal half of Claspin (Figure S3A), consistent with findings of Yoo et al. in *Xenopus* [3]. We were able to confirm this interaction further by coimmunoprecipitation of Myc-tagged kinase-dead Plk1 with endogenous Claspin in vivo and vice




Figure 2. Claspín Protein Levels Are Regulated via Proteasome-Dependent Degradation

(A) U2OS cells were treated with MG132 for the indicated times before Western-blot analysis with the indicated antibodies. (B) U2OS cells were incubated with MG132 for 3 hr before treatment with UV light (40 J/m²). Cells were collected 1 hr after UV treatment for analysis of Claspín expression. (C) Claspín expression in U2OS cells treated with nocodazole for 8 hr in the presence or absence of MG132. Untreated cells were taken along as a control. (D) HEK 293T cells were transfected with indicated plasmids. After 36 hr, cells were lysed and extracts were incubated with Ni-NTA beads. After washing, the resin was analyzed for Claspín ubiquitination. (E) U2OS cells were incubated for 3 hr in the presence of MG132 before lysis. Extracts were used for immunoprecipitations with Claspín or control IgG antibodies. Analysis was done by Western blotting with antibodies against Claspín and ubiquitin.

versa (Figure 3D and data not shown). This interaction of Plk1 with Claspín is mediated through the conserved Polo-box domain (PBD), as demonstrated by pull-down experiments with a GST fusion of the PBD of Plk1 (Figure 3E).

Using RNA-interference to deplete Plk1, we assessed whether Plk1 is required for the degradation of Claspín at mitotic onset. Indeed, depletion of Plk1 in HeLa and U2OS cells was able to prevent the degradation of Claspín in mitosis (Figure 3F), indicating that Plk1 is essential for mitotic Claspín degradation. Interestingly, whereas control-transfected mitotic cells show loss of Claspín during mitosis, no clear reduction in Claspín levels was observed in Plk1-depleted mitotic cells (Figure S3B).

β-TrCP Targets Claspín for Degradation

The SCF (Skp1-Cullin-F-box-protein) ubiquitin ligase complex is responsible for the degradation of multiple proteins during mitotic onset. In particular, key mitotic-entry regulators, such as Wee1 and Emi1 [5–7], are degraded by the β-TrCP-SCF complex in a Plk1-dependent fashion. To test whether Claspín might also be degraded by the β-TrCP-SCF complex, we downregulated β-TrCP1/2 by shRNA. Importantly, downregulation of β-TrCP1/2 resulted in elevation of Claspín protein levels during mitosis (Figure 4A), and we could demonstrate an interaction between Claspín, β-TrCP1/2 and Skp1 (Figure S4A and data not shown). Interestingly, a DSGxxS motif that is present in most SCF substrates [8] can be recognized in a conserved region in the N terminus of Claspín (Figure S4B). To investigate whether this DSGxxS motif is required for Claspín targeting for proteasomal degradation, we generated point mutants in which the two essential serines were replaced by alanines (Claspín S30/34A). Strikingly, the S30/34A mutant displayed a

reduced affinity for β-TrCP1/2 as compared to wt-Claspín (Figure 4B), and disruption of the recognition motif abolished Claspín degradation in mitotic cells (Figure 4C).

Together, these results show the significant role for the β-TrCP1/2-SCF complex in Claspín degradation at mitotic onset. Also, they suggest that Plk1 might act at multiple points to promote checkpoint recovery. We have previously demonstrated that Plk1 promotes proteasomal degradation of the Cdk-inhibitory kinase Wee1 [7], which acts at the bottom of the checkpoint signaling cascade. Here we show that Plk1 is also involved in promoting the degradation of Claspín. Similar to Wee1, Plk1 can promote the proteasomal degradation of Claspín in a β-TrCP1/2-dependent fashion. This degradation requires the presence of a phosphodegron motif (DSGxxS), which is also found in other β-TrCP1/2 substrates, such as Emi1 [5, 6]. Presently, we do not know whether Plk1 directly phosphorylates the serines in the phosphodegron motif or whether Plk1-dependent degradation of Claspín by the β-TrCP1/2-SCF complex occurs in a more complex fashion, as was shown for Wee1 [9].

Claspín Degradation Is Required for Checkpoint Recovery

The downregulation of Claspín levels at mitotic entry might act to prevent Chk1 activation during mitosis. Indeed, we found that Chk1 can be activated in mitotic cells expressing the nondegradable S30/34A mutant of Claspín (Figure 4D), indicating that the degradation of Claspín is required to inactivate the checkpoint in mitosis. Similarly, degradation of Claspín might be a way to decrease Chk1 activity during recovery from a checkpoint arrest. To test this latter hypothesis, we arrested U2OS cells in G2 with a DNA-damaging agent and then treated them with caffeine to induce checkpoint




Figure 3. Chk1 and Plk1 Both Regulate Claspin Levels

(A) HEK 293T cells were transfected with wild-type or kinase-dead versions of Flag-Chk1. WCE were analyzed by Western blot for Claspin expression.

(B) U2OS cells were transfected with control (Luciferase) or Chk1 siRNA for 72 hr. WCE were analyzed by Western blotting.

(C) 293T cells were transfected with Myc-tagged versions of wild-type or kinase-dead Plk1. After 36 hr, WCE were made and analyzed.

(D) Immunocomplexes isolated with control or Myc antibodies from U2OS cells expressing Myc-kd-Plk1 were analyzed by Western blotting for associated Claspin.

(E) U2OS cells were transfected with HA-Claspin and treated for 16 hr with taxol. Alternatively, cells were pretreated with doxorubicin (0.5 μ M) for 1 hr. WCE were subsequently incubated with recombinant GST-PBD. WCE and pull-down fractions were analyzed by Western blot.

(F) U2OS or HeLa cells were transfected with the pSuper or pSuper-Plk1 plasmids and subsequently arrested with thymidine for 24 hr. S-phase (2 hr after release) and mitotic shake-off fractions (18 hr after release, in the presence of nocodazole) were collected. WCE were analyzed by Western blot.

recovery. Caffeine inhibits the checkpoint kinases ATM and ATR and silences the checkpoint, allowing cells to recover from the arrest [10]. Indeed, whereas Chk1 phosphorylation is lost upon caffeine treatment in control cells, Plk1 depletion resulted in less-efficient Chk1 dephosphorylation after checkpoint silencing (Figure 4E). As expected, the status of Chk1 phosphorylation completely matched the presence or absence of Claspin (Figure 4E). To confirm that Claspin degradation is a key event in checkpoint recovery, we next assessed the effects of expressing the nondegradable Claspin mutant S30/34A. Expression of this mutant significantly lowered the extent of checkpoint recovery in response to different types of DNA damage. When caffeine was

used to silence the G2 DNA-damage checkpoint induced by doxorubicin or camptothecin, a clear decrease in checkpoint recovery was visible in cells expressing non-degradable Claspin (Figure 4F). These results indicate that degradation of Claspin indeed is an important requirement during checkpoint recovery induced by caffeine treatment. We next investigated spontaneous checkpoint recovery by using hydroxyurea. Eighteen hours after a 1 hr pulse with 10 mM hydroxyurea, most cells had recovered, as evidenced by a high mitotic index (Figure 4F). However, cells expressing the nondegradable Claspin displayed a significant decrease in checkpoint recovery (Figure 4F). Together, these results demonstrate that the Plk1-dependent process of Claspin




Figure 4. β-TrCP1/2-SCF-Dependent Degradation of Claspin Is Required for Checkpoint Recovery

(A) U2OS cells were transfected with pSuper-GFP or pSuper-β-TrCP1/2 in combination with pBabePuro. Twelve hours after transfection, cells were incubated with puromycin for 24 hr. Cells were then left untreated or incubated with nocodazole for 16 hr. Mitotic cells were collected by gentle shake-off, and WCE were prepared and analyzed by Western blot.

(B) HEK 293T cells were transfected with Flag-β-TrCP2 together with HA-wt-Claspin or HA-S30/34A-Claspin. Immunoprecipitations for Flag were analyzed by Western blot with HA and Flag antibodies.

(C) U2OS cells were transfected with HA-wt-Claspin or HA-S30/34A-Claspin. Cells were treated with nocodazole for 12 hr, then, mitotic cells were collected by gentle shake-off, and WCE were prepared and analyzed by Western blot.

(D) U2OS cells were left untransfected (left panel) or were transfected with either HA-wt-Claspin or HA-S30/34A-Claspin (right panel). Mitotic cells were obtained by treatment of the cells with nocodazole followed by mitotic shake off. Where indicated, cells were treated with UV light (20 J/m²), and WCE were made 90 min later. Claspin and HA antibodies were used for detection of endogenous and exogenous Claspin, respectively.

(E) U2OS cells were transfected with pSuper or pSuper-Plk1 and were synchronized with a 24 hr thymidine block. Eight hours after release, cells were treated for 1 hr with doxorubicin (0.5 μM) or left untreated. Subsequently, nocodazole was added. After 16 hr, all doxorubicin-treated cells were arrested in G₂. These cells were left untreated or were treated with 5 mM caffeine for 8 hr. WCE were prepared and analyzed by Western blotting with the indicated antibodies (left panel). In parallel, DNA content and MPM2 positivity was assessed by flow cytometry (right panel).

(F) U2OS cells were transfected with HA-wt-Claspin or HA-S30/34A-Claspin and treated as for Figure 4E. In addition, similar experiments were performed with camptothecin and hydroxyurea. Subsequently, cells were fixed and stained for MPM2. Relative mitotic indices are shown (mitotic indices of HA-wt-Claspin were put to 100%). Recovery after hydroxyurea was assessed in the absence of caffeine.

degradation is a key event in allowing recovery after an arrest by the DNA-damage checkpoint.

Taken together, our observations indicate that the degradation of Claspin serves to silence the DNA-damage checkpoint at mitotic entry. Moreover, we find that expression of nondegradable Claspin inhibits mitotic entry after checkpoint recovery. This effect is seen in cultures that have been arrested in G2 via a variety of DNA damaging agents and that are subsequently induced to enter mitosis by treatment with caffeine. In addition, the effect of nondegradable Claspin on recovery is even more pronounced in cultures that are allowed to recover spontaneously from a hydroxyurea-induced S-phase arrest, representing a physiological condition of checkpoint recovery. In contrast, expression of nondegradable Claspin has no effect on mitotic entry in an unperturbed cell cycle, indicating that Claspin degradation is not required per se for mitotic entry under all conditions. Importantly, expression of a nondegradable Claspin behaves in a similar way as Plk1 depletion, which we previously showed to lead to a stringent block in checkpoint recovery while failing to block mitotic entry in cells that did not activate the G2 DNA-damage checkpoint [10]. Thus, Plk1-induced Claspin degradation appears to positively regulate checkpoint recovery, and interference in this pathway severely limits the capacity of a cell to recover from genotoxic stress.

These results are in good agreement with data from yeast, where the budding yeast polo-homolog Cdc5 is required for the inactivation of the DNA checkpoint kinase Rad53 [11]. Moreover, Polo-like kinase appears to function both in checkpoint adaptation, as was shown in *Xenopus* [3], and in checkpoint recovery, as we demonstrate here in human cells. Nonetheless, there is a clear difference in the mechanism by which Claspin function is controlled in these two different organisms. Plx1-mediated phosphorylation in *Xenopus* leads to displacement of Claspin from the chromatin, without concomitant Claspin degradation [3]. In contrast, in human cells Plk1-mediated phosphorylation causes destruction of Claspin, pointing to a more rigorous, irreversible control mechanism. Interestingly, such a fundamental difference is not without precedent; others have shown that interference with the function of Orc1 occurs through its displacement from chromatin in *Xenopus*, whereas this is mediated via protein destruction in human cells [12]. Also, although binding of Plx1 to xClaspin requires ATR-dependent phosphorylation, it is likely that the priming of Claspin for binding to Plk1 in human cells involves alternative kinases [13]. The identity of these kinases and the mechanism by which Plk1 is reactivated after genotoxic stress and allowed to trigger downregulation of Claspin remain the most interesting questions to resolve because they will help us understand how a cell switches from a checkpoint-arrested state to a state of active recovery. Clearly, our data show that the action of Plk1 during recovery is not restricted to Wee1 but that it also regulates a core checkpoint component that acts at the top of the signaling cascade. In light of our data, one could even speculate that mere activation of Plk1 in DNA-damaged cells might be sufficient to silence the checkpoint and promote checkpoint adaptation.

Supplemental Data

Supplemental Data include four figures and Experimental Procedures and can be found online at: <http://www.current-biology.com/cgi/content/full/16/19/1950/DC1/>.

Acknowledgments

We acknowledge Drs. J. Bartek, D. Bohmann, J. Chen, S. Jackson, S. Nijman, P. Reaper, and N. Watanabe for providing reagents and the Medema and Freire labs as well as Dr. P. Reaper for fruitful discussions. This work was made possible by grants from the Dutch Cancer Society to M.v.V. (2004-3063) and V.S. (2005-3412), from the Association for International Cancer Research to V.S. (05-005), from ZonMW to R.M. (VICI 918.46.616), and from the Spanish Science and Education Ministry to R.F. (grants GEN2003-20243-C08-07 and SAF2004-07856). R.F. is also supported by Fondo de Investigaciones Sanitarias (FIS). I.M. is supported by a FUNCIS fellowship.

Received: May 26, 2006

Revised: August 7, 2006

Accepted: August 8, 2006

Published online: August 24, 2006

References

1. Kumagai, A., and Dunphy, W.G. (2000). Claspin, a novel protein required for the activation of Chk1 during a DNA replication checkpoint response in *Xenopus* egg extracts. *Mol. Cell* 6, 839–849.
2. Chini, C.C., and Chen, J. (2003). Human claspin is required for replication checkpoint control. *J. Biol. Chem.* 278, 30057–30062.
3. Yoo, H.Y., Kumagai, A., Shevchenko, A., Shevchenko, A., and Dunphy, W.G. (2004). Adaptation of a DNA replication checkpoint response depends upon inactivation of Claspin by the Polo-like kinase. *Cell* 117, 575–588.
4. Chini, C.C., Wood, J., and Chen, J. (2006). Chk1 is required to maintain Claspin stability. *Oncogene* 25, 4165–4171.
5. Hansen, D.V., Loktev, A.V., Ban, K.H., and Jackson, P.K. (2004). Plk1 regulates activation of the anaphase promoting complex by phosphorylating and triggering SCFbetaTrCP-dependent destruction of the APC Inhibitor Emi1. *Mol. Biol. Cell* 15, 5623–5634.
6. Moshe, Y., Boulaire, J., Pagano, M., and Hershko, A. (2004). Role of Polo-like kinase in the degradation of early mitotic inhibitor 1, a regulator of the anaphase promoting complex/cyclosome. *Proc. Natl. Acad. Sci. USA* 101, 7937–7942.
7. Watanabe, N., Arai, H., Nishihara, Y., Taniguchi, M., Watanabe, N., Hunter, T., and Osada, H. (2004). M-phase kinases induce phospho-dependent ubiquitination of somatic Wee1 by SCFbeta-TrCP. *Proc. Natl. Acad. Sci. USA* 101, 4419–4424.
8. Fuchs, S.Y., Spiegelman, V.S., and Kumar, K.G. (2004). The many faces of beta-TrCP E3 ubiquitin ligases: reflections in the magic mirror of cancer. *Oncogene* 23, 2028–2036.
9. Watanabe, N., Arai, H., Iwasaki, J., Shiina, M., Ogata, K., Hunter, T., and Osada, H. (2005). Cyclin-dependent kinase (CDK) phosphorylation destabilizes somatic Wee1 via multiple pathways. *Proc. Natl. Acad. Sci. USA* 102, 11663–11668.
10. van Vugt, M.A., Bras, A., and Medema, R.H. (2004). Polo-like kinase-1 controls recovery from a G2 DNA damage-induced arrest in mammalian cells. *Mol. Cell* 15, 799–811.
11. Pelliccioli, A., Lee, S.E., Lucca, C., Foiani, M., and Haber, J.E. (2001). Regulation of *Saccharomyces* Rad53 checkpoint kinase during adaptation from DNA damage-induced G2/M arrest. *Mol. Cell* 7, 293–300.
12. Mendez, J., Zou-Yang, X.H., Kim, S.Y., Hidaka, M., Tansey, W.P., and Stillman, B. (2002). Human origin recognition complex large subunit is degraded by ubiquitin-mediated proteolysis after initiation of DNA replication. *Mol. Cell* 9, 481–491.
13. van Vugt, M.A., and Medema, R.H. (2004). Checkpoint adaptation and recovery: Back with Polo after the break. *Cell Cycle* 3, 1383–1386.