

Binding structure of the leucine aminopeptidase inhibitor microginin FR1

Manuel Kraft^{a,1}, Christian Schleberger^{b,1}, Jürgen Weckesser^a, Georg E. Schulz^{b,*}

^a Institut für Biologie II, Microbiologie, Albert-Ludwigs-Universität, Schänzlestr. 1, D-79104 Freiburg im Breisgau, Germany

^b Institut für Organische Chemie und Biochemie, Albert-Ludwigs-Universität, Albertstr. 21, D-79104 Freiburg im Breisgau, Germany

Received 9 November 2006; revised 22 November 2006; accepted 23 November 2006

Available online 4 December 2006

Edited by Hans Eklund

Abstract Natural bioactive compounds are of general interest for pharmaceutical research because they may serve as leads in drug development campaigns. Among them, microginins are linear peptides known to inhibit various exopeptidases. The crystal structure of microginin FR1 from *Microcystis* sp. bound to bovine lens leucine aminopeptidase was established at 1.73 Å resolution. The observed binding structure could be beneficial for the design of potent aminopeptidase inhibitors.

© 2006 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

Keywords: ACE inhibition; Bestatin; Bioactive peptides; Microcystins; X-ray diffraction analysis

1. Introduction

Water blooms in eutrophic lakes are often caused by cyanobacteria [1]. Field populations of the common bloom-forming genus *Microcystis* can be toxic because some strains produce microcystins, which are hepatotoxic, promote tumors and affect the development of fish and amphibians [2–4]. In addition, cyanobacteria produce non-toxic peptides, some of which show remarkable bioactivities and have therefore attracted the attention of the pharmaceutical industry [5,6]. Of particular interest are phosphatase inhibitors [7] and selective protease inhibitors, as for example the elastase inhibitor scyptolin A [8].

Among these peptides are microginins (Fig. 1), the first of which was isolated from *Microcystis aeruginosa* (NIES-100) [9]. Microginins contain the lipidic β -amino acid Ahda (3-amino-2-hydroxy-decanoic acid) and inhibit the angiotensin-converting enzyme (ACE) [9] as well as aminopeptidases but do not affect papain, trypsin, chymotrypsin or elastase [10]. Related to the microginins are the well-known inhibitors bestatin [11] and amastatin [12] produced by *Streptomyces*. Both of them inhibit aminopeptidases [13]; bestatin causes a reduction of the HIV-infection rate [14]. Here we report the binding structure of microginin FR1 that was extracted from a water bloom in a local lake [15] to a leucine aminopeptidase (LAP).

2. Materials and methods

Microginin FR1 was isolated from water bloom material in Lake Waltershofen near Freiburg and the IC₅₀ value for bovine lens LAP was determined as described [16]. The enzyme was isolated from calf lenses as described [17]. The calf eyes were received 5 h after slaughtering and stored on wet ice. The isolated enzyme ran through an additional gel permeation column (Superdex-200 26/60, Amersham) equilibrated with buffer A (50 mM Tris-HCl pH 7.8, 50 μ M ZnSO₄ and 200 mM NaCl). LAP-containing fractions were pooled and concentrated to 7 mg/ml in buffer A. Microginin FR1 dissolved in DMSO was added at a concentration of 5 mM to the enzyme solution, incubated for 30 min at 37 °C and centrifuged. Using the hanging drop set-up, 10 μ l of the enzyme-plus-inhibitor solution was equilibrated against buffer B (50 mM Tris-HCl pH 7.8, 50 μ M ZnSO₄, 50% (v/v) 2-methyl-2,4-pentanediol) [18]. The crystals were flash-frozen to 100 K without any further cryo-protectant.

X-ray diffraction data were collected at beamline BL14-1 of BESSY (Berlin) and processed using the programs XDS and XSCALE [19]. The structure of bovine lens LAP (Protein Data Bank code 1BLL) [18] was used as a starting model for structure determination. After rigid body refinement with the program REFMAC5 [20], model bias was removed by a simulated annealing run starting at a temperature of 2500 K with CNS [21]. An initial map was calculated using RESOLVE [22]. After manual adjustments with the program COOT [23], the model was refined in REFMAC5 and the inhibitor was placed in the resulting (F_o–F_c) difference electron density map. An energy-minimized conformation of the inhibitor and the corresponding dictionary files were generated with PRODRG [24]. Water molecules were placed using COOT and the refinement was finalized with REFMAC5. The coordinates and structure factors are deposited in the Protein Data Bank code 2J9A.

3. Results and discussion

Purified microginin FR1 was added in excess to a solution of LAP from bovine lens, and cocrystals were obtained by the hanging drop technique. X-ray diffraction data were collected to 1.73 Å resolution and the structure was determined (Table 1). An initial (F_o–F_c)-electron density map showed the inhibitor binding conformation of the first three residues unambiguously. After refinement, the remaining two residues were also placed unambiguously although with higher *B*-factors (Fig. 2A).

Bovine lens LAP is a homo-hexamer with D₃ symmetry (Fig. 2B). Two large contacts around the twofold axes are formed by the N- and C-terminal domains in different subunit pairings, tightly interconnecting the hexamer. The C-terminal domains form a further contact around the threefold axis. The active centers are on the inside of the hexamer lining a large central cavity. They are shielded by the N-terminal domains but accessible through six channels with a width appropriate for an extended peptide chain (Fig. 2C). With such a topology, the

*Corresponding author. Fax: +49 761 203 6161.

E-mail address: georg.schulz@ocbc.uni-freiburg.de (G.E. Schulz).

¹These authors have contributed equally.

Abbreviations: ACE, angiotensin-converting enzyme; Ahda, (2S, 3R)-3-amino-2-hydroxy-decanoic-acid; LAP, leucine aminopeptidase




Fig. 1. Covalent structure of the pentapeptide microginin FR1 with its characteristic Ahda residue [15]. The related dipeptide bestatin [11] and tetrapeptide amastatin [12] carry a modified Ahda as the first residue. In bestatin and amastatin the atoms C5–C10 of Ahda are replaced by phenyl and isopropyl, respectively. The other residues of bestatin and amastatin are -Leu and -Val-Val-Asp, respectively. All amino acid residues have the L-configuration. The position of the scissile bond of processed peptides is indicated by the assigned subsites S1 and S1'.

enzyme can only cleave unfolded peptides, because folded peptides or proteins cannot enter the central cavity.

The Ahda-Ala part of microginin FR1 fits snugly into the active center pocket (Fig. 3), binding with full occupancy as indicated by the crystallographic *B*-factors matching those of the environment. The decanoic acid moiety of Ahda is accommodated in a spacious pocket extension formed by Leu269, Met270, Asp365, Ala451, Met454 and Thr455. This extension is for the most part nonpolar but contains Asp365 at one side, suggesting that a positive charge at this part of the inhibitor would increase the binding strength. The amino group of Ahda interacts with one of the two zinc ions at the active center (Zn1) as well as with Thr359-O, while the hydroxyl group of Ahda interacts with Zn2 and with a water molecule tightly bound to the backbone amides of Glu334 and Gly335. The 1-carbonyl of Ahda forms a hydrogen bond to Lys262-N ϵ . The alanine of microginin FR1 forms hydrogen bonds with Leu360-O and Gly362-N reminiscent of a piece of parallel

β -sheet. Ahda and alanine occupy subsites S1 and S1' (Fig. 1), thus covering the active center of the LAP.

Instead of the carbonate or the three water molecules found in other bovine lens LAP structures [25,26], we observed a chloride ion bound to Arg366 close to the metal ion Zn2 (Fig. 3A). Since a water molecule placed at this position caused positive difference electron density and since the observed density is at the level of several sulfur atoms, a chloride ion from the crystallization buffer is the most likely interpretation.

The three C-terminal residues of the inhibitor are involved in less clearly defined binding patterns. The third residue *N*-methyl-leucine binds with its carbonyl oxygen through a water network to Arg336 and Ala363-O. The remaining two tyrosines of microginin FR1 show rather high *B*-factors (Fig. 3A). The first tyrosine binds with its hydroxyl through Arg425' to Asp365. The hydroxyl of the second tyrosine forms hydrogen bonds through water molecules and directly to the backbone of residues 423 and 424. Despite their high *B*-factors, the removal of these tyrosines weaken the inhibitory capacity appreciably [16]. Other inhibitor complexes with bovine lens LAP are known [25].

The cyanobacterial peptide microginin FR1 represents a large group of known microginins characterized by the N-terminal residue Ahda [9,10,27,28]. The IC₅₀ values of microginin FR1 for bovine lens cytosolic LAP (EC 3.4.11.1) was determined as 1.3 μ M, which is lower than the value 16 μ M for ACE [15]. The IC₅₀ values of the other microginins for ACE are in the range of 10 μ M or above [9,16,28]. Stronger inhibition was observed for the microsomal LAP (EC 3.4.11.2). Microginin FR1 shows an IC₅₀ value of 6 nM with the porcine kidney microsomal LAP, while other microginins range from 16 nM to around 10 μ M [9,10,16,28]. The related inhibitors bestatin and amastatin show IC₅₀ values in the micromolar range for cytosolic and microsomal LAP [13,29].

The inhibited enzymes cytosolic LAP, microsomal LAP and ACE are all exopeptidases with one or two zinc ions in the active center. However, their structures are completely different [30–32]. All discussed inhibitors have Ahda-related first residues (Fig. 1), which bind to the metal centers of cytosolic and microsomal LAP as shown in Fig. 3 and elsewhere [18,30,32,33]. The reported binding structure to cytosolic

Table 1
Structure determination^a

| | |
|--|-------------------------|
| <i>Data collection</i> | |
| Resolution (Å) | 35 – 1.73 (1.80 – 1.73) |
| Unique reflections | 62705 (6822) |
| Multiplicity | 6.0 (5.4) |
| Completeness (%) | 99.4 (98.1) |
| <i>R</i> _{sym} (%) | 6.9 (47) |
| <i>I</i> / σ _{<i>I</i>} | 19.0 (3.5) |
| Wilson <i>B</i> -factor (Å ²) | 23.1 |
| <i>Refinement</i> | |
| Protein atoms | 3921 |
| Solvent atoms | 539 |
| Microginin FR1 atoms | 52 |
| <i>R</i> _{cryst} / <i>R</i> _{free} (5% test set) | 14.5/17.4 |
| Average <i>B</i> -factor (Å ²) | 19 |
| Rmsd bond lengths (Å)/bond angles (°) | 0.019/1.596 |
| Ramachandran:favored/allowed (%) | 93.1/6.9 |

^aThe data were collected at a wavelength of 0.9184 Å at BESSY (Berlin). The space group was P6₃22 with unit cell parameters *a* = *b* = 130.0 Å, *c* = 120.8 Å and one subunit in the asymmetric unit.




Fig. 2. Stereoview of microginin FR1 bound to bovine lens LAP. (A) The final model of one subunit of the enzyme represented as a ribbon plot with the bound inhibitor shown in its ($2F_o - F_c$)-electron density at a contour level of 0.8σ . The two zinc ions of the active center are given for reference (pink). The N-terminal (residues 1–158) and C-terminal (159–485) domains are in yellow and red colors, respectively. The threefold (green) and twofold (black) axes of the hexamer are given. (B) Surface representation of the hexameric enzyme viewed along the threefold axis with colored subunits and labeled domains. The twofold axes are black. The entrances to the active centers are marked by pink sticks. (C) Hexamer in the inflated-stick-representation with the same colors, axes and sticks (pink) as in panel B. The hexamer is cut parallel but 6 Å above the plane of the twofold axes. Microginin FR1 is pink and labeled (Inh).




Fig. 3. Close-up stereoview of the active center of bovine lens LAP. (A) The binding structure of the inhibitor microginin FR1. Chain cuts are marked by halos. The two zinc ions are gray, the water molecules are blue and the chloride ion is green. Residues of a neighboring subunit are marked by a prime. Hydrogen bonds are black dashed lines. The zinc coordinations are red dashed lines (distances 2.0–2.3 Å). Van-der-Waals contacts are green dotted lines (all unlabeled distances are between 3.6 Å and 4.6 Å). The *B*- factors of the inhibitor residues Ahda-Ala-mLeu-Tyr-Tyr are 15, 13, 21, 51 and 81 Å², respectively. (B) Bound microginin FR1 with zinc ions (pink) superimposed on the bound inhibitors amastatin (blue) [18] and L-leucinephosphonate (pink) [33]. A structure of bound bestatin was reported [30] but not deposited in the Protein Data Bank.

LAP resembles the known binding structure of the microginin-related bestatin to microsomal LAP [32] but differs in detail.

Among the wide variety of peptides produced by cyanobacteria, the microginins with their lipidic β -amino acid Ahda inhibit both types of leucine aminopeptidases, cytosolic (EC 3.4.11.1) and microsomal (EC 3.4.11.2), the structures of which are quite different [30,32]. Since these aminopeptidases are of medicinal relevance [13,14,34] and since the inhibition occurs at nanomolar to micromolar concentrations, the microginins constitute a welcome lead for the design of novel drugs. The microginins may also be used as a starting point for ACE inhibitor design.

Acknowledgements: We thank Dr. Erschig for the allocation of calf eyes and the team of BESSY (Berlin) for their help in data collection. The project was supported by the Deutsche Forschungsgemeinschaft under grant GRK-434.

References

- [1] Hallegraef, G.M. (1993) A review of harmful algal blooms and their apparent global increase. *Phycologia* 32, 79–99.
- [2] Carmichael, W.W. (1996) Toxic microcystin and the environment in: *Toxic Microcystin* (Watanabe, M., Harada, K., Carmichael, W.W. and Fujiki, H., Eds.), pp. 1–11, CRC Press, Boca Raton, FL, USA.

- [3] Obereim, A., Becker, J., Codd, G.A. and Steinberg, C. (1999) Effects of cyanobacterial toxins and aqueous crude extracts of cyanobacteria on the development of fish and amphibians. *Environ. Toxicol.* 14, 77–88.
- [4] Nishiwaki-Matsushima, R., Ohta, T., Nishiwaki, S., Suganuma, M., Kohyama, K., Ishikawa, T., Carmichael, W.W. and Fujiki, H. (1992) Liver tumor promotion by the cyanobacterial cyclic peptide toxin microcystin-LR. *J. Cancer Res. Clin. Oncol.* 118, 420–424.
- [5] Namikoshi, M. and Rinehart, K.L. (1996) Bioactive compounds produced by cyanobacteria. *J. Ind. Microbiol.* 17, 373–384.
- [6] Weckesser, J., Martin, C. and Jakobi, C. (1996) Cyanopeptolins, depsipeptides from cyanobacteria. *Syst. Appl. Microbiol.* 19, 133–138.
- [7] Goldberg, J., Huang, H.B., Kwon, Y.G., Greengard, P., Nairn, A.C. and Kuriyan, J. (1995) Three-dimensional structure of the catalytic subunit of protein serine/threonine phosphatase-1. *Nature* 376, 745–753.
- [8] Matern, U., Schleberger, C., Jelakovic, S., Weckesser, J. and Schulz, G.E. (2003) Binding structure of elastase inhibitor scyptolin A. *Chem. Biol.* 10, 997–1001.
- [9] Okino, T., Matsuda, H., Murakami, M. and Yamaguchi, K. (1993) Microginin, an angiotensin-converting enzyme inhibitor from the blue-green alga *Microcystis aeruginosa*. *Tetrahedron Lett.* 34, 501–504.
- [10] Ishida, K., Matsuda, H., Murakami, M. and Yamaguchi, K. (1997) Microginins 299-A and -B, leucine aminopeptidase inhibitors from the cyanobacterium *Microcystis aeruginosa* (NIES-299). *Tetrahedron* 53, 10281–10288.
- [11] Umezawa, H., Aoyagi, T., Suda, H., Hamada, M. and Takeuchi, T. (1976) Bestatin, an inhibitor of aminopeptidase B, produced by actinomycetes. *J. Antibiot.* 29, 97–99.
- [12] Aoyagi, T., Tobe, H., Kojima, F., Hamada, M., Takeuchi, T. and Umezawa, H. (1978) Amastatin, an inhibitor of aminopeptidase A, produced by actinomycetes. *J. Antibiot.* 31, 636–638.
- [13] Rich, D.H., Moon, B.J. and Harbeson, S. (1984) Inhibition of aminopeptidases by amastatin and bestatin derivatives. Effect of inhibitor structure on slow-binding processes. *J. Med. Chem.* 27, 417–422.
- [14] Pulido-Cejudo, G., Conway, B., Proulx, P., Brown, R. and Izaguirre, C.A. (1997) Bestatin-mediated inhibition of leucine aminopeptidase may hinder HIV infection. *Antiviral Res.* 36, 167–177.
- [15] Neumann, U., Forchert, A., Flury, T. and Weckesser, J. (1997) Microginin FR1, a linear peptide from a water bloom of *Microcystis* species. *FEMS Microbiol. Lett.* 153, 475–478.
- [16] Ishida, K., Kato, T., Murakami, M., Watanabe, M. and Watanabe, M.F. (2000) Microginins, zinc metalloproteases inhibitors from the cyanobacterium *Microcystis aeruginosa*. *Tetrahedron* 56, 8643–8656.
- [17] Allen, M.P., Yamada, A.H. and Carpenter, F.H. (1983) Kinetic parameters of metal-substituted leucine aminopeptidase from bovine lens. *Biochemistry* 22, 3778–3783.
- [18] Kim, H. and Lipscomb, W.N. (1993) X-ray crystallographic determination of the structure of bovine lens leucine aminopeptidase complexed with amastatin: Formulation of a catalytic mechanism featuring a *gem*-diolate transition state. *Biochemistry* 32, 8465–8478.
- [19] Kabsch, W. (1993) Automatic processing of rotation diffraction data from crystals of initially unknown symmetry and cell constants. *J. Appl. Crystallogr.* 26, 795–800.
- [20] Murshudov, G.N., Vagin, A.A. and Dodson, E.J. (1997) Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr. D* 53, 240–255.
- [21] Brünger, A.T., Adams, P.D., Clore, G.M., DeLano, W.L., Gros, P., Grosse-Kunstleve, R.W., Jiang, J.S., Kuszewski, J., Nilges, M., Pannu, N.S., Read, R.J., Rice, L.M., Simonson, T. and Warren, G.L. (1998) Crystallography & NMR system: A new software suite for macromolecular structure determination. *Acta Crystallogr. D* 54, 905–921.
- [22] Terwilliger, T.C. (2000) Maximum-likelihood density modification. *Acta Crystallogr. D* 56, 965–972.
- [23] Emsley, P. and Cowtan, K. (2004) COOT: model-building tools for molecular graphics. *Acta Crystallogr. D* 60, 2126–2132.
- [24] Schüttelkopf, A.W. and van Aalten, D.M. (2004) PRODRG: a tool for high-throughput crystallography of protein–ligand complexes. *Acta Crystallogr. D* 60, 1355–1363.
- [25] Sträter, N. and Lipscomb, W.N. (1995) Two-metal ion mechanism of bovine lens leucine aminopeptidase: active site solvent structure and binding mode of L-leucinol, a *gem*-diolate transition state analogue, by X-ray crystallography. *Biochemistry* 34, 14792–14800.
- [26] Sträter, N., Sun, L., Kantrowitz, E.R. and Lipscomb, W.N. (1999) A bicarbonate ion as a general base in the mechanism of peptide hydrolysis by dizinc leucine aminopeptidase. *Proc. Natl. Acad. Sci. USA* 96, 11151–11155.
- [27] Ishida, K., Matsuda, H. and Murakami, M. (1998) Four new microginins, linear peptides from the cyanobacterium *Microcystis aeruginosa*. *Tetrahedron* 54, 13475–13484.
- [28] Sano, T., Takagi, H., Morrison, L.F., Metcalf, J.S., Codd, G.A. and Kaya, K. (2005) Leucine aminopeptidase M inhibitors, cyanostatin A and B, isolated from cyanobacterial water blooms in Scotland. *Phytochemistry* 66, 543–548.
- [29] Wilkes, S.H. and Prescott, J.M. (1985) The slow, tight binding of bestatin and amastatin to aminopeptidases. *J. Biol. Chem.* 260, 13154–13162.
- [30] Burley, S.K., David, P.R., Sweet, R.M., Taylor, A. and Lipscomb, W.N. (1992) Structure determination and refinement of bovine lens leucine aminopeptidase and its complex with bestatin. *J. Mol. Biol.* 224, 113–140.
- [31] Natesh, R., Schwager, S.L.U., Sturrock, E.D. and Acharya, K.R. (2003) Crystal structure of the human angiotensin-converting enzyme-lisinopril complex. *Nature* 421, 551–554.
- [32] Addlagatta, A., Gay, L. and Matthews, B.W. (2006) Structure of aminopeptidase N from *Escherichia coli* suggests a compartmentalized, gated active site. *Proc. Natl. Acad. Sci. USA* 103, 13339–13344.
- [33] Sträter, N. and Lipscomb, W.N. (1995) Transition state analogue L-leucinephosphonic acid bound to bovine lens leucine aminopeptidase: X-ray structure at 1.65 Å resolution in a new crystal form. *Biochemistry* 34, 9200–9210.
- [34] Hooper, N.M. and Lendeckel, U., Eds., (2004). *Aminopeptidases in Biology and Disease*, Springer Verlag, Berlin.