

Contents lists available at ScienceDirect

Cancer Epidemiology

The International Journal of Cancer Epidemiology, Detection, and Prevention

journal homepage: www.cancerepidemiology.net

European Code against Cancer 4th Edition: Alcohol drinking and cancer[☆]



Chiara Scoccianti^a, Michele Cecchini^{b,1}, Annie S. Anderson^c, Franco Berrino^d,
Marie-Christine Boutron-Ruault^e, Carolina Espina^a, Timothy J. Key^f, Michael Leitzmann^g,
Teresa Norat^h, Hilary Powersⁱ, Martin Wiseman^j, Isabelle Romieu^{a,*}

^a International Agency for Research on Cancer (IARC), 150 Cours Albert Thomas, 69372 Lyon Cedex 08, France^b Health Policy Analyst OECD, 2 rue André Pascal, 75775 Paris Cedex 16, France^c Centre for Research into Cancer Prevention & Screening, Level 7, Mailbox 7, Ninewells Hospital & Medical School, Dundee, DD1 9SY, Scotland, United Kingdom^d Fondazione IRCCS Istituto Nazionale dei Tumori, 1 via Venezian, 20133 Milan, Italy^e Institut Gustave Roussy, 114 rue Edouard Vaillant, 94805 Villejuif, France^f Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, United Kingdom^g Department of Epidemiology and Preventive Medicine, University of Regensburg, 93042 Regensburg, Germany^h Department of Epidemiology and Biostatistics, School of Public Health Imperial College London, St Mary's Campus, London W2 1 PG, United Kingdomⁱ Human Nutrition Unit, The Medical School, Beech Hill Road, Sheffield, S10 2RX, United Kingdom^j World Cancer Research Fund International, Second Floor, 22 Bedford Square, London WC1B 3HH, United Kingdom

ARTICLE INFO

Article history:

Received 2 July 2014

Received in revised form 10 November 2014

Accepted 16 January 2015

Available online 24 June 2015

Keywords:

Alcohol drinking

Cancer

Ethanol

Acetaldehyde

Disease

Primary prevention

Europe

ABSTRACT

Alcohol consumption is the third leading risk factor for disease and mortality in Europe. The International Agency for Research on Cancer (IARC) Monographs provide strengthened evidence that the consumption of alcoholic beverages is causally associated with cancers of the oral cavity, pharynx, larynx, oesophagus, liver, colorectum and female breast, even for low and moderate alcohol intakes. The risk of cancer increases in a dose-dependent manner, and the higher the amount of alcohol consumed, the higher the risk of developing cancer. Several biological mechanisms explain the carcinogenicity of alcohol; among them, ethanol and its genotoxic metabolite acetaldehyde play a major role. Taking all this evidence into account, a recommendation of the 4th edition of the European Code against Cancer (ECAC) is: "If you drink alcohol of any type, limit your intake. Not drinking alcohol is better for cancer prevention."

© 2015 International Agency for Research on Cancer; Licensee ELSEVIER Ltd <https://creativecommons.org/licenses/by-nc-nd/3.0/igo/>

[☆] This is an Open Access article published under the CC BY NC ND 3.0 IGO license which permits users to download and share the article for non-commercial purposes, so long as the article is reproduced in the whole without changes, and provided the original source is properly cited. This article shall not be used or reproduced in association with the promotion of commercial products, services or any entity. There should be no suggestion that IARC endorses any specific organisation, products or services. The use of the IARC logo is not permitted. This notice should be preserved along with the article's original URL.

Abbreviations: IARC, International Agency for Research on Cancer; EU, European Union; RR, relative risk; UADT, upper aero digestive tract; ADH, alcohol dehydrogenase; ALDH, acetaldehyde dehydrogenase; MTHFR, methylenetetrahydrofolate reductase; ER, estrogen receptor; PR, progesterone receptor; WHO, World Health Organization.

* Corresponding author at: IARC European Code against Cancer Secretariat, 150 Cours Albert Thomas, F-69372 Lyon Cedex 08, France. Tel.: +33 4 72 73 84 85.

E-mail address: secretariat@cancer-code-europe.iarc.fr (I. Romieu).

¹ The views expressed are those of the author and not necessarily those of the OECD, or its member countries.

1. Introduction

Alcohol consumption is linked to a large number of health impairments, chronic diseases and deaths worldwide [1]. The 2012 Monograph of the International Agency for Research on Cancer (IARC) provides stronger evidence on the carcinogenicity of alcohol by tumour sites and by mechanisms of alcohol carcinogenesis even for low and moderate alcohol intakes [2,3]. The IARC Monographs reached the conclusion: "alcohol consumption is carcinogenic to humans (Group 1); ethanol in alcoholic beverages is carcinogenic to humans (Group 1); acetaldehyde associated with the consumption of alcoholic beverages is carcinogenic to humans (Group 1)" [2]. Overall, there is no consistent difference in cancer risk between different types of alcoholic beverage [4–6]. Europe is the region of highest alcohol consumption in the world, with an average consumption of more than twice the global average; it has

<http://dx.doi.org/10.1016/j.canep.2015.01.007>

1877-7821/© 2015 International Agency for Research on Cancer; Licensee ELSEVIER Ltd <https://creativecommons.org/licenses/by-nc-nd/3.0/igo/>




Fig. 1. Average European adult alcohol consumption in litres/capita/year among populations aged ≥ 15 years (years 2010 and 1980). From calculations provided by the OECD based on "Health at a glance: Europe 2012. OECD Publishing, 2012" [8].

a high prevalence of hazardous drinkers and an average alcohol-attributable cancer burden which also exceeds by far the global average [1]. Taken together, the 4th edition of the European Code Against Cancer (ECAC) [7] advocates action-oriented recommendations for the general public. The ECAC recommends decreasing or cutting alcohol consumption in order to prevent several types of cancer and to improve overall health.

1.1. European alcohol consumption among adult and young generations

Per capita alcohol consumption has been falling in the European Union (EU) as a whole over the past three decades, while remaining particularly high compared to the global average. The most recent sales data show that individuals aged ≥ 15 years drink on average 10.7 l of pure alcohol per year [8]. Consumption tends to be higher in the Central-Eastern and Eastern countries such as, for example, Latvia, Romania, Lithuania and Austria, all of which have an average consumption >12 l per capita. At the other end of the spectrum, Mediterranean countries (e.g., Italy, Malta and Greece) and Nordic Countries (e.g., Norway, Sweden and Iceland) have relatively lower levels of consumption, in the region of 7–8 l of pure alcohol per adult person (Fig. 1). Gender, age and socio-economic status are key factors in determining levels of alcohol consumption [9]. Men are more likely to consume alcohol than women, and to drink more when they do [10], particularly in Central, Western and Northern EU countries. Compared to older adults, young and middle-aged people tend to drink higher volumes of alcohol [11]. Women with higher level of education tend to drink more alcohol while the opposite is generally true for men [10]. Hazardous drinking behaviours, such as binge drinking (i.e., the occasional consumption of ≥ 60 g of pure alcohol on the same occasion and at least one day in the last month), have been increasing over the past 20 years [13], especially in Germany and Ireland and among younger generations [14–17]. The prevalence of binge drinkers doubled in France and has increased by about 30% in Germany between 2002 and 2008 [13]; binge drinking was reported by 36% of girls and 40% of boys in 2010 [8].

1.2. Effect of alcohol drinking on coronary heart disease

Light to moderate alcohol consumption might be associated with a reduced risk of coronary heart disease. A small reduction in risk has been suggested for the consumption of one drink every

second day (not including as 'abstainers' those who reduced or stopped drinking [18]), and is probably confined to middle-aged or older individuals [18–20]. Similarly, a recent systematic review suggests that light to moderate alcohol consumption might be associated with a reduced risk of cardiovascular outcome, in particular coronary heart disease mortality compared to stroke [21], but again reverse causation and residual confounding cannot be excluded. While there is some evidence that low doses of alcohol may raise blood levels of high-density lipoprotein and improve reduction in coagulation [22,23], high doses may precipitate cardiac arrhythmias, myocardial ischaemia or infarction, and coronary death [24–26].

2. Association with cancer

2.1. Cancer types associated with alcohol drinking

As established by the IARC Monographs, consumption of alcoholic beverages is causally associated with cancers of the oral cavity, pharynx, larynx, oesophagus, colorectum, liver (hepatocellular carcinoma), and female breast in a dose-dependent manner [2]. Summary relative risks (RR) supporting these associations both for light and heavy drinking are indicated in Table 1. The relationship between alcohol consumption and cancer risk is monotonic and without a threshold (Fig. 2). Alcohol consumption accounts for about 3% and 10% of total cancers diagnosed in women and men, respectively [27]. In both genders, the alcohol-attributable fraction is high for upper aero-digestive tract (25–44%), liver (18–33%), and colorectal (4–17%) cancers, and in women for breast cancer (about 5%), with variation across EU countries related to different levels of exposure to alcohol [27]. Drinking patterns play an important role in modulating the relationship between alcohol and cancer risk. The strongest associations are observed for heavy drinking, in particular regular heavy drinking. Any reduction in alcohol consumption has a beneficial effect on reducing the risk of cancer.

2.1.1. Neoplasms of the upper digestive tract: oral cavity, pharynx, larynx and squamous-cell carcinoma of the oesophagus

Results from cohort studies and recent meta-analyses provide convincing evidence that the consumption of alcoholic beverages increases the risk of neoplasms of the upper digestive and respiratory tracts (UADT) even in the absence of smoking. Significant dose–response relationships are found with different

Table 1
Meta-analyses on alcohol consumption and cancer risk at different intakes.

Cancer	ICD-10 code	Reference	RR (95%CI) Light drinkers [#]	RR (95%CI) Heavy drinkers [#]
Oral cavity	C00–C13	Turati et al. [89]	1.17 (1.01–1.35)	4.64 (3.78–5.70)
Pharynx			1.23 (0.87–1.73)	6.62 (4.72–9.29)
OP cancers		Turati et al. [33]	2.55 (2.15–3.02)	5.40 (4.49–6.50)
Larynx	C32	Islami et al.* [31]	0.88 (0.71–1.08)	2.62 (2.13–3.23)
Oesophagus	C15	Tramacere et al. [90]	0.92 (0.78–1.09)	1.16 (0.92–1.46)
		Islami et al. [32]	1.32 (0.90–1.60)	3.35 (2.35–4.78)
Colorectal	C18–C21	Fedirko et al. [43]	1.21 (1.13–1.28)	1.52 (1.27–1.81)
Liver	C22	Corrao et al.* [48]	1.19 (1.12–1.27)	1.40 (1.25–1.56)
Female breast	C50	Bagnardi et al. [44]	1.05 (1.02–1.08)	
		Corrao et al.* [48]	1.25 (1.20–1.29)	1.55 (1.44–1.67)

ICD-10: International Classification of Diseases, 10th Revision [91].

[#] Light drinker (≤ 1 drink/g or 12.5 g/day); heavy drinker (≥ 4 drinks/day or 50 g/day).

* Light drinker (≤ 25 g/day); heavy drinker (≥ 50 g/day).

metrics of exposure to alcohol such as level, frequency [28] and duration [29] of consumption, and for moderate intake in women [4,30,31]. Among non-smokers, reported RRs for oesophageal squamous-cell carcinoma and laryngeal cancer range from 0.74 (95%CI: 0.47–1.16) for light intakes to 3.09 (95%CI: 1.75–5.46) for high intakes [31,32]. Risks for oropharyngeal cancer are as high as 5.40 (95%CI: 4.49–6.50) [33]. The combined exposure to alcohol drinking and tobacco smoking results in a supra-multiplicative synergistic effect which enhances the risk of these neoplasms up to 14-fold, among heavy smokers and heavy drinkers (four or more drinks/day) [34,35] (Fig. 3).

There is evidence that the risk of UADT cancer decreases with time since drinking cessation, without however ever falling to that of lifetime abstainers. A recent meta-analysis reports an average 2% decreased risk of pharyngeal and laryngeal cancers per year of cessation [36]. A trend for decreased risk is suggested for laryngeal, oropharyngeal and oesophageal cancers among former drinkers. After at least 6 years of drinking cessation, odds ratios decreased to 1.24 (95%CI: 0.68–2.47) [37], 1.74 (95%CI: 0.77–3.92) [38], and 0.85 (95%CI: 0.78–0.92) for laryngeal, oropharyngeal and oesophageal cancers, respectively [39,40].

2.1.2. Neoplasms of the lower digestive tract: colorectum and liver

Results from cohort studies and recent meta-analyses provide overall evidence of a linear dose–response relationship between average alcohol consumption and cancers of the liver and colorectum. Summary effect estimates for colorectal cancer are about 11% (RR = 1.11; 95%CI: 0.90–1.38) at one drink/day (corresponding to 10–12 g of ethanol) and 40% (RR = 1.41; 95%CI: 1.16–1.72) at more than 4–5 drinks/day [41]. The risk is higher for drinkers with a family history of colorectal cancer (RR = 2.80; 95%CI: 2.00–3.91 at ≥ 30 g/d compared with non-drinkers with no family history) [42]. Overall, a slightly higher risk is reported for colon cancer (RR = 1.16, 95%CI: 0.97–1.39) than for rectal cancer (RR = 1.11, 95%CI: 0.97–1.29) at one drink/day [41], and the risk is greater in men (RR 1.24; 95%CI: 1.13–1.28) than in women (RR 1.08; 95%CI: 1.03–1.13) at less than 1 drink/day [43], compared to never drinkers. In both genders combined, dose–response analyses show RRs of 1.07 (95%CI: 1.04–1.10), 1.38 (95%CI: 1.28–1.50) and 1.82 (95%CI: 1.41–2.35) at 10, 50 and 100 g/day of alcohol consumption, respectively [43]. Associations of alcohol consumption with hepatocellular carcinoma are generally found at high intakes [44] with significant increased risks of about 30–40% at consumption levels >40 g/day [45–48]. An increased risk at low to moderate alcohol consumption is also reported [4,47,49].

2.1.3. Neoplasms of the female breast

A positive association between alcoholic beverage consumption and breast cancer risk is supported by more than 100 epidemiological studies [2]. The risk is increased by 8% for post-menopausal

breast cancer, 9% for pre-menopausal breast cancer, and 10% for overall breast cancer [50], per 10 g/day of ethanol. While studies on binge drinking and cancer risk are still sparse, and a standardised measure of exposure is lacking, the available evidence shows that risk is increased between 33% (95%CI: 1.11–1.59, for monthly binge drinking) [51] and 55% (95%CI: 1.07–2.26, for weekly binge drinking) [52].

The degree of mammary development may determine a susceptibility window for exposure to alcohol consumption, which may be further defined by hormonal status or first full-term pregnancy [53]. However, at present few studies have investigated whether and to what extent the age at exposure increases cancer risk, and the evidence is generally limited. In the Nurses' Health Study (NHS) II, alcohol consumption during early adulthood was dose-dependently associated with an increased risk of proliferative benign breast disease, which may lead to invasive breast cancer later in life [54]. In contrast, a NHS prospective analysis reported similar breast cancer risks for categories of age at alcohol exposure of 18–40 years and >40 years (RRs = 1.21, 95%CI: 0.88–1.67 and 1.18, 95%CI: 1.03–1.34, respectively, for a daily intake of ≥ 20 g compared to none) [51]. A large Danish cohort of post-menopausal women also could not confirm a risk difference according to the age at which drinking was started [55].


2.2. Possible biological mechanisms

Alcoholic beverages mainly comprise ethanol and water and, in minor percentages, other volatile and non-volatile compounds. Ethanol is the most important carcinogen; moreover, the metabolism of ethanol activates various pro-carcinogens present, as well as inhaled or ingested, in alcoholic beverages. Regarding breast cancer, ethanol most likely acts as both a weak cumulative carcinogen and a tumour promoter of pre-existing breast cancer cells [56]. For cancers of the oral cavity, pharynx, larynx and oesophagus, the higher than multiplicative risk observed among drinkers and smokers is explained by the synergistic effect of carcinogens contained in tobacco smoke and in alcoholic beverages which leads to mucosal hyperproliferation [57–60].


The risk of digestive tract cancers is modulated by variants of the ethanol-metabolising enzymes, the alcohol and acetaldehyde dehydrogenases (ADH and ALDH). The *ADH1B* 47Arg allele is strongly associated with increased UADT cancer risk (RR = 3.47; 95%CI: 2.76–4.36 for the Arg/Arg genotype compared to the His/His genotype) [61]. The fast-metabolising *ADH1C**1 variant, which is more frequent in Caucasians [62], may increase the risk of head and neck cancer [63,64]. The *ALDH2* Lys487 allele promotes acetaldehyde accumulation and is associated with higher risk of oesophageal cancer [2].

Ethanol also disrupts folate metabolism regulated by methylenetetrahydrofolate reductase (MTHFR) [65]; an individual's

Neoplasms of the upper digestive tract



Neoplasms of the lower digestive tract



Other neoplasms




Fig. 2. Relationship between average daily alcohol consumption and relative risk of cancer. Plots of the relative risk functions for all cancers causally associated with alcohol consumption. The relationship between alcohol consumption and cancer risk is monotonic and without a threshold. From Corrao et al. [48] <http://dx.doi.org/10.1016/j.ypmed.2003.11.027> PMID:15066364.

genotype for the C677T *MTHFR* variant modulates the effect of alcohol consumption on cancer risk. The CC or CT variants may increase colorectal cancer risk among alcohol drinkers (RR = 1.87; 95%CI: 1.29–2.71) [66]. The TT variant decreases *MTHFR* activity by 70% and appears to modulate risks of breast cancer, in association

with low folate intake [67,68], and of oral cancer [69], at daily consumption of ≥ 2 drinks.

The alcohol-related increase in androgen and oestrogen levels [70] has been suggested as a key mechanism for breast cancer's development. Most studies observe overall a stronger association




Fig. 3. Estimated relative risk (RR) for the interaction between tobacco smoking and alcohol drinking on cancers of the upper respiratory tract (reference category, risk = 1). Combined exposure to alcohol drinking and tobacco smoking increases the risk of upper digestive and respiratory tract neoplasms in a supra-multiplicative manner. From P. Boyle, P. Autier, H. Bartelink, et al. European Code Against Cancer and scientific justification: third version (2003) *Annals of Oncology* 2003, 14: 973–1005, by permission of Oxford University Press [7].

with the most common oestrogen-receptor-positive (ER+) and/or progesterone-receptor-positive (PR+) markers compared with ER– and/or PR– breast tumours, for the highest versus the lowest alcohol consumption group [71–73]. A meta-analysis showed that for each additional drink, breast cancer risk increases by 12% among ER+ subjects, by 11% among ER+PR+ subjects, by 15% among ER+PR– subjects, and by 7% for all ER– subjects, and no association was apparent with ER–/PR– or ER–/PR+ tumours when considered separately [74].

Other ethanol-mediated toxic effects associated with cancer development are cirrhosis of the liver and gastro-oesophageal reflux disease, which leads to hyperproliferation of the oesophageal mucosa. Also, ethanol-associated immune suppression may facilitate tumour cell spread [65].

3. Justification for recommendation

The consumption of alcoholic beverages is causally associated with cancers of the oral cavity, pharynx, larynx, oesophagus, liver, colorectum and female breast. Risk estimates of alcohol-related cancer incidence and mortality emphasise a dose-dependent relationship that does not vary by beverage type and does not show a threshold of intake, since the adverse effect is observed even at consumption of less than one alcoholic drink per day [12,75]. Ethanol carcinogenesis may not uniformly target tissues and may depend on drinking patterns and timing of exposure, as based on observed differences in cancer risk across organ sites.

Taking all these issues into account, the ECAC has developed the following recommendation:

“If you drink alcohol of any type, limit your intake. Not drinking alcohol is better for cancer prevention.”

3.1. Alcohol-attributable burden of disease

Worldwide, alcohol consumption is the second leading risk factor after tobacco in terms of morbidity, and it accounts for a significant number of deaths, particularly in high-income countries. In Europe, alcohol is the third leading risk factor for disease and mortality after tobacco and high blood pressure [76]. Alcohol is causally associated with about 60 types of disease, including heart disease, stroke and vascular diseases, as well as liver cirrhosis and cancer. Fetal exposure to alcohol strongly increases the risk of birth defects and intellectual impairments. Alcohol-use disorders (e.g.,

acute alcohol intoxication, alcohol-dependence syndrome) account for a significant share of total morbidity, particularly among men (44.5%) [1]. Alcohol also contributes to death and disability through accidents and injuries, assault, violence, homicide and suicide.

In the EU in 2004, almost 95,000 men and more than 25,000 women, aged 15–64, died prematurely of alcohol-attributable causes. This means that one in seven male and one in 13 female premature deaths in this age category were caused by alcohol [76]. In men in 2004, approximately two in five deaths were due to liver cirrhosis, one in three to injuries, and one in five (15.9%) to cancer. In women, more than two thirds of alcohol-attributable deaths arose from liver cirrhosis and cancer (30.7%; the largest proportion of which, 21%, involves breast cancer), with cardiovascular disease other than ischaemic heart disease as a distant third cause.

Almost 90% of the burden of disease attributable to alcohol is caused by heavy drinking, both regular and irregular. This has important implications for prevention and alcohol policy: any measure which aims to successfully reduce alcohol-attributable harm has to involve cutting down regular and irregular heavy drinking occasions.

3.2. Public policies on alcohol taxation as measures of prevention

Negative health effects and social harm caused by detrimental alcohol consumption can be reduced through effective prevention strategies. In 1979, the World Health Assembly called upon World Health Organization (WHO) member states to develop and adopt appropriate legislation and measures to tackle alcohol misuse [77]. Such efforts culminated with the endorsement, in 2010, of the global strategy on the harmful use of alcohol [78] that supports ten target areas for national actions, including: health sector response, community actions, drink-driving policies, limitation of the availability of alcohol, action on marketing and pricing policies, reducing the negative consequences of intoxication and reducing the public health effect of illegally and informally produced alcohol. Based on four key criteria (namely effectiveness, affordability, efficiency and acceptability/feasibility), the WHO has more recently identified restricted access to retail alcohol, limitation of alcohol advertising, and taxes on alcohol as the best-performing interventions to tackle harmful alcohol use [79]. Both the enforcement of drink-driving laws and brief advice for excessive drinkers are valuable interventions, although their efficiency in some settings may be less attractive.

In particular, alcohol prices – and consequently affordability – are a strong determinant of alcohol consumption and can be altered by using taxes or direct price controls, including minimum price policies [80]. However, public policies on alcohol taxation are not always homogeneous across the EU. Countries with stricter and more comprehensive alcohol policies generally have lower levels of alcohol consumption. Observational studies suggest that the effects of taxation would be larger for moderate than heavy drinkers [81], women [82], and young consumers [83]. On average, a 10% increase in the price of alcohol is correlated with a 4.4% decrease in consumption, with some variability depending on the type of alcoholic beverage [84]. Sadly, between 1996 and 2004 affordability has increased by 50% or more in Estonia, the United Kingdom, Czech Republic, Finland and Ireland, and by a smaller amount in Portugal, Spain, Denmark, Sweden, France, Austria, the Netherlands, Greece, Germany, Poland and Belgium – in particular for young people [85]. Time-series analyses in Canada (British Columbia) showed that a 10% increase in the minimum price for alcohol reduces the consumption of spirits by 6.8%, of wine by 8.9%, of alcopops (flavoured, often sweet, alcoholic beverages) by 13.9%, and of beer by 1.5% [86]. The most common

approach in European countries is based on a combination of excise duty and value-added taxes, which should account for potential effects of the type of beverage consumed. As an example, the introduction of a tax on alcopops in Germany simply shifted consumption from spirit-based to beer-based beverages [87].

4. Conclusion

Worldwide, alcohol consumption is the second leading risk factor in terms of morbidity, and it accounts for a significant number of deaths, particularly in high-income countries. In Europe, it has been estimated that about 10% (95%CI: 7–13%) of all cancer cases in men and 3% (95%CI: 1–5%) of all cancer cases in women are attributable to alcohol consumption [12]. There is strong evidence that people can reduce their risk of cancer by limiting or cutting their consumption of alcoholic drinks. Overall, in European populations the risk of cancer in men who consume less than two alcoholic drinks (less than 20 g of pure alcohol) per day and in women who consume less than one alcoholic drink (less than 10 g of pure alcohol) per day is 6% lower (hazard ratio = 0.94; 95%CI: 0.90–0.96) than that in people with higher alcohol intakes [88]. On average, reducing the consumption from four or more to one or less alcoholic drinks per day may reduce the risk of liver cancer by 21% [48], the risk of colorectal cancer by 31% [43], and the risk of female breast cancer by 30% [48]. The WHO have stated clearly that increasing price, reducing availability and banning advertising are the most effective policy measures that should be implemented at the European level in order to decrease the alcohol-related burden of disease.

European Code Against Cancer.:

EUROPEAN CODE AGAINST CANCER:
<p>12 ways to reduce your cancer risk</p> <ol style="list-style-type: none"> 1. Do not smoke. Do not use any form of tobacco 2. Make your home smoke free. Support smoke-free policies in your workplace 3. Take action to be a healthy body weight 4. Be physically active in everyday life. Limit the time you spend sitting 5. Have a healthy diet: <ul style="list-style-type: none"> • Eat plenty of whole grains, pulses, vegetables and fruits • Limit high-calorie foods (foods high in sugar or fat) and avoid sugary drinks • Avoid processed meat; limit red meat and foods high in salt 6. If you drink alcohol of any type, limit your intake. Not drinking alcohol is better for cancer prevention 7. Avoid too much sun, especially for children. Use sun protection. Do not use sunbeds 8. In the workplace, protect yourself against cancer-causing substances by following health and safety instructions 9. Find out if you are exposed to radiation from naturally high radon levels in your home; take action to reduce high radon levels 10. For women: <ul style="list-style-type: none"> • Breastfeeding reduces the mother's cancer risk. If you can, breastfeed your baby • Hormone replacement therapy (HRT) increases the risk of certain cancers. Limit use of HRT

11. Ensure your children take part in vaccination programmes for:
 - Hepatitis B (for newborns)
 - Human papillomavirus (HPV) (for girls)
12. Take part in organised cancer screening programmes for:
 - Bowel cancer (men and women)
 - Breast cancer (women)
 - Cervical cancer (women)

The European Code Against Cancer focuses on actions that individual citizens can take to help prevent cancer. Successful cancer prevention requires these individual actions to be supported by governmental policies and actions.

The resulting recommendation of the 4th edition of the ECAC targeted to individuals is to limit or cut the consumption of alcoholic beverages.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgments

The European Code against Cancer project was co-funded by the European Union [grant agreements No. 2010 53 04 and No. 2011 53 05] and the International Agency for Research on Cancer. The authors alone are responsible for the views expressed in this manuscript.

References

- [1] World Health Organization. Global Health Risks: Mortality and Burden of Disease Attributable to Selected Major Risks. Geneva: World Health Organization, 2009.
- [2] IARC Working Group on the Evaluation of Carcinogenic Risks to Human. Personal Habits and Indoor Combustions. Volume 100 E. A Review of Human Carcinogens, 100. 2012.
- [3] Coglianò VJ, Baan R, Straif K, Grosse Y, Lauby-Secretan B, El Ghissassi F, et al. Preventable exposures associated with human cancers. *J. Natl. Cancer Inst.* 2011;103(24):1827–39.
- [4] Allen NE, Beral V, Casabonne D, Kan SW, Reeves GK, Brown A, et al. Moderate alcohol intake and cancer incidence in women. *J. Natl. Cancer Inst.* 2009;101(5):296–305.
- [5] Newcomb PA, Nichols HB, Beasley JM, Egan K, Titus-Ernstoff L, Hampton JM, et al. No difference between red wine or white wine consumption and breast cancer risk. *Cancer Epidemiol. Biomarkers Prev.* 2009;18(3):1007–10.
- [6] Marron M, Boffetta P, Moller H, Ahrens W, Pohlabein H, Benhamou S, et al. Risk of upper aerodigestive tract cancer and type of alcoholic beverage: a European multicenter case-control study. *Eur. J. Epidemiol.* 2012;27(7):499–517.
- [7] Boyle P, Autier P, Bartelink H, Baselga J, Boffetta P, Burn J, et al. European Code Against Cancer and scientific justification: third version (2003). *Ann. Oncol.* 2003;14(7):973–1005.
- [8] OECD. Health at a Glance: Europe 2012. OECD Publishing; 2012.
- [9] Kanny D, Liu Y, Brewer RD. Binge drinking - United States, 2009. *MMWR Surveill Summ.* 2011;60 Suppl:101–4.
- [10] Bloomfield K, Grittner U, Kramer S, Gmel G. Social inequalities in alcohol consumption and alcohol-related problems in the study countries of the EU concerted action 'Gender, Culture and Alcohol Problems: a Multi-national Study'. *Alcohol Alcohol Suppl.* 2006;41(1):i26–36.
- [11] Grittner U, Kuntsche S, Graham K, Bloomfield K. Social inequalities and gender differences in the experience of alcohol-related problems. *Alcohol Alcohol* 2012;47(5):597–605.
- [12] Devaux M, Sassi F. Alcohol Consumption and Harmful Drinking: Trends and Social Disparities in OECD Countries. OECD Publishing; 2013.
- [13] Polednak AP. Recent trends in incidence rates for selected alcohol-related cancers in the United States. *Alcohol Alcohol* 2005;40(3):234–8.
- [14] MacArthur GJ, Smith MC, Melotti R, Heron J, Macleod J, Hickman M, et al. Patterns of alcohol use and multiple risk behaviour by gender during early and late adolescence: the ALSPAC cohort. *J. Public Health (Oxf)* 2012;34 Suppl. 1:i20–30.

- [15] Rebholz CE, Kuehni CE, Strippoli MP, Rueegg CS, Michel G, Hengartner H, et al. Alcohol consumption and binge drinking in young adult childhood cancer survivors. *Pediatr. Blood Cancer* 2012;58(2):256–64.
- [16] Anderson P, Baumberg B. Alcohol in Europe. A Public Health Perspective. A Report for the European Commission. London: HAPI, 2006.
- [17] Fillmore KM, Stockwell T, Chikritzhs T, Bostrom A, Kerr W. Moderate alcohol use and reduced mortality risk: systematic error in prospective studies and new hypotheses. *Ann. Epidemiol.* 2007;17(5 Suppl.):S16–23.
- [18] Gunzerath L, Faden V, Zakhari S, Warren K. National Institute on Alcohol Abuse and Alcoholism report on moderate drinking. *Alcohol Clin. Exp. Res.* 2004;28(6):829–47.
- [19] Hvidtfeldt UA, Tolstrup JS, Jakobsen MU, Heitmann BL, Gronbaek M, O'Reilly E, et al. Alcohol intake and risk of coronary heart disease in younger, middle-aged, and older adults. *Circulation* 2010;121(14):1589–97.
- [20] Jimenez M, Chiuev SE, Glynn RJ, Stampfer MJ, Camargo Jr CA, Willett WC, et al. Alcohol consumption and risk of stroke in women. *Stroke* 2012;43(4):939–45.
- [21] Ronskley PE, Brien SE, Turner BJ, Mukamal KJ, Ghali WA. Association of alcohol consumption with selected cardiovascular disease outcomes: a systematic review and meta-analysis. *BMJ* 2011;342:d671.
- [22] Klatsky AL, Armstrong MA, Friedman GD. Risk of cardiovascular mortality in alcohol drinkers, ex-drinkers and nondrinkers. *Am. J. Cardiol.* 1990;66(17):1237–42.
- [23] Imhof A, Koenig W. Alcohol inflammation and coronary heart disease. *Addict. Biol.* 2003;8(3):271–7.
- [24] Rehm J, Room R, Graham K, Monteiro M, Gmel G, Sempos CT. The relationship of average volume of alcohol consumption and patterns of drinking to burden of disease: an overview. *Addiction* 2003;98(9):1209–28.
- [25] Britton A, Marmot M. Different measures of alcohol consumption and risk of coronary heart disease and all-cause mortality: 11-year follow-up of the Whitehall II Cohort Study. *Addiction* 2004;99(1):109–16.
- [26] Trevisan M, Dorn J, Falkner K, Russell M, Ram M, Muti P, et al. Drinking pattern and risk of non-fatal myocardial infarction: a population-based case-control study. *Addiction* 2004;99(3):313–22.
- [27] Schutze M, Boeing H, Pischon T, Rehm J, Kehoe T, Gmel G, et al. Alcohol attributable burden of incidence of cancer in eight European countries based on results from prospective cohort study. *BMJ* 2011;342:d1584.
- [28] Friberg JT, Yuan JM, Wang R, Koh WP, Lee HP, Yu MC. A prospective study of tobacco and alcohol use as risk factors for pharyngeal carcinomas in Singapore Chinese. *Cancer* 2007;109(6):1183–91.
- [29] Muwonge R, Ramadas K, Sankila R, Thara S, Thomas G, Vinoda J, et al. Role of tobacco smoking, chewing and alcohol drinking in the risk of oral cancer in Trivandrum, India: a nested case-control design using incident cancer cases. *Oral Oncol.* 2008;44(5):446–54.
- [30] Rota M, Bellocco R, Scotti L, Tramacere I, Jenab M, Corrao G, et al. Random-effects meta-regression models for studying nonlinear dose-response relationship, with an application to alcohol and esophageal squamous cell carcinoma. *Stat. Med.* 2010;29(26):2679–87.
- [31] Islami F, Tramacere I, Rota M, Bagnardi V, Fedirko V, Scotti L, et al. Alcohol drinking and laryngeal cancer: overall and dose-risk relation—a systematic review and meta-analysis. *Oral Oncol.* 2010;46(11):802–10.
- [32] Islami F, Fedirko V, Tramacere I, Bagnardi V, Jenab M, Scotti L, et al. Alcohol drinking and esophageal squamous cell carcinoma with focus on light-drinkers and never-smokers: a systematic review and meta-analysis. *Int. J. Cancer* 2011;129(10):2473–84.
- [33] Turati F, Garavello W, Tramacere I, Pelucchi C, Galeone C, Bagnardi V, et al. A meta-analysis of alcohol drinking and oral and pharyngeal cancers: results from subgroup analyses. *Alcohol Alcohol* 2013;48(1):107–18.
- [34] Hashibe M, Brennan P, Chuang SC, Boccia S, Castellsague X, Chen C, et al. Interaction between tobacco and alcohol use and the risk of head and neck cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *Cancer Epidemiol. Biomarkers Prev.* 2009;18(2):541–50.
- [35] Pelucchi C, Tramacere I, Boffetta P, Negri E, La Vecchia C. Alcohol consumption and cancer risk. *Nutr. Cancer* 2011;63(7):983–90.
- [36] Ahmad, Kiadaliri A, Jarl J, Gavrilidis G, Gerdtham UG. Alcohol drinking cessation and the risk of laryngeal and pharyngeal cancers: a systematic review and meta-analysis. *PLoS One* 2013;8(3):e58158.
- [37] Altieri A, Bosetti C, Talamini R, Gallus S, Franceschi S, Levi F, et al. Cessation of smoking and drinking and the risk of laryngeal cancer. *Br. J. Cancer* 2002;87(11):1227–9.
- [38] Castellsague X, Quintana MJ, Martinez MC, Nieto A, Sanchez MJ, Juan A, et al. The role of type of tobacco and type of alcoholic beverage in oral carcinogenesis. *Int. J. Cancer* 2004;108(5):741–9.
- [39] Jarl J, Gerdtham UG. Time pattern of reduction in risk of oesophageal cancer following alcohol cessation—a meta-analysis. *Addiction* 2012;107(7):1234–43.
- [40] Rehm J, Patra J, Popova S. Alcohol drinking cessation and its effect on esophageal and head and neck cancers: a pooled analysis. *Int. J. Cancer* 2007;121(5):1132–7.
- [41] World Cancer Research Fund/American Institute for Cancer Research. Continuous Update Project Report. Food, Nutrition Physical Activity and the Prevention of Colorectal Cancer; 2010.
- [42] Cho EL, Lee JE, Rimm EB, Fuchs CS, Giovannucci EL. Alcohol consumption and the risk of colon cancer by family history of colorectal cancer. *Am. J. Clin. Nutr.* 2012;95(2):413–9.
- [43] Fedirko V, Tramacere I, Bagnardi V, Rota M, Scotti L, Islami F, et al. Alcohol drinking and colorectal cancer risk: an overall and dose-response meta-analysis of published studies. *Ann. Oncol.* 2011;22(9):1958–72.
- [44] Bagnardi V, Rota M, Botteri E, Tramacere I, Islami F, Fedirko V, et al. Light alcohol drinking and cancer: a meta-analysis. *Ann. Oncol.* 2013;24(2):301–8.
- [45] Huxley RR, nsary-Moghaddam A, Clifton P, Czernichow S, Parr CL, Woodward M. The impact of dietary and lifestyle risk factors on risk of colorectal cancer: a quantitative overview of the epidemiological evidence. *Int. J. Cancer* 2009;125(1):171–80.
- [46] Hassan MM, Spitz MR, Thomas MB, El-Deeb AS, Glover KY, Nguyen NT, et al. Effect of different types of smoking and synergism with hepatitis C virus on risk of hepatocellular carcinoma in American men and women: case-control study. *Int. J. Cancer* 2008;123(8):1883–91.
- [47] Shih WL, Chang HC, Liaw YF, Lin SM, Lee SD, Chen PJ, et al. Influences of tobacco and alcohol use on hepatocellular carcinoma survival. *Int. J. Cancer* 2012;131(11):2612–21.
- [48] Corrao G, Bagnardi V, Zambon A, La Vecchia C. A meta-analysis of alcohol consumption and the risk of 15 diseases. *Prev. Med.* 2004;38(5):613–9.
- [49] Gwack J, Hwang SS, Ko KP, Jun JK, Park SK, Chang SH, et al. Fasting serum glucose and subsequent liver cancer risk in a Korean prospective cohort. *J. Prev. Med. Public Health* 2007;40(1):23–8.
- [50] World Cancer Research Fund/American Institute for Cancer Research. Continuous Update Project Report. Food, Nutrition, Physical Activity and the Prevention of Breast Cancer; 2008.
- [51] Chen WY, Rosner B, Hankinson SE, Colditz GA, Willett WC. Moderate alcohol consumption during adult life, drinking patterns, and breast cancer risk. *JAMA* 2011;306(17):1884–90.
- [52] Jeffreys M, McKenzie F, Firestone R, Gray M, Cheng S, Moala A, et al. A multi-ethnic breast cancer case-control study in New Zealand: evidence of differential risk patterns. *Cancer Causes Control* 2013;24(1):135–52.
- [53] Berkey CS, Willett WC, Frazier AL, Rosner B, Tamimi RM, Rockett HR, et al. Prospective study of adolescent alcohol consumption and risk of benign breast disease in young women. *Pediatrics* 2010;125(5):e1081–87.
- [54] Liu Y, Tamimi RM, Berkey CS, Willett WC, Collins LC, Schnitt SJ, et al. Intakes of alcohol and folate during adolescence and risk of proliferative benign breast disease. *Pediatrics* 2012;129(5):e1192–98.
- [55] Tjonneland A, Christensen J, Thomsen BL, Olsen A, Stripp C, Overvad K, et al. Lifetime alcohol consumption and postmenopausal breast cancer rate in Denmark: a prospective cohort study. *J. Nutr.* 2004;134(1):173–8.
- [56] Brooks PJ, Zakhari S. Moderate alcohol consumption and breast cancer in women: from epidemiology to mechanisms and interventions. *Alcohol Clin. Exp. Res.* 2013;37(1):23–30.
- [57] Lachenmeier DW, Monakhova YB. Short-term salivary acetaldehyde increase due to direct exposure to alcoholic beverages as an additional cancer risk factor beyond ethanol metabolism. *J. Exp. Clin. Cancer Res.* 2011;30:3.
- [58] Ferreira Antunes JL, Toporcov TN, Biazevic MG, Boing AF, Scully C, Petti S. Joint and independent effects of alcohol drinking and tobacco smoking on oral cancer: a large case-control study. *PLoS One* 2013;8(7):e68132.
- [59] Hashibe M, Brennan P, Benhamou S, Castellsague X, Chen C, Curado MP, et al. Alcohol drinking in never users of tobacco, cigarette smoking in never drinkers, and the risk of head and neck cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *J. Natl. Cancer Inst.* 2007;99(10):777–89.
- [60] Zaridze D, Lewington S, Boroda A, Scelo G, Karpov R, Lazarev A, et al. Alcohol and mortality in Russia: prospective observational study of 151 000 adults. *Lancet* 2014 Jan 30.
- [61] Guo H, Zhang G, Mai R. Alcohol dehydrogenase-1B Arg47His polymorphism and upper aerodigestive tract cancer risk: a meta-analysis including 24,252 subjects. *Alcohol Clin. Exp. Res.* 2012;36(2):272–8.
- [62] Brennan P, Lewis S, Hashibe M, Bell DA, Boffetta P, Bouchardy C, et al. Pooled analysis of alcohol dehydrogenase genotypes and head and neck cancer: a HuGE review. *Am. J. Epidemiol.* 2004;159(1):1–16.
- [63] Cadoni G, Boccia S, Petrelli L, Di Giannantonio P, Arzani D, Giorgio A, et al. A review of genetic epidemiology of head and neck cancer related to polymorphisms in metabolic genes, cell cycle control and alcohol metabolism. *Acta Otorhinolaryngol. Ital.* 2012;32(1):1–11.
- [64] Chang JS, Straif K, Guha N. The role of alcohol dehydrogenase genes in head and neck cancers: a systematic review and meta-analysis of ADH1B and ADH1C. *Mutagenesis* 2012;27(3):275–86.
- [65] Seitz HK, Stickel F. Molecular mechanisms of alcohol-mediated carcinogenesis. *Nat. Rev. Cancer* 2007;7(8):599–612.
- [66] Kim J, Cho YA, Kim DH, Lee BH, Hwang DY, Jeong J, et al. Dietary intake of folate and alcohol MTHFR C677 T polymorphism, and colorectal cancer risk in Korea. *Am. J. Clin. Nutr.* 2012;95(2):405–12.
- [67] Platek ME, Shields PG, Marian C, McCann SE, Bonner MR, Nie J, et al. Alcohol consumption and genetic variation in methylenetetrahydrofolate reductase and 5-methyltetrahydrofolate-homocysteine methyltransferase in relation to breast cancer risk. *Cancer Epidemiol. Biomarkers Prev.* 2009;18(9):2453–9.
- [68] Alshatwi AA. Breast cancer risk, dietary intake, and methylenetetrahydrofolate reductase (MTHFR) single nucleotide polymorphisms. *Food Chem. Toxicol.* 2010;48(7):1881–5.
- [69] Supic G, Jovic N, Kozomara R, Zeljic K, Magic Z. Interaction between the MTHFR C677 T polymorphism and alcohol—impact on oral cancer risk and multiple DNA methylation of tumor-related genes. *J. Dent. Res.* 2011;90(1):65–70.
- [70] Fernandez SV. Estrogen, alcohol consumption, and breast cancer. *Alcohol Clin. Exp. Res.* 2011;35(3):389–91.
- [71] Zhang SM, Lee IM, Manson JE, Cook NR, Willett WC, Buring JE. Alcohol consumption and breast cancer risk in the Women's Health Study. *Am. J. Epidemiol.* 2007;165(6):667–76.

- [72] Lew JQ, Freedman ND, Leitzmann MF, Brinton LA, Hoover RN, Hollenbeck AR, et al. Alcohol and risk of breast cancer by histologic type and hormone receptor status in postmenopausal women: the NIH-AARP Diet and Health Study. *Am. J. Epidemiol.* 2009;170(3):308–17.
- [73] Li CI, Chlebowski RT, Freiberg M, Johnson KC, Kuller L, Lane D, et al. Alcohol consumption and risk of postmenopausal breast cancer by subtype: the women's health initiative observational study. *J. Natl. Cancer Inst.* 2010;102(18):1422–31.
- [74] Suzuki R, Orsini N, Mignone L, Saji S, Wolk A. Alcohol intake and risk of breast cancer defined by estrogen and progesterone receptor status—a meta-analysis of epidemiological studies. *Int. J. Cancer* 2008;122(8):1832–41.
- [75] Nelson DE, Jarman DW, Rehm J, Greenfield TK, Rey G, Kerr WC, et al. Alcohol-attributable cancer deaths and years of potential life lost in the United States. *Am. J. Public Health* 2013;103(4):641–8.
- [76] Anderson P, Moller L, Galea G. Alcohol in the European Union Consumption, Harm and Policy Approaches. London: HAPI, 2012.
- [77] Thirty-ninth World Health Assembly Evaluates Progress towards the Goal of Health for all. Resolution WHA32.40. Development of the WHO programme on alcohol-related problems. World Health Assembly. Geneva, 1986.
- [78] World Health Organization. Global Status Report on Alcohol and Health; 2011.
- [79] World Health Organization. A Comprehensive Global Monitoring Framework and Voluntary Global Targets for the Prevention and Control of NCDs. Geneva; 2011.
- [80] Wall M, Casswell S. Affordability of alcohol as a key driver of alcohol demand in New Zealand: a co-integration analysis. *Addiction* 2013;108(1):72–9.
- [81] Manning WG, Blumberg L, Moulton LH. The demand for alcohol: the differential response to price. *J. Health Econ.* 1995;14(2):123–48.
- [82] Elder RW, Lawrence B, Ferguson A, Naimi TS, Brewer RD, Chattopadhyay SK, et al. The effectiveness of tax policy interventions for reducing excessive alcohol consumption and related harms. *Am. J. Prev. Med.* 2010;38(2):217–29.
- [83] Xu X, Chaloupka FJ. The effects of prices on alcohol use and its consequences. *Alcohol Res. Health* 2011;34(2):236–45.
- [84] Wagenaar AC, Salois MJ, Komro KA. Effects of beverage alcohol price and tax levels on drinking: a meta-analysis of 1003 estimates from 112 studies. *Addiction* 2009;104(2):179–90.
- [85] Rabinovich L, Brutscher P, de Vries H, Tiessen J, Clift J, Reding A. The Affordability of Alcoholic Beverages in the European Union - Understanding the Link between Alcohol Affordability, Consumption and Harms; 2009.
- [86] Stockwell T, Auld MC, Zhao J, Martin G. Does minimum pricing reduce alcohol consumption? The experience of a Canadian province. *Addiction* 2012;107(5):912–20.
- [87] Anderson P, Suhrcke M, Brookes C. An overview of the Market for Alcohol Beverages of Potentially Particular Appeal to Minors. London: HAPI, 2012.
- [88] Romaguera D, Vergnaud AC, Peeters PH, van Gils CH, Chan DS, Ferrari P, et al. Is concordance with World Cancer Research Fund/American Institute for Cancer Research guidelines for cancer prevention related to subsequent risk of cancer? Results from the EPIC study. *Am. J. Clin. Nutr.* 2012;96(1):150–63.