

Multiple Flag Varieties of Finite Type

Peter Magyar, Jerzy Weyman, and Andrei Zelevinsky*

Department of Mathematics, Northeastern University, Boston, Massachusetts 02115

Received April 9, 1998; accepted April 30, 1998

We classify all products of flag varieties with finitely many orbits under the diagonal action of the general linear group. We also classify the orbits in each case and construct explicit representatives. © 1999 Academic Press

1. INTRODUCTION

For a reductive group G , the Schubert (or Bruhat) decomposition describes the orbits of a Borel subgroup B acting on the flag variety G/B . It is the starting point for analyzing the geometry and topology of G/B , and is also significant for the representation theory of G . This decomposition states that $G/B = \coprod_{w \in W} B \cdot wB$, where W is the Weyl group. An equivalent form which is more symmetric is the G -orbit decomposition of the double flag variety: $(G/B)^2 = \coprod_{w \in W} G \cdot (eB, wB)$. One can easily generalize the Schubert decomposition by considering G -orbits on a product of two *partial* flag varieties $G/P \times G/Q$, where P and Q are parabolic subgroups. The crucial feature in each case is that the number of orbits is *finite* and has a rich combinatorial structure.

Here we address the more general question: for which tuples of parabolic subgroups (P_1, \dots, P_k) does the group G have finitely many orbits when acting diagonally in the product of several flag varieties $G/P_1 \times \dots \times G/P_k$? As before, this is equivalent to asking when $G/P_2 \times \dots \times G/P_k$ has finitely many P_1 -orbits. (If $P_1 = B$ is a Borel subgroup, this is one definition of a spherical variety. Thus, our problem includes that of classifying the multiple flag varieties of spherical type.) To the best of our knowledge, the problem of classifying all finite-orbit tuples for an arbitrary G is still open (although in the special case when $k = 3$, $P_1 = B$, and P_2 and P_3 are maximal parabolic subgroups, such a classification was given in [6]).

* The research of Peter Magyar, Jerzy Weyman, and Andrei Zelevinsky was supported in part by an NSF Postdoctoral Fellowship and NSF Grants DMS-9700884 and DMS-9625511, respectively.

In this paper, we present a complete solution of the classification problem for $G = GL_n$. For this case, the partial flag varieties G/P consist of all flags of subspaces with some fixed dimensions in an n -dimensional vector space V . Our classification theorem (Theorem 2.2) provides the list of all dimension types such that GL_n has finitely many orbits in the corresponding product of flag varieties. We also classify the orbits in each case and construct explicit representatives (standard forms). Precise formulations of the main results will be given in the next section; the proofs are given in Sections 3 and 4. In Section 4.3 we also discuss some partial results on the generalized Bruhat order given by adjacency of orbits.

We use results and ideas from the theory of quiver representations. In fact, our key criterion for finite type (Proposition 3.3 below) is very close to (but distinct from) the characterization of quiver representations of finite type due to V. Kac [5].

2. MAIN RESULTS

2.1. Classification theorem. Let $\mathbf{a} = (a_1, \dots, a_p)$ be a nonnegative list of integers with sum equal to n . We call such a list a *composition* of n , and a_1, \dots, a_p the *parts* of \mathbf{a} . Throughout this paper, all vector spaces are over a fixed algebraically closed field. For a vector space V of dimension n , we denote by $\text{Fl}_{\mathbf{a}}(V)$ the variety of flags $A = (0 = A_0 \subset A_1 \subset \dots \subset A_p = V)$ of vector subspaces in V such that

$$\dim(A_i/A_{i-1}) = a_i \quad (i = 1, \dots, p).$$

A tuple of compositions $(\mathbf{a}_1, \dots, \mathbf{a}_k)$ of the same number n is said to be of *finite type* if the group $GL(V)$ (acting diagonally) has finitely many orbits in the multiple flag variety $\text{Fl}_{\mathbf{a}_1}(V) \times \dots \times \text{Fl}_{\mathbf{a}_k}(V)$. We say that a composition is *trivial* if it has only one non-zero part n . Then the corresponding flag variety consists of a single point, so adding any number of trivial compositions to a tuple gives essentially the same multiple flag variety, and does not affect the finite type property.

THEOREM 2.1. *If a tuple of non-trivial compositions $(\mathbf{a}_1, \dots, \mathbf{a}_k)$ is of finite type then $k \leq 3$.*

In other words, a multiple flag variety of finite type cannot have more than 3 non-trivial factors. Thus any tuple of compositions of finite type can be made into a triple by adding or removing trivial compositions, and we need only classify *triples* of finite type. We will write $(\mathbf{a}, \mathbf{b}, \mathbf{c})$ instead of $(\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3)$. We denote by $\min(\mathbf{a})$ the minimum of the non-zero parts of a composition \mathbf{a} . Now we can formulate our first main theorem.

THEOREM 2.2 *Let \mathbf{a} , \mathbf{b} and \mathbf{c} be compositions and let p , q and r denote their respective numbers of non-zero parts. Assume without loss of generality that $p \leq q \leq r$. Then the triple $(\mathbf{a}, \mathbf{b}, \mathbf{c})$ is of finite type if and only if it belongs to one of the following classes:*

- $(A_{q,r}) \quad (p, q, r) = (1, q, r), 1 \leq q \leq r.$
- $(D_{r+2}) \quad (p, q, r) = (2, 2, r), 2 \leq r.$
- $(E_6) \quad (p, q, r) = (2, 3, 3).$
- $(E_7) \quad (p, q, r) = (2, 3, 4).$
- $(E_8) \quad (p, q, r) = (2, 3, 5).$
- $(E_{r+3}^{(a)}) \quad (p, q, r) = (2, 3, r), 3 \leq r, \min(\mathbf{a}) = 2.$
- $(E_{r+3}^{(b)}) \quad (p, q, r) = (2, 3, r), 3 \leq r, \min(\mathbf{b}) = 1.$
- $(S_{q,r}) \quad (p, q, r) = (2, q, r), 2 \leq q \leq r, \min(\mathbf{a}) = 1.$

The types in Theorem 2.2 have some obvious overlaps. The type $A_{q,r}$ covers all multiple flag varieties with less than three non-trivial factors. The type $S_{q,r}$ appeared in Brion [4]. For relations with the classification of quiver representations of finite type due to V. Kac [5], see Remark 3.4.

Note that, for each of the first five types in Theorem 2.2, there are no restrictions on the dimensions of subspaces in the corresponding flag varieties; only the last three types $E^{(a)}$, $E^{(b)}$ and S involve such restrictions. The first five types are naturally related to the simply-laced Dynkin graphs (as suggested by their names). Let $T = T_{p,q,r}$ denote the graph with $p + q + r - 2$ vertices that consists of 3 chains with p , q , and r vertices, joined together at a common endpoint. We see that the cases in our classification with no restrictions on dimensions are precisely those for which T is one of the Dynkin graphs A_n , D_n , E_6 , E_7 , or E_8 . This is of course no coincidence: we will see that this part of our classification is equivalent to Gabriel’s classification of quivers of finite type (and follows from the Cartan–Killing classification of graphs that give rise to positive-definite quadratic forms).

2.2. Classification of orbits

Now we describe a combinatorial parametrization of the set of $GL(V)$ -orbits in $\text{Fl}_{\mathbf{a}}(V) \times \text{Fl}_{\mathbf{b}}(V) \times \text{Fl}_{\mathbf{c}}(V)$ for any triple $(\mathbf{a}, \mathbf{b}, \mathbf{c})$ of compositions of finite type. For a composition $\mathbf{a} = (a_1, \dots, a_p)$, we write

$$|\mathbf{a}| = a_1 + \dots + a_p, \quad \|\mathbf{a}\|^2 = a_1^2 + \dots + a_p^2;$$

the number p of parts of \mathbf{a} will be denoted $\ell(\mathbf{a})$ and called the *length* of \mathbf{a} .

For any positive integers p, q , and r , let $A_{p, q, r}$ denote the additive semigroup of all triples of compositions $(\mathbf{a}, \mathbf{b}, \mathbf{c})$ such that $(\ell(\mathbf{a}), \ell(\mathbf{b}), \ell(\mathbf{c})) = (p, q, r)$, and $|\mathbf{a}| = |\mathbf{b}| = |\mathbf{c}|$. (Here, in contrast to the notation of Theorem 2.2, the numbers p, q, r include the zero parts of $\mathbf{a}, \mathbf{b}, \mathbf{c}$.) For every $(\mathbf{a}, \mathbf{b}, \mathbf{c}) \in A_{p, q, r}$, we set

$$Q(\mathbf{a}, \mathbf{b}, \mathbf{c}) = \dim GL(V) - \dim \mathrm{Fl}_{\mathbf{a}}(V) - \dim \mathrm{Fl}_{\mathbf{b}}(V) - \dim \mathrm{Fl}_{\mathbf{c}}(V),$$

where V is a vector space of dimension $n = |\mathbf{a}| = |\mathbf{b}| = |\mathbf{c}|$. An easy calculation shows that

$$Q(\mathbf{a}, \mathbf{b}, \mathbf{c}) = \frac{1}{2}(\|\mathbf{a}\|^2 + \|\mathbf{b}\|^2 + \|\mathbf{c}\|^2 - n^2). \quad (2.1)$$

The function Q is called the *Tits quadratic form*.

Let $\Pi_{p, q, r}$ denote the set of all triples $\mathbf{d} = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in A_{p, q, r}$ of finite type such that $Q(\mathbf{d}) = 1$.

THEOREM 2.3. *Let $(\mathbf{a}, \mathbf{b}, \mathbf{c}) \in A_{p, q, r}$ be a triple of compositions of finite type. Then $GL(V)$ -orbits in $\mathrm{Fl}_{\mathbf{a}}(V) \times \mathrm{Fl}_{\mathbf{b}}(V) \times \mathrm{Fl}_{\mathbf{c}}(V)$ are in natural bijection with families of nonnegative integers $M = (m_{\mathbf{d}})$ indexed by $\mathbf{d} \in \Pi_{p, q, r}$ such that, in the semigroup $A_{p, q, r}$,*

$$\sum_{\mathbf{d} \in \Pi_{p, q, r}} m_{\mathbf{d}} \mathbf{d} = (\mathbf{a}, \mathbf{b}, \mathbf{c}).$$

The set $\Pi_{p, q, r}$ can be explicitly described as follows. For a composition \mathbf{a} , we denote by \mathbf{a}^+ the partition obtained from \mathbf{a} by removing all zero parts and rearranging the non-zero parts in weakly decreasing order. (For example, if $\mathbf{a} = (0, 2, 1, 0, 3, 2)$ then $\mathbf{a}^+ = (3, 2, 2, 1)$.) We denote $(a^p) = \underbrace{(a, \dots, a)}_{p \text{ parts}}$.

THEOREM 2.4. *A triple $(\mathbf{a}, \mathbf{b}, \mathbf{c}) \in A_{p, q, r}$ belongs to $\Pi_{p, q, r}$ if and only if the (unordered) triple of partitions $\{\mathbf{a}^+, \mathbf{b}^+, \mathbf{c}^+\}$ is one of the following:*

$$\begin{aligned} & \{(1), (1), (1)\}, \quad \{(3^2), (2^3), (2, 1, 1, 1, 1)\}, \quad \{(4, 2), (2^3), (1^6)\}, \\ & \{(m+1, m), (m, m, 1), (1^{2m+1})\} \quad (m \geq 2), \\ & \{(m, m), (m, m-1, 1), (1^{2m})\} \quad (m \geq 2), \\ & \{(n-1, 1), (1^n), (1^n)\} \quad (n \geq 2). \end{aligned}$$

Remark 2.5. Except for $((3^2), (2^3), (2, 1, 1, 1, 1))$, all of the triples in Theorem 2.4 are “spherical”, meaning that one of the compositions $\mathbf{a}, \mathbf{b}, \mathbf{c}$ is equal to (1^n) , or equivalently one of the factors of the triple flag variety

is a complete flag variety. Our spherical cases are identical to C. Simpson's list [9] of certain local systems on \mathbf{P}^1 with three punctures. Such local systems are equivalent to triples of matrices $X_1, X_2, X_3 \in GL_n$ with $X_1 X_2 X_3 = I$, up to simultaneous conjugation. Simpson classifies the triples of semi-simple conjugacy classes C_1, C_2, C_3 of GL_n , one of which is regular, such that the product $C_1 \times C_2 \times C_3$ contains a unique solution to the equation $X_1 X_2 X_3 = I$ (up to simultaneous conjugation). These are called *rigid* local systems. The last indecomposable type on our list corresponds to the local system associated to the Pochhammer hypergeometric function.

To describe the bijection in Theorem 2.3, we introduce the following additive category $\mathcal{F}_{p,q,r}$. The *objects* of $\mathcal{F}_{p,q,r}$ are families $(V; A, B, C)$, where V is a finite-dimensional vector space, and (A, B, C) is a triple of flags in V belonging to $\text{Fl}_{\mathbf{a}}(V) \times \text{Fl}_{\mathbf{b}}(V) \times \text{Fl}_{\mathbf{c}}(V)$ for some $(\mathbf{a}, \mathbf{b}, \mathbf{c}) \in A_{p,q,r}$. The triple $\mathbf{d} = (\mathbf{a}, \mathbf{b}, \mathbf{c})$ is called the *dimension vector* of $(V; A, B, C)$. A *morphism* from $(V; A, B, C)$ to $(V'; A', B', C')$ in $\mathcal{F}_{p,q,r}$ is a linear map $f: V \rightarrow V'$ such that $f(A_i) \subset A'_i$, $f(B_i) \subset B'_i$, and $f(C_i) \subset C'_i$ for all i . Direct sum of objects is taken componentwise on each member of each flag in the objects.

Comparing definitions, we see that isomorphism classes of objects in $\mathcal{F}_{p,q,r}$ with a given dimension vector $(\mathbf{a}, \mathbf{b}, \mathbf{c}) \in A_{p,q,r}$ are naturally identified with $GL(V)$ -orbits in $\text{Fl}_{\mathbf{a}}(V) \times \text{Fl}_{\mathbf{b}}(V) \times \text{Fl}_{\mathbf{c}}(V)$. The advantage of dealing with $\mathcal{F}_{p,q,r}$ is that this category admits direct sums, and so every object $(V; A, B, C)$ of $\mathcal{F}_{p,q,r}$ can be decomposed into a direct sum of indecomposable objects. By the Krull–Schmidt theorem (see Section 3.1), such a decomposition is unique up to an automorphism of $(V; A, B, C)$. So the isomorphism class of an object is determined by the multiplicities of the non-isomorphic indecomposable objects in its decomposition. Theorem 2.3 now becomes a consequence of the following.

THEOREM 2.6. *For every $\mathbf{d} \in \Pi_{p,q,r}$, there exists a unique isomorphism class $I_{\mathbf{d}}$ of indecomposable objects in $\mathcal{F}_{p,q,r}$ with the dimension vector \mathbf{d} . For every triple $(\mathbf{a}, \mathbf{b}, \mathbf{c}) \in A_{p,q,r}$ of finite type, any object in $\mathcal{F}_{p,q,r}$ with the dimension vector $(\mathbf{a}, \mathbf{b}, \mathbf{c})$ decomposes (uniquely) into a direct sum of objects $I_{\mathbf{d}}$.*

COROLLARY 2.7. *The bijection in Theorem 2.3 sends a family $M = (m_{\mathbf{d}}) (\mathbf{d} \in \Pi_{p,q,r})$ to the $GL(V)$ -orbit Ω_M in $\text{Fl}_{\mathbf{a}}(V) \times \text{Fl}_{\mathbf{b}}(V) \times \text{Fl}_{\mathbf{c}}(V)$ corresponding to the isomorphism class $\bigoplus_{\mathbf{d} \in \Pi_{p,q,r}} m_{\mathbf{d}} I_{\mathbf{d}}$ of objects in $\mathcal{F}_{p,q,r}$.*

2.3. Representatives of orbits. By Corollary 2.7, in order to give an explicit representative of each $GL(V)$ -orbit in a multiple flag variety of finite type, it is enough to exhibit a triple of flags that represents every indecomposable object $I_{\mathbf{d}}$ in Theorem 2.6. All possible dimension vectors $\mathbf{d} = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in \Pi_{p,q,r}$ are described in Theorem 2.4. Note that vanishing of

some part a_i in a composition \mathbf{a} means that in any flag $A \in \text{Fl}_{\mathbf{a}}(V)$ the subspace A_i coincides with A_{i-1} . Thus in constructing a representative for $I_{\mathbf{a}}$, we can assume without loss of generality that none of the compositions \mathbf{a} , \mathbf{b} and \mathbf{c} have zero parts. So it is enough to treat all the dimension vectors \mathbf{d} obtained from the triples $(\mathbf{a}^+, \mathbf{b}^+, \mathbf{c}^+)$ in Theorem 2.4 by some permutations of the parts. In particular, we assume that $p = \ell(\mathbf{a}) \leq 2$; thus a flag $A \in \text{Fl}_{\mathbf{a}}(V)$ is determined by one vector subspace A_1 in V . Under these assumptions, we will show that a triple of flags $(V; A, B, C)$ representing $I_{\mathbf{a}}$ can be presented in a standard form according to the following definition.

DEFINITION 2.8. An object $(V; A, B, C)$ in $\mathcal{F}_{p,q,r}$ is presented in *standard form* if V is given a basis e_1, \dots, e_n with the following properties:

- (1) Each subspace B_i of the flag B has a basis consisting of the first $b_1 + \dots + b_i$ standard basis vectors e_1, e_2, \dots , while each C_i has a basis consisting of the last $c_1 + \dots + c_i$ basis vectors e_n, e_{n-1}, \dots
- (2) $p \leq 2$, and the vector subspace $A_1 \subset V$ of dimension $a = a_1$ has basis vectors $\sum_{l \in S_1} e_l, \dots, \sum_{l \in S_a} e_l$ for some subsets $S_1, \dots, S_a \subset \{1, \dots, n\}$.
- (3) The subsets S_k satisfy: $|\bigcup_{k \neq k'} (S_k \cap S_{k'})| \leq 2$; that is, there are at most two elements which belong to more than one of the S_k .

THEOREM 2.9. Let $\mathbf{d} \in \Pi_{p,q,r}$ be a triple of compositions obtained from some triple $(\mathbf{a}^+, \mathbf{b}^+, \mathbf{c}^+)$ in Theorem 2.4 by permutations of the parts. Then the corresponding indecomposable object $I_{\mathbf{d}}$ in $\mathcal{F}_{p,q,r}$ can be presented in standard form with the collection of subsets S_1, \dots, S_a chosen as follows:

- $\mathbf{d} = ((1), (1), (1))$: $S_1 = \{1\}$.
- $\mathbf{d} = ((4, 2), (2^3), (1^6))$: $S_1 = \{1, 5\}$, $S_2 = \{2, 3\}$, $S_3 = \{2, 5, 6\}$, $S_4 = \{4, 5\}$.
- $\mathbf{d} = ((3^2), (2^3), (2, 1, 1, 1, 1))$: $S_1 = \{1, 2, 3\}$, $S_2 = \{1, 6\}$, $S_3 = \{2, 4, 5\}$.
- $\mathbf{d} = ((3^2), (2^3), (1, 1, 2, 1, 1))$: $S_1 = \{1, 5, 6\}$, $S_2 = \{2, 3, 6\}$, $S_3 = \{4, 5\}$.
- $\mathbf{d} = ((3^2), (2^3), (1, 1, 1, 1, 2))$: $S_1 = \{1, 4, 6\}$, $S_2 = \{2, 4, 5\}$, $S_3 = \{2, 3\}$.
- $\mathbf{d} = ((3^2), (2^3), (1, 2, 1, 1, 1))$: $S_1 = \{1, 2, 4, 6\}$, $S_2 = \{1, 3\}$, $S_3 = \{1, 5\}$.
- $\mathbf{d} = ((3^2), (2^3), (1, 1, 1, 2, 1))$: $S_1 = \{1, 2, 4, 6\}$, $S_2 = \{1, 3\}$, $S_3 = \{1, 5\}$.
- $\mathbf{d} = ((2, 4), (2^3), (1^6))$: $S_1 = \{1, 2, 3, 6\}$, $S_2 = \{1, 4, 5\}$.
- $\mathbf{d} = ((m, m+1), (1, m, m), (1^{2m+1}))$:
 $S_k = \{1, k+1, 2m+2-k\}$ ($1 \leq k \leq m$).
- $\mathbf{d} = ((m+1, m), (1, m, m), (1^{2m+1}))$: $S_1 = \{1, 2\}$,
 $S_k = \{1, k+1, 2m+3-k\}$ ($2 \leq k \leq m$), $S_{m+1} = \{1, m+2\}$.
- $\mathbf{d} = ((m, m+1), (m, 1, m), (1^{2m+1}))$:
 $S_k = \{k, m+1, 2m+2-k\}$ ($1 \leq k \leq m$).
- $\mathbf{d} = ((m+1, m), (m, 1, m), (1^{2m+1}))$: $S_1 = \{1, m+1\}$,
 $S_k = \{k, m+1, 2m+3-k\}$ ($2 \leq k \leq m$), $S_{m+1} = \{m+1, m+2\}$.
- $\mathbf{d} = ((m, m+1), (m, m, 1), (1^{2m+1}))$:
 $S_k = \{k, 2m+1-k, 2m+1\}$ ($1 \leq k \leq m$).

- $\mathbf{d} = ((m + 1, m), (m, m, 1), (1^{2m+1})): S_1 = \{1, 2m + 1\},$
- $S_k = \{k, 2m + 2 - k, 2m + 1\} (2 \leq k \leq m), S_{m+1} = \{m + 1, 2m + 1\}.$
- $\mathbf{d} = ((m, m), (1, m - 1, m), (1^{2m})):$
- $S_k = \{1, k + 1, 2m + 1 - k\} (1 \leq k \leq m - 1), S_m = \{1, m + 1\}.$
- $\mathbf{d} = ((m, m), (1, m, m - 1), (1^{2m})):$
- $S_1 = \{1, 2\}, S_k = \{1, k + 1, 2m + 2 - k\} (2 \leq k \leq m).$
- $\mathbf{d} = ((m, m), (m - 1, 1, m)(1^{2m})):$
- $S_k = \{k, m, 2m + 1 - k\} (1 \leq k \leq m - 1), S_m = \{m, m + 1\}.$
- $\mathbf{d} = ((m, m), (m, 1, m - 1), (1^{2m})):$
- $S_1 = \{1, m + 1\}, S_k = \{k, m + 1, 2m + 2 - k\} (2 \leq k \leq m).$
- $\mathbf{d} = ((m, m), (m - 1, m, 1), (1^{2m})):$
- $S_k = \{k, 2m - k, 2m\} (1 \leq k \leq m - 1), S_m = \{m, 2m\}.$
- $\mathbf{d} = ((m, m), (m, m - 1, 1), (1^{2m})):$
- $S_1 = \{1, 2m\}, S_k = \{k, 2m + 1 - k, 2m\} (2 \leq k \leq m).$
- $\mathbf{d} = ((n - 1, 1), (1^n), (1^n)): S_k = \{1, k + 1\} 1 \leq k \leq n - 1.$
- $\mathbf{d} = ((1, n - 1), (1^n), (1^n)): S_1 = \{1, 2, \dots, n\}.$

For any composition \mathbf{a} , let \mathbf{a}_{red} denote the composition obtained from \mathbf{a} by removing all zero parts and keeping the non-zero parts in the same order. For a dimension vector $\mathbf{d} = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in A_{p, q, r}$, we set $\mathbf{d}_{\text{red}} = (\mathbf{a}_{\text{red}}, \mathbf{b}_{\text{red}}, \mathbf{c}_{\text{red}})$ and call \mathbf{d}_{red} the *reduced dimension vector* of \mathbf{d} . Thus $\mathbf{d} \in \Pi_{p, q, r}$ if and only if \mathbf{d}_{red} is one of the triples in Theorem 2.9.

Finite-type indecomposables in Theorem 2.9 are closely related to *exceptional pairs* in the sense of [8]; see Remark 4.3.

EXAMPLE 2.10. Type $A_{q, r}$: two flags. Let $\mathbf{b} = (b_1, \dots, b_q)$ and $\mathbf{c} = (c_1, \dots, c_r)$ be two compositions of n . We can identify a pair of partial flags $(B, C) \in \text{Fl}_{\mathbf{b}}(V) \times \text{Fl}_{\mathbf{c}}(V)$ with the object $(V; A, B, C)$ in the category $\mathcal{F}_{1, q, r}$, where A is the trivial flag $(0 = A_0 \subset A_1 = V)$. An indecomposable summand of $(V; A, B, C)$ can only have the reduced dimension vector $((1), (1), (1))$. The indecomposable objects with this reduced dimension vector are of the form $I_{ij} = (V'; A', B', C')$ where $1 \leq i \leq q$ and $1 \leq j \leq r$: here $\dim V' = 1$, A' is the trivial flag in V' , $B' = (0 = B'_0 = \dots = B'_{i-1} \subset B'_i = \dots = B'_q = V')$, and $C' = (0 = C'_0 = \dots = C'_{j-1} \subset C'_j = \dots = C'_r = V')$.

It follows that $GL(V)$ -orbits in $\text{Fl}_{\mathbf{b}}(V) \times \text{Fl}_{\mathbf{c}}(V)$ are parametrized by $q \times r$ nonnegative integer matrices $M = (m_{ij})$ with row sums b_1, \dots, b_q and column sums c_1, \dots, c_r ; the orbit Ω_M corresponds to a direct sum $\bigoplus_{i, j} m_{ij} I_{ij}$. (In particular, if B and C are complete flags, we obtain the usual parametrization of orbits by permutation matrices). A representative of Ω_M can be given as follows: V has a basis $\{e_{ijk} : 1 \leq i \leq q, 1 \leq j \leq r, 1 \leq k \leq m_{ij}\}$, each B_i is spanned by the $e_{i'j'k}$ with $i' \leq i$, and each C_j is spanned by the $e_{i'j'k}$ with $j' \leq j$.

EXAMPLE 2.11. Type $S_{q,r}$: two flags and a line. As in Example 2.10, let \mathbf{b} and \mathbf{c} be any two compositions of n , but now let us take $\mathbf{a} = (1, n-1)$. Let $(V; A, B, C)$ be a triple of flags of type $(S_{q,r})$ with the dimension vector $(\mathbf{a}, \mathbf{b}, \mathbf{c}) \in A_{2,q,r}$. By inspection of the cases in Theorem 2.9, we see that an indecomposable summand of $(V; A, B, C)$ can only have the reduced dimension vector $((1), (1), (1))$ or $((1, t-1), (1'), (1'))$ for some $t = 2, \dots, n$. The corresponding indecomposable objects are of the following form. First each I_{ij} in the previous example can be also considered as an indecomposable object in the present situation: we take V' , B' , and C' as above, and define the flag A' as $(0 = A'_0 = A'_1 \subset A'_2 = V')$; by abuse of notation, we denote this indecomposable object in $\mathcal{F}_{2,q,r}$ by the same symbol I_{ij} .

Besides these indecomposables, the object $(V; A, B, C)$ must have precisely one indecomposable summand $(V'; A', B', C')$ with $\dim A'_1 = 1$ (since $\dim A_1 = 1$). Such indecomposables are indexed by non-empty sets $\Delta = \{(i_1, j_1), (i_2, j_2), \dots, (i_t, j_t)\}$ with $1 \leq i_1 < \dots < i_t \leq q$ and $r \geq j_1 > \dots > j_t \geq 1$ (such a Δ can be pictured as the outer corners of a Young diagram contained in a $q \times r$ rectangle). The indecomposable object I_Δ is represented by the following triple of flags $(V'; A', B', C')$: the space V' has basis e_1, \dots, e_t , the subspace A'_1 is spanned by $e_1 + \dots + e_t$, each B'_i is spanned by the e_l with $i_l \leq i$, and each C'_j is spanned by the e_l with $j_l \leq j$.

We see that $GL(V)$ -orbits in $\text{Fl}_{\mathbf{a}}(V) \times \text{Fl}_{\mathbf{b}}(V) \times \text{Fl}_{\mathbf{c}}(V)$ correspond to objects $I_\Delta \oplus \bigoplus m'_{ij} I_{ij}$ in $\mathcal{F}_{2,q,r}$ with the dimension vector $(\mathbf{a}, \mathbf{b}, \mathbf{c})$. It is convenient to use the numbers $m_{ij} = m'_{ij} + 1$ for $(i, j) \in \Delta$ and $m_{ij} = m'_{ij}$ otherwise. In this notation, the orbits are parametrized by pairs $(\Delta, M = (m_{ij}))$ where Δ is any set as above, and M is a $q \times r$ nonnegative integer matrix with row sums b_1, \dots, b_q and column sums c_1, \dots, c_r , and with $m_{ij} > 0$ for all $(i, j) \in \Delta$. A representative of the orbit $\Omega_{\Delta, M}$ is $(V; A, B, C)$ where $(V; B, C)$ is the representative of Ω_M constructed in the previous example, and A_1 is spanned by the vector $\sum_{(i,j) \in \Delta} e_{ij} 1$.

3. PROOFS OF THEOREMS 2.1–2.6

For a k -tuple of positive integers (p_1, \dots, p_k) , we define the graph T_{p_1, \dots, p_k} , the semigroup A_{p_1, \dots, p_k} , the Tits quadratic form $Q(\mathbf{d})$ on A_{p_1, \dots, p_k} , and the additive category $\mathcal{F}_{p_1, \dots, p_k}$ analogously to their counterparts for $k=3$. Also let Π_{p_1, \dots, p_k} be the set of k -tuples of compositions of finite type $\mathbf{d} \in A_{p_1, \dots, p_k}$ with $Q(\mathbf{d}) = 1$. When there is no risk of ambiguity, we drop the subscripts (p_1, \dots, p_k) and write A, \mathcal{F}, Π , etc.

3.1. *Proof of Theorem 2.6.* The only “non-elementary” part of our argument is the following proposition.

PROPOSITION 3.1. *Suppose $\mathbf{d} \in A$ is the dimension vector of an indecomposable object of \mathcal{F} with $Q(\mathbf{d}) \geq 1$. Then $Q(\mathbf{d}) = 1$, and there is a unique isomorphism class $I_{\mathbf{d}}$ of indecomposable objects with the dimension vector \mathbf{d} .*

Proof. Consider $T = T_{p_1, \dots, p_k}$ as a directed graph with all edges pointing toward the central vertex where all the chains are joined. Let \mathcal{C} be the category of quiver representations of T : recall that such a representation is specified by attaching a finite-dimensional vector space to every vertex of T , and a linear map between the corresponding spaces to every arrow (directed edge) of T . There is an obvious functor from \mathcal{F} to \mathcal{C} : given a tuple of flags, the corresponding quiver representation associates the flag subspaces to vertices of T , and inclusion maps to arrows. The image of this functor lies in the subcategory \mathcal{I} of \mathcal{C} consisting of quiver representations with all arrows represented by injective linear maps. In fact, our functor allows us to identify the isomorphism classes of indecomposable objects in \mathcal{F} with those in \mathcal{I} . Note that \mathcal{I} is a full additive subcategory of \mathcal{C} , and that indecomposables of \mathcal{I} are also indecomposables of \mathcal{C} , since an injective linear map can never have a non-injective map as a direct summand.

In view of this translation, our proposition follows from general results due to V. Kac ([5, Theorem 1]) which provide a description of dimension vectors for indecomposable quiver representations of an arbitrary finite directed graph. (These dimension vectors turn out to be in a natural bijection with positive roots of the simply-laced Kac–Moody Lie algebra corresponding to the graph.) Kac shows that the dimension vectors of indecomposables all have $Q(\mathbf{d}) \leq 1$; and if an indecomposable has $Q(\mathbf{d}) = 1$ (the case of a real root), then there is a unique indecomposable of dimension \mathbf{d} up to isomorphism. This directly implies our Proposition. ■

Note that \mathcal{F} is not an abelian category (since it does not always admit quotients). However, the Krull–Schmidt Theorem (as in [1]) still applies. That is, each object of \mathcal{F} has a unique splitting into indecomposables.

In general, the condition that $\mathbf{d} \in A$ has $Q(\mathbf{d}) = 1$ does not imply the existence of an indecomposable object $I_{\mathbf{d}}$ in \mathcal{F} with the dimension vector \mathbf{d} . However, we now show that if \mathbf{d} is of finite type with $Q(\mathbf{d}) = 1$ (i.e., if $\mathbf{d} \in \Pi$) then $I_{\mathbf{d}}$ exists and has an important additional property. For any two (isomorphism classes of) objects F and F' in \mathcal{F} , let us denote

$$\langle F', F \rangle = \dim \operatorname{Hom}_{\mathcal{F}}(F', F). \tag{3.1}$$

We say that $F \in \mathcal{F}$ is a *Schur indecomposable* if $\langle F, F \rangle = 1$ (which clearly implies that F is indeed an indecomposable object in \mathcal{F}).

PROPOSITION 3.2. *If $\mathbf{d} \in \Pi$ then there exists a Schur indecomposable $I_{\mathbf{d}}$ with the dimension vector \mathbf{d} .*

Proof. Since \mathbf{d} is of finite type, the corresponding multiple flag variety $\text{Fl}_{\mathbf{d}}(V) = \text{Fl}_{\mathbf{a}_1}(V) \times \cdots \times \text{Fl}_{\mathbf{a}_k}(V)$ has a (dense) Zariski open orbit Ω . Let $I_{\mathbf{d}}$ be the corresponding isomorphism class in \mathcal{F} , and let F be any representative of $I_{\mathbf{d}}$; by abuse of notation, we can think of F as a point in Ω . Then we have

$$\langle I_{\mathbf{d}}, I_{\mathbf{d}} \rangle = \dim \text{Stab}_{GL(V)}(F) = \dim GL(V) - \dim \text{Fl}_{\mathbf{d}}(V) = Q(\mathbf{d}) = 1.$$

Therefore, $I_{\mathbf{d}}$ is a Schur indecomposable, as desired. \blacksquare

By Propositions 3.1 and 3.2, for every $\mathbf{d} \in \Pi$, there exists a unique isomorphism class $I_{\mathbf{d}}$ of indecomposable objects in \mathcal{F} with the dimension vector \mathbf{d} . Now the proof of Theorem 2.6 (and hence that of Theorem 2.3 and Corollary 2.7) can be concluded as follows.

We say that a non-zero $\mathbf{d}' \in \Lambda$ is a *summand* of $\mathbf{d} \in \Lambda$ if $\mathbf{d} - \mathbf{d}' \in \Lambda$. It follows from the Krull–Schmidt theorem that if \mathbf{d} is of finite type then every summand of \mathbf{d} is also of finite type. Thus every object in \mathcal{F} whose dimension vector is of finite type decomposes (uniquely) into a direct sum of objects $I_{\mathbf{d}}$ for $\mathbf{d} \in \Pi$, and we are done. \blacksquare

3.2. Proof of Theorems 2.1, 2.2, and 2.4. The following criterion reduces the classification of tuples of compositions of finite type to an “elementary” problem about the Tits form.

PROPOSITION 3.3. *A tuple of compositions $\mathbf{d} \in \Lambda$ is of finite type if and only if $Q(\mathbf{d}') \geq 1$ for any summand \mathbf{d}' of \mathbf{d} .*

Proof. Let $\text{Fl}_{\mathbf{d}}(V)$ be the multiple flag variety corresponding to $\mathbf{d} \in \Lambda$. First suppose \mathbf{d} is of finite type. Since the one-dimensional subgroup of scalar matrices in $GL(V)$ acts trivially on $\text{Fl}_{\mathbf{d}}(V)$, we must have $\dim GL(V) - 1 \geq \dim \text{Fl}_{\mathbf{d}}(V)$, i.e., $Q(\mathbf{d}) \geq 1$. We have already noticed that any summand \mathbf{d}' of \mathbf{d} must also be of finite type, hence we must have $Q(\mathbf{d}') \geq 1$.

Conversely, suppose $Q(\mathbf{d}') \geq 1$ for any summand \mathbf{d}' of \mathbf{d} . In view of Proposition 3.1, this implies that every indecomposable summand of an object in \mathcal{F} with the dimension vector \mathbf{d} is uniquely determined by its dimension vector. Therefore, the isomorphism classes of objects with the dimension vector \mathbf{d} are in a bijection with partitions of \mathbf{d} into the sum of dimension vectors of indecomposables. Since there are finitely many such partitions, \mathbf{d} must be of finite type, and we are done. \blacksquare

Remark 3.4. The criterion in Proposition 3.3 is almost identical to that of V. Kac [5, Proposition 2.4], for finite-type quiver varieties: the quiver variety with a given dimension vector \mathbf{d} has finitely many orbits exactly if $Q(\mathbf{d}') \geq 1$ for all *quiver* summands \mathbf{d}' of \mathbf{d} . (A quiver summand need not

have positive jumps in dimension along each flag; only the dimension of each space must be positive.

To illustrate the difference between the two criteria, consider the triple of compositions $\mathbf{d} = ((3, 1), (1^4), (1^4))$ of finite type $S_{4,4}$. The corresponding space of quiver representations has infinitely many orbits because it has the quiver summand $\mathbf{d}' = ((2, 2), (1^4), (1^4))$ with $Q(\mathbf{d}') = 0$; in terms of dimensions rather than dimension jumps, ${}^{1234321}_3 = {}^{1234321}_2 + {}^{0000000}_1$. In fact, the infinitely many generic objects of dimension \mathbf{d}' are all subobjects of the unique generic indecomposable $I_{\mathbf{d}}$, but there are no quotient objects in the flag category \mathcal{F} , only in the quiver category \mathcal{C} . (We thank C. Ringel for this example.)

The classification of finite types in the rest of this section closely follows the Cartan–Killing classification, which in our terminology amounts to finding all graphs T_{p_1, \dots, p_k} with positive-definite Tits form.

Proof of Theorem 2.1. Clearly, if \mathbf{d} is a k -tuple of compositions of finite type then any subtuple of \mathbf{d} is also of finite type. Thus it suffices to show that a quadruple \mathbf{d} of non-trivial compositions cannot be of finite type. But any such quadruple has a summand \mathbf{d}' with the reduced dimension vector $((1^2), (1^2), (1^2), (1^2))$. A calculation shows that $Q(\mathbf{d}') = 0$, so by Proposition 3.3, \mathbf{d} cannot be of finite type. ■

Next, beginning the proof of Theorem 2.2, we eliminate those dimension vectors with a summand corresponding to the minimal imaginary root of an affine root system. Let $N_{p,q,r}$ be the set of all $\mathbf{d}' = (\mathbf{a}', \mathbf{b}', \mathbf{c}') \in A_{p,q,r}$ such that $\{\mathbf{a}'_{\text{red}}, \mathbf{b}'_{\text{red}}, \mathbf{c}'_{\text{red}}\}$ is one of the following three triples:

$$\{(1^3), (1^3), (1^3)\}, \quad \{(2^2), (1^4), (1^4)\}, \quad \{(3^2), (2^3), (1^6)\}. \quad (3.2)$$


The dimension vectors are associated via the Kac correspondence with minimal imaginary roots for the affine Lie algebras \hat{E}_6 , \hat{E}_7 and \hat{E}_8 , respectively (cf. the proof of Proposition 3.1). (Note that the quadruple $((1^2), (1^2), (1^2), (1^2))$ that appeared in the proof of Theorem 2.1 corresponds to the minimal imaginary root for \hat{D}_4 .) Using formula (2.1), we find $Q(\mathbf{d}') = 0$ for any $\mathbf{d}' \in N_{p,q,r}$.

Without loss of generality, we can assume that a triple $\mathbf{d} = (\mathbf{a}, \mathbf{b}, \mathbf{c})$ is reduced, i.e., all the compositions \mathbf{a} , \mathbf{b} and \mathbf{c} have all parts non-zero. Thus, $\mathbf{d} \in A_{p,q,r}$, where p , q , and r have the same meaning as in Theorem 2.2.

LEMMA 3.5. *Let $\mathbf{d} \in A_{p,q,r}$ be a reduced triple of compositions. Then exactly one of the following holds:*

- (i) \mathbf{d} belongs to one of the types A–S in Theorem 2.2.
- (ii) The triple \mathbf{d} has some $\mathbf{d}' \in N_{p,q,r}$ as a summand.

Proof. Following the usual classification of Dynkin diagrams, we present the proof in the schematic form of a tree of implications:



The root of the tree is our

HYPOTHESIS. $\mathbf{d} \in A_{p,q,r}$ is a reduced triple of compositions, and $1 \leq p \leq q \leq r$.

The arrows coming from a statement point to all possible cases resulting from the statement. We employ the abuse of notation $\mathbf{d} = A$, $\mathbf{d} = D$, etc. to indicate that \mathbf{d} belongs to the corresponding type in Theorem 2.2. Similarly we write $\mathbf{d} \geq \widehat{E}_6$, etc. to indicate that \mathbf{d} has a summand corresponding to the given affine type. The lemma follows because every case ends in (i) or (ii), and these conditions are clearly disjoint.

Combining Lemma 3.5 with Proposition 3.3, we prove one direction of Theorem 2.2: if \mathbf{d} is of finite type then it necessarily belongs to one of the types A–S. It remains to show that each of the conditions A–S is *sufficient* for $(\mathbf{a}, \mathbf{b}, \mathbf{c})$ to be of finite type. The following lemma is an immediate consequence of Lemma 3.5.

LEMMA 3.6. *If $\mathbf{d} \in A_{p,q,r}$ is of one of the types A–S then the same is true for any summand of \mathbf{d} .*

The remaining part of Theorem 2.2 now follows by combining Proposition 3.3 with Lemma 3.6 and the following.

LEMMA 3.7. *Suppose $\mathbf{d} \in A_{p,q,r}$ is of one of the types A–S. Then $Q(\mathbf{d}) \geq 1$.*

Thus it only remains to prove Lemma 3.7, which we deduce from formula (2.1) and the following elementary estimates.

LEMMA 3.8. *Let \mathbf{b} be a reduced composition with $|\mathbf{b}| = n$ and $\ell(\mathbf{b}) = q$. Then*

- (1) $\|\mathbf{b}\|^2 \geq n$, with equality precisely when $\mathbf{b} = (1^n)$;
- (2) if $q = 3$ then $\|\mathbf{b}\|^2 \geq 3(n - 2)$, with equality precisely when $\max(b_1, b_2, b_3) \leq 2$;
- (3) if $q = 2$, and $n = 2m$ is even then $\|\mathbf{b}\|^2 \geq 2m^2$, with equality precisely when $\mathbf{b} = (m, m)$;
- (4) if $q = 2$, and $n = 2m + 1$ is odd then $\|\mathbf{b}\|^2 \geq 2m^2 + 2m + 1$, with equality precisely when $\mathbf{b}^+ = (m + 1, m)$.

Proof. Easy. For example, part (2) is a consequence of the identity:

$$\|\mathbf{b}\|^2 - 3(n - 2) = \sum_{i=1}^3 (b_i^2 - 3b_i + 2) = \sum_{i=1}^3 (b_i - 1)(b_i - 2).$$

The other parts are even simpler. ■

Proof of Lemma 3.7. Suppose $\mathbf{d} = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in A_{p,q,r}$ is of one of the types A–S.

Case 1. Suppose \mathbf{d} is of one of the types A, D, E_6 , E_7 , or E_8 . Then the form Q is positive definite (this is the Cartan–Killing classification), so $Q(\mathbf{d}) \geq 1$. Furthermore, the equality $Q(\mathbf{d}) = 1$ occurs precisely when \mathbf{d} corresponds to a positive root of the associated simple Lie algebra (cf. proof of Proposition 3.1).

Case 2. Suppose \mathbf{d} is of type $E^{(a)}$. Now the desired inequality $Q(\mathbf{d}) \geq 1$ follows from the equality

$$\|\mathbf{a}\|^2 - n^2 = 2^2 + (n - 2)^2 - n^2 = 8 - 4n$$

and the inequalities $\|\mathbf{b}\|^2 \geq 3(n - 2)$ and $\|\mathbf{c}\|^2 \geq n$ (Lemma 3.8, parts (1), (2)). The equality $Q(\mathbf{d}) = 1$ occurs precisely when $(\mathbf{a}^+, \mathbf{b}^+, \mathbf{c}^+) = ((2, 2), (2, 1, 1), (1^4)), ((3, 2), (2, 2, 1), 1^5),$ or $((4, 2), (2^3), (1^6))$.

Case 3. Suppose \mathbf{d} is of type $E^{(b)}$. Let \mathbf{b}' be the composition obtained from \mathbf{b} by removing a part equal to 1, so that we have $|\mathbf{b}'| = n - 1$, $\ell(\mathbf{b}') = 2$, and $\|\mathbf{b}\|^2 = \|\mathbf{b}'\|^2 + 1$. If $n = 2m$ is even then $Q(\mathbf{d}) \geq 1$ follows from the inequalities

$$\|\mathbf{a}\|^2 \geq 2m^2, \quad \|\mathbf{b}'\|^2 \geq 2m^2 - 2m + 1, \quad \|\mathbf{c}\|^2 \geq 2m$$

(Lemma 3.8, parts (1), (3) and (4)). The equality $Q(\mathbf{d}) = 1$ occurs precisely when $(\mathbf{a}^+, \mathbf{b}^+, \mathbf{c}^+) = ((m, m), (m, m - 1, 1), (1^{2m}))$.

If $n = 2m + 1$ is odd then $Q(\mathbf{d}) \geq 1$ follows from the inequalities

$$\|\mathbf{a}\|^2 \geq 2m^2 + 2m + 1, \quad \|\mathbf{b}'\|^2 \geq 2m^2, \quad \|\mathbf{c}\|^2 \geq 2m$$

(Lemma 3.8, parts (1), (3) and (4)). The equality $Q(\mathbf{d}) = 1$ occurs precisely when $(\mathbf{a}^+, \mathbf{b}^+, \mathbf{c}^+) = ((m+1, m), (m, m, 1), (1^{2m+1}))$.

Case 4. Suppose \mathbf{d} is of type S . Then $Q(\mathbf{d}) \geq 1$ follows from

$$\|\mathbf{a}\|^2 - n^2 = 1 + (n-1)^2 - n^2 = 2 - 2n$$

and $\|\mathbf{b}\|^2 \geq n$, $\|\mathbf{c}\|^2 \geq n$ (Lemma 3.8, part (1)). The equality $Q(\mathbf{d}) = 1$ occurs precisely when $(\mathbf{a}^+, \mathbf{b}^+, \mathbf{c}^+) = ((n-1, 1), (1^n), (1^n))$.

This completes the proofs of Lemma 3.7 and Theorem 2.2. \blacksquare

As a by-product of the above argument (the examination of the equality $Q(\mathbf{d}) = 1$), we immediately obtain Theorem 2.4.

4. ORBIT REPRESENTATIVES AND GENERALIZED BRUHAT ORDER

4.1. Morphisms of standard triples of flags. We have seen (Theorem 2.6) that the finite-type indecomposable objects of the triple flag category \mathcal{F}_{pqr} occur only in dimensions $\mathbf{d} \in \Pi_{pqr}$, and that each such dimension contains a unique indecomposable isomorphism class $I_{\mathbf{d}}$. Furthermore, by Proposition 3.2, this $I_{\mathbf{d}}$ is characterized among all objects of dimension \mathbf{d} by the Schur condition $\langle I_{\mathbf{d}}, I_{\mathbf{d}} \rangle = 1$. Thus, to prove Theorem 2.9, we need only show that each of the standard forms listed there is Schur. We will deduce the Schur property from a general formula for $\langle F', F \rangle$, where F' is an arbitrary object presented in the standard form.

Let $F' = (V'; A', B', C')$ be an object with $\dim V' = n$, presented in the standard form of Definition 2.8. Then the flags (B', C') can be encoded by a family

$$\Delta = ((i_1, j_1), \dots, (i_n, j_n))$$

of n pairs of indices satisfying $1 \leq i_1 \leq \dots \leq i_n \leq q$ and $r \geq j_1 \geq \dots \geq 1$. These are defined in terms of the dimension vectors (b_1, \dots, b_q) and (c_1, \dots, c_r) of B' and C' by:

$$i_l = \min\{i \mid b_1 + \dots + b_i \geq l\}$$

$$j_l = \min\{j \mid c_1 + \dots + c_j \geq n + 1 - l\}.$$

This means that each subspace B'_i (resp. C'_j) is spanned by the standard basis vectors e_i such that $i_l \leq i$ (resp. $j_l \leq j$). The set Δ reflects the decomposition of the pair of flags B', C' : in the terminology of Example 2.10, the triple $(V'; B', C') \in \mathcal{F}_{q,r}$ is a direct sum $\bigoplus_{l=1}^n I_{i_l j_l}$.

We see that the standard form $F' = (V'; A', B', C')$ is completely determined by the following combinatorial data: a family \mathcal{A} and a collection of subsets S_1, \dots, S_a in $\{1, \dots, n\}$. The sets S_k must satisfy condition (3) in Definition 2.8, which says that the set

$$K := \bigcup_{k \neq k'} (S_k \cap S_{k'})$$

has at most two elements.

PROPOSITION 4.1. *Let $F' = (V'; A', B', C')$ be an object of \mathcal{F}_{pqr} presented in the standard form of Definition 2.8, encoded as above by a set \mathcal{A} , and by subsets S_1, \dots, S_a . Let $F = (V; A, B, C)$ be any object of \mathcal{F}_{pqr} , and define*

$$D_l = B_{i_l} \cap C_{j_l}, \quad E_k = \sum_{l \in S_k \setminus K} D_l.$$

Let $v \mapsto \bar{v}$ denote the natural projection $V \rightarrow V/A_1$, so that $\bar{U} = (U + A_1)/A_1 \subset V/A_1$ for any subspace $U \subset V$. Then the dimension $\langle F', F \rangle$ of the space of homomorphisms in \mathcal{F}_{pqr} from F' to F is given as follows:

(1) *If $K = \{\mu, \nu\}$ for some indices $\mu \neq \nu$, then*

$$\begin{aligned} \langle F', F \rangle = & \sum_{l=1}^n \dim D_l - \dim \bar{D}_\mu - \dim \bar{D}_\nu - \sum_{k=1}^a \dim \bar{E}_k \\ & + \dim \left(\bar{D}_\mu \cap \bar{D}_\nu \cap \bigcap_{|S_k \cap K|=1} \bar{E}_k \right) \\ & + \dim \left(\bigcap_{K \subset S_k} \bar{E}_k \cap \left(\left(\bar{D}_\mu \cap \bigcap_{S_k \cap K = \{\mu\}} \bar{E}_k \right) \right. \right. \\ & \left. \left. + \left(\bar{D}_\nu \cap \bigcap_{S_k \cap K = \{\nu\}} \bar{E}_k \right) \right) \right). \end{aligned} \tag{4.1}$$

(2) *If $K = \{\mu\}$ for some index μ then*

$$\langle F', F \rangle = \sum_{l=1}^n \dim D_l - \dim \bar{D}_\mu - \sum_{k=1}^a \dim \bar{E}_k + \dim \left(\bar{D}_\mu \cap \bigcap_{\mu \in S_k} \bar{E}_k \right). \tag{4.2}$$

(3) *If $K = \emptyset$ (i.e., all S_k are pairwise disjoint) then*

$$\langle F', F \rangle = \sum_{l=1}^n \dim D_l - \sum_{k=1}^a \dim \bar{E}_k. \tag{4.3}$$

Proof. We will only prove the most complicated formula (4.1). A morphism from F' to F is a linear map from V' to V , and so is determined

by the images of the basis vectors e_1, \dots, e_n ; let us denote these images by v_1, \dots, v_n . By the definition, the vectors v_l must satisfy the following conditions:

$$v_l \in D_l \quad (1 \leq l \leq n), \quad \sum_{l \in S_k} v_l \in A_1 \quad (1 \leq k \leq a). \quad (4.4)$$

Thus $\langle F', F \rangle$ is equal to the dimension of the subspace $U \subset V^n$ formed by n -tuples (v_1, \dots, v_n) satisfying (4.4). Clearly, $U = \text{Ker}(\varphi)$, where $\varphi: \bigoplus_{l=1}^n D_l \rightarrow (V/A_1)^a$ is the linear map

$$\varphi(v_1, \dots, v_n) \mapsto \left(\sum_{l \in S_1} \bar{v}_l, \dots, \sum_{l \in S_a} \bar{v}_l \right).$$

Thus we have

$$\langle F', F \rangle = \sum_{l=1}^n \dim D_l - \text{rk}(\varphi). \quad (4.5)$$

Consider the subspace

$$W = \bar{D}_\mu \oplus \bar{D}_\nu \oplus \bigoplus_{k=1}^a \bar{E}_k \subset (V/A_1)^{a+2}.$$

Then the map $\varphi: \bigoplus_{l=1}^n D_l \rightarrow (V/A_1)^a$ can be factored as $\varphi = \varphi_2 \circ \varphi_1$:

$$\bigoplus_{l=1}^n D_l \xrightarrow{\varphi_1} W \xrightarrow{\varphi_2} (V/A_1)^a,$$

where

$$\varphi_1: (v_1, \dots, v_n) \mapsto \left(\bar{v}_\mu, \bar{v}_\nu, \sum_{l \in S_1 \setminus \{\mu, \nu\}} \bar{v}_l, \dots, \sum_{l \in S_a \setminus \{\mu, \nu\}} \bar{v}_l \right),$$

and

$$\begin{aligned} \varphi_2: (w^{(\mu)}, w^{(\nu)}, w_1, \dots, w_a) \\ \mapsto (\chi_1(\mu) w^{(\mu)} + \chi_1(\nu) w^{(\nu)} + w_1, \dots, \chi_a(\mu) w^{(\mu)} + \chi_a(\nu) w^{(\nu)} + w_a). \end{aligned}$$

(Here χ_k stands for the indicator function of the set S_k , i.e., $\chi_k(l) = 1$ if $l \in S_k$, otherwise $\chi_k(l) = 0$.) Since the sets $S_1 \setminus \{\mu, \nu\}, \dots, S_a \setminus \{\mu, \nu\}$ are pairwise disjoint, the map φ_1 is surjective. It follows that

$$\begin{aligned} \text{rk}(\varphi) &= \text{rk}(\varphi_2) = \dim W - \dim \text{Ker}(\varphi_2) \\ &= \dim \bar{D}_\mu + \dim \bar{D}_\nu + \sum_{k=1}^a \dim \bar{E}_k - \dim \text{Ker}(\varphi_2). \end{aligned} \quad (4.6)$$

It remains to compute $\dim \text{Ker}(\varphi_2)$. The definition of φ_2 implies that the projection $(w^{(\mu)}, w^{(\nu)}, w_1, \dots, w_a) \mapsto (w^{(\mu)}, w^{(\nu)})$ restricts to an isomorphism between $\text{Ker}(\varphi_2)$ and the space of pairs $(w^{(\mu)}, w^{(\nu)})$ such that

$$w^{(\mu)} \in \bar{D}_\mu \cap \bigcap_{\substack{\mu \in S_k \\ \nu \notin S_k}} \bar{E}_k, \quad w^{(\nu)} \in \bar{D}_\nu \cap \bigcap_{\substack{\mu \notin S_k \\ \nu \in S_k}} \bar{E}_k,$$

$$w^{(\mu)} + w^{(\nu)} \in \bigcap_{\{\mu, \nu\} = S_k} \bar{E}_k.$$

It follows that

$$\begin{aligned} \dim \text{Ker}(\varphi_2) = & \dim \left(\left(\bar{D}_\mu \cap \bigcap_{\substack{\mu \in S_k \\ \nu \notin S_k}} \bar{E}_k \right) \cap \left(\bar{D}_\nu \cap \bigcap_{\substack{\mu \notin S_k \\ \nu \in S_k}} \bar{E}_k \right) \right) \\ & + \dim \left(\bigcap_{\{\mu, \nu\} = S_k} \bar{E}_k \cap \left(\left(\bar{D}_\mu \cap \bigcap_{\substack{\mu \in S_k \\ \nu \notin S_k}} \bar{E}_k \right) + \left(\bar{D}_\nu \cap \bigcap_{\substack{\mu \notin S_k \\ \nu \in S_k}} \bar{E}_k \right) \right) \right) \end{aligned} \tag{4.7}$$

Combining (4.5), (4.6), and (4.7) we obtain the desired formula (4.1). ■

4.2. Proof of Theorem 2.9. Let $F \in \mathcal{F}_{p, q, r}$ be one of the standard-form triples of flags in Theorem 2.9. It suffices to show that F is a Schur indecomposable, i.e., that $\langle F, F \rangle = 1$ (cf. (3.1) and Proposition 3.2).

In the first and last case on the list, $\mathbf{d} = ((1), (1), (1))$ and $\mathbf{d} = ((1, n-1), (1^n), (1^n))$, the equality $\langle F, F \rangle = 1$ follows at once from (4.3).

In each of the first four 6-dimensional cases on the list, the desired equality $\langle F, F \rangle = 1$ is a direct consequence of (4.1). It is also easy to check by an independent calculation that every morphism from F to itself is scalar. For instance, let us do this for $\mathbf{d} = ((4, 2), (2^3), (1^6))$. Let (x_{ij}) be a 6×6 matrix that represents a morphism $\varphi: V \rightarrow V$ in the standard basis e_1, \dots, e_6 . The condition that φ preserves the flags B and C means that the only non-zero matrix entries can be $x_{11}, x_{21}, x_{22}, x_{33}, x_{43}, x_{44}, x_{55}, x_{65}$, and x_{66} . Thus we have $\varphi(e_2 + e_3) = x_{22}e_2 + x_{33}e_3 + x_{43}e_4$; the condition that this vector lies in A_1 implies that $x_{22} = x_{33}$ and $x_{43} = 0$. Similarly, the condition that $\varphi(e_4 + e_5) \in A_1$ implies that $x_{44} = x_{55}$ and $x_{65} = 0$. Finally, the two remaining conditions that $\varphi(e_1 + e_5)$ and $\varphi(e_2 + e_5 + e_6)$ lie in A_1 imply that $x_{11} = x_{55}$, $x_{21} = 0$; and $x_{22} = x_{55} = x_{66}$. Combining all these equalities, we see that φ is scalar, as desired.

For the rest of the list, the equality $\langle F, F \rangle = 1$ can be checked case by case with the help of (4.2). To simplify this procedure, we observe that all these cases satisfy the following strengthened form of condition (3) in Definition 2.8:

(3') Each set S_k has at least two elements, $\bigcup_{k=1}^a S_k = \{1, \dots, n\}$, and there exists an index μ such that $S_k \cap S_{k'} = \{\mu\}$ for all $k \neq k'$.

Assuming (3'), we will give combinatorial conditions on subsets S_k that are necessary and sufficient for the corresponding object F to be Schur indecomposable. This requires some terminology.

Let $F = (V; A, B, C)$ be a triple of flags with the dimension vector $(\mathbf{a}, \mathbf{b}, \mathbf{c})$; let $n = \dim V$. We associate to \mathbf{b} the subdivision of $[1, n] = \{1, \dots, n\}$ into consecutive blocks $[1, b_1], [b_1 + 1, b_1 + b_2], \dots$ of sizes b_1, \dots, b_q . The blocks of this subdivision will be called \mathbf{b} -blocks. We define the \mathbf{c} -blocks similarly, except going the opposite way (so that the first \mathbf{c} -block is $[n - c_1 + 1, n]$). We say that an index $l \in [1, n]$ is \mathbf{b} -separated (resp. \mathbf{c} -separated) from a subset $S \subset [1, n]$ if no element of S smaller (resp. larger) than l lies in the same \mathbf{b} -block (resp. \mathbf{c} -block) with l . If l is both \mathbf{b} -separated and \mathbf{c} -separated from S , we say that l is \mathbf{bc} -separated from S .

PROPOSITION 4.2. *Suppose $F = (V; A, B, C) \in \mathcal{F}_{p, q, r}$ has the dimension vector $(\mathbf{a}, \mathbf{b}, \mathbf{c})$ and is presented in a standard form satisfying (3'). Let $a = a_1$, and denote $S'_0 = \{\mu\}$, $S'_k = S_k \setminus \{\mu\}$.*

Then F is a Schur indecomposable if and only if the subsets S'_k for $k = 0, \dots, a$ satisfy the following conditions:

- (1) *No two elements of the same S'_k lie in the same \mathbf{b} -block or in the same \mathbf{c} -block.*
- (2) *For every two distinct indices j and k , each $l \in S'_j$ is either \mathbf{b} -separated or \mathbf{c} -separated from S'_k .*
- (3) *In the situation of (2), S'_j contains an index \mathbf{bc} -separated from S'_k .*
- (4) *For every j , any two elements of S'_j are equivalent to each other with respect to the equivalence relation generated by the following: $l \sim l'$ if, for some $k \neq j$, both l and l' are \mathbf{bc} -separated from S'_k .*

Proof. We compute $\langle F, F \rangle$ using (4.2). Note that under the condition (3'), we have $E_k = \sum_{l \in S'_k} D_l$ for $k = 1, \dots, a$. Also define $E_0 = D_\mu$. Then formula (4.2) further simplifies as follows:

$$\langle F, F \rangle = \sum_{l=1}^n \dim D_l - \sum_{k=0}^a \dim \bar{E}_k + \dim \left(\bigcap_{i=0}^a \bar{E}_k \right). \quad (4.8)$$

Tracing the definitions, we observe that each subspace $D_l = B_{i_l} \cap C_{j_l}$ is spanned by all the basis vectors e_ρ such that ρ is not \mathbf{bc} -separated from $\{l\}$. In particular, $e_l \in D_l$. It follows that $\bar{e}_\mu \in \bar{E}_k$ for all k , and it is also clear that $\bar{e}_\mu \neq 0$. Thus the last term of formula 4.8 must contribute exactly 1 to $\langle F, F \rangle$, and the first two terms must contribute 0. We thus find that $\langle F, F \rangle = 1$ if and only if:

- (i) for each $k = 0, \dots, a$, the sum $\sum_{l \in S'_k} D_l$ is direct;
- (ii) for each k , we have $A_1 \cap E_k = 0$; and
- (iii) $\bigcap_{i=0}^a \overline{E}_k = \langle \bar{e}_\mu \rangle$, a one-dimensional space.

It is now completely straightforward to show that conditions (i)–(iii) are equivalent to conditions (1)–(4) in our proposition. To be more precise, (i) translates into (1) and (2), (ii) translates into (3), and (iii) into (4). ■

Now an easy inspection shows that all the remaining cases in Theorem 2.9 satisfy conditions (1)–(4) in Proposition 4.2. In most of these cases, the inspection is simplified even more by the following observation: if $\mathbf{c} = (1^n)$ then condition (2) is automatic. This completes the proof of Theorem 2.9. ■

It is easy to show that in the four exceptional 6-dimensional cases there exist no subset S_1, \dots, S_a satisfying the conditions in Proposition 4.2 (this check starts with the observation that the case $j = 0$ in condition (3) means that an index μ must be the minimal element of its \mathbf{b} -block and the maximal element of its \mathbf{c} -block). This justifies our efforts in obtaining (4.1).

Remark 4.3. Our finite-type indecomposables $I_{\mathbf{a}}$ are Schur objects of \mathcal{F} , also known in quiver theory as exceptional objects. It is possible to obtain the list of representatives in Theorem 2.9 by a recursive procedure which is a special case of the mutations of exceptional pairs studied by Rudakov, Schofield, Crawley–Boevey, and Ringel (see [8]). In our situation, this procedure relies on the following simple general proposition.

PROPOSITION 4.4. *Suppose there is a short exact sequence*

$$0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$$

in \mathcal{F} with the following properties: both F' and F'' are Schur indecomposables, and $\langle F', F'' \rangle = \langle F, F' \rangle \langle F'', F \rangle = 0$. Then F is a Schur indecomposable.

It turns out that, for every dimension vector \mathbf{d} in Theorem 2.9, one can construct a short exact sequence as in Proposition 4.4 such that F has dimension vector \mathbf{d} , and one of the Schur indecomposables F' and F'' has reduced dimension vector $((1), (1), (1))$. The other summand is smaller than \mathbf{d} and is also on our list so we can assume that we already know its “nice” presentation; we can then use an explicit form of the short exact sequence to construct a “nice” presentation for F . For instance, if $\mathbf{d} = ((4, 3), (3, 1, 3), (1^7))$ then we can choose the dimension vectors of F' and F'' to be respectively $\mathbf{d}' = ((3, 3), (3, 1, 2), (1, 1, 0, 1, 1, 1, 1))$, and $\mathbf{d}'' = ((1, 0), (0, 0, 1), (0, 0, 1, 0, 0, 0, 0))$. Iterating this procedure, one can construct representatives for all Schur indecomposables on our list (this was in fact our original way to do it).

4.3. *Generalized Bruhat order.* Having determined the orbits in triple flag varieties of finite type, we naturally ask how they fit together. Recall that a parametrization of orbits is given by Theorem 2.3 and Corollary 2.7. We define the partial order (called *degeneration order* or *generalized Bruhat order*) on the set of families $M = (m_{\mathbf{d}})_{(\mathbf{d} \in \Pi_{p,q,r})}$ by setting $M \leq^{\text{deg}} M'$ if the orbit Ω_M lies in the Zariski closure of $\Omega_{M'}$.

Recall from Theorem 2.6 that the orbit Ω_M corresponds to the isomorphism class $\bigoplus_{\mathbf{d} \in \Pi_{p,q,r}} m_{\mathbf{d}} I_{\mathbf{d}}$ in the category $\mathcal{F}_{p,q,r}$; denote this isomorphism class by F_M . The following proposition is a special case of a result due to C. Riedtmann (cf. [2, 3, 7]).

PROPOSITION 4.5. *If $M \leq^{\text{deg}} M'$ then $\langle I_{\mathbf{d}}, F_M \rangle \geq \langle I_{\mathbf{d}}, F_{M'} \rangle$ for any $\mathbf{d} \in \Pi_{p,q,r}$.*

It would be interesting to know if the converse statement is also true, i.e., if the degeneration order $M \leq^{\text{deg}} M'$ is given by the inequalities $\langle I_{\mathbf{d}}, F_M \rangle \geq \langle I_{\mathbf{d}}, F_{M'} \rangle$ for all $\mathbf{d} \in \Pi_{p,q,r}$. This is true when the graph $T_{p,q,r}$ is one of the Dynkin graphs A_n, D_n, E_6, E_7 , or E_8 , as a consequence of general results due to K. Bongartz (cf. [2, Section 4], [3, Section 5.2]).

Note that Theorem 2.9 and formulas (4.1), (4.3), and (4.8) allow us to compute $\langle I_{\mathbf{d}}, F_M \rangle$ explicitly for all $\mathbf{d} \in \Pi_{p,q,r}$. In particular, it is easy to compute $\langle I_{\mathbf{d}}, I_{\mathbf{d}'} \rangle$ for any two Schur indecomposables of finite type. Knowing these numbers yields an explicit formula for $\langle F_M, F_M \rangle$:

$$\langle F_M, F_M \rangle = \sum_{\mathbf{d}, \mathbf{d}' \in \Pi_{p,q,r}} \langle I_{\mathbf{d}}, I_{\mathbf{d}'} \rangle m_{\mathbf{d}} m_{\mathbf{d}'}. \quad (4.9)$$

This yields a formula for the (co)dimension of any orbit Ω_M .

PROPOSITION 4.6. *The codimension of the orbit Ω_M in a multiple flag variety $\text{Fl}_{\mathbf{a}}(V) \times \text{Fl}_{\mathbf{b}}(V) \times \text{Fl}_{\mathbf{c}}(V)$ of finite type is given by*

$$\text{codim}(\Omega_M) = \langle F_M, F_M \rangle - Q(\mathbf{a}, \mathbf{b}, \mathbf{c}). \quad (4.10)$$

Proof. This follows at once from the formula

$$\langle F_M, F_M \rangle = \dim \text{Stab}_{GL(V)}(F) = \dim GL(V) - \dim \Omega_M,$$

where F is any representative of Ω_M (cf. the proof of Proposition 3.2). ■

EXAMPLE 4.7. Type $A_{q,r}$: two flags. We use the notation of Example 2.10, so that Ω_M denotes the orbit in $\text{Fl}_{\mathbf{b}}(V) \times \text{Fl}_{\mathbf{c}}(V)$ corresponding to a

$q \times r$ nonnegative integer matrix $M = (m_{ij})$ with row sums b_1, \dots, b_q and column sums c_1, \dots, c_r . Formula (4.3) specializes to

$$\langle I_{ij}, F \rangle = \dim(B_i \cap C_j)$$

for any pair of flags $F = (V; B, C)$. It follows that

$$\langle I_{ij}, F_M \rangle = \sum_{k=1}^i \sum_{l=1}^j m_{kl}. \tag{4.11}$$

By Proposition 4.5 and results of Bongartz quoted above, the degeneration order $M \stackrel{\text{deg}}{\leq} M'$ is given by the inequalities

$$\sum_{k=1}^i \sum_{l=1}^j m_{kl} \geq \sum_{k=1}^i \sum_{l=1}^j m'_{kl}$$

for all i and j . (If $\mathbf{b} = \mathbf{c} = (1^n)$, this is Ehresmann's original description of the Bruhat order on the symmetric group.) Finally, (4.10) implies the following formula for the codimension of an orbit Ω_M in $\text{Fl}_{\mathbf{b}}(V) \times \text{Fl}_{\mathbf{c}}(V)$:

$$\text{codim}(\Omega_M) = \sum_{k < i, l < j} m_{ij} m_{kl}. \tag{4.12}$$

EXAMPLE 4.8. Type $S_{q,r}$: two flags and a line. We will use the notation of Example 2.11. In particular, $\Omega_{A,M}$ denotes the orbit in $\text{Fl}_{\mathbf{a}}(V) \times \text{Fl}_{\mathbf{b}}(V) \times \text{Fl}_{\mathbf{c}}(V)$ corresponding to a $q \times r$ nonnegative integer matrix $M = (m_{ij})$ as above and to a non-empty set $A = \{(i_1, j_1), (i_2, j_2), \dots, (i_t, j_t)\}$ such that $1 \leq i_1 < \dots < i_t \leq q$, $r \geq j_1 > \dots > j_t \geq 1$, and $m_{ij} > 0$ for $(i, j) \in A$. For any triple of flags $F = (V; A, B, C)$, recall that A_1 is the only proper subspace in the flag A , and $\dim A_1 = 1$. Using (4.3), we obtain

$$\begin{aligned} \langle I_{ij}, F \rangle &= \dim(B_i \cap C_j), \\ \langle I_{A'}, F \rangle &= \dim \left(A_1 \cap \sum_{(i,j) \in A'} (B_i \cap C_j) \right) + \sum_{(i,j) \in A'} \dim(B_i \cap C_j) \\ &\quad - \dim \left(\sum_{(i,j) \in A'} (B_i \cap C_j) \right). \end{aligned}$$

These formulas imply that

$$\begin{aligned} \langle I_{ij}, F_{A,M} \rangle &= \sum_{k=1}^i \sum_{l=1}^j m_{kl}, \\ \langle I_{A'}, F_{A,M} \rangle &= \delta_{A \leq A'} + \sum_{(i,j) \in \text{In}(A')} \sum_{k=1}^i \sum_{l=1}^j m_{kl}, \end{aligned} \tag{4.13}$$

where we use the following notation: $\Delta \leq \Delta'$ means that for any $(k, l) \in \Delta$ there exists $(i, j) \in \Delta'$ such that $k \leq i$ and $l \leq j$; the δ -symbol has the usual indicator meaning; and the operation $\Delta' \mapsto \text{In}(\Delta')$ is defined by

$$\text{In}(\{(i_1, j_1), \dots, (i_t, j_t)\}) = \{(i_1, j_2), (i_2, j_3), \dots, (i_{t-1}, j_t)\}.$$

Finally, (4.10) implies the following formula for the codimension of an orbit $\Omega_{\Delta, M}$ in $\text{Fl}_{\mathbf{a}}(V) \times \text{Fl}_{\mathbf{b}}(V) \times \text{Fl}_{\mathbf{c}}(V)$:

$$\text{codim}(\Omega_{\Delta, M}) = \sum_{k < i, l < j} m_{ij} m_{kl} + \sum_{\{(i, j)\} \not\leq \Delta} m_{ij}. \quad (4.14)$$

ACKNOWLEDGMENTS

We thank Michel Brion and Claus Ringel for helpful references and discussions. This paper was completed during Andrei Zelevinsky's stay at the Institut Fourier, St Martin d'Hères (France) in February–March 1998; he gratefully acknowledges the warm hospitality of Michel Brion and financial support by CNRS, France.

REFERENCES

1. M. Atiyah, On the Krull–Schmidt theorem with application to sheaves, *Bull. Soc. Math. France* **84** (1956), 307–317.
2. K. Bongartz, On degenerations and extensions of finite dimensional modules, *Adv. Math.* **121** (1996), 245–287.
3. K. Bongartz, Degenerations for representations of tame quivers, *Ann. Sci. Éc. Norm. Sup.* **28**, No. 4 (1995), 647–668.
4. M. Brion, Groupe de Picard et nombres caractéristiques des variétés sphériques, *Duke Math. J.* **58** (1989), 397–424.
5. V. Kac, Infinite root systems, representations of graphs and invariant theory, *Invent. Math.* **56** (1980), 57–92.
6. P. Littelmann, On spherical double cones, *J. Algebra* **166** (1994), 142–157.
7. C. Riedtmann, Degenerations for representations of quivers with relations, *Ann. Sci. Éc. Norm. Sup.* **19**, No. 4 (1986), 275–301.
8. C. M. Ringel, Exceptional modules are tree modules, preprint, 1997.
9. C. T. Simpson, Products of matrices, in “Differential Geometry, Global Analysis, and Topology (Halifax, NS, 1990),” CMS Conf. Proc., Vol. 12, Amer. Math. Soc., Providence, RI, 1991.