



Infectio

Asociación Colombiana de Infectología

www.elsevier.es/infectio



LETTER TO THE EDITOR

Epidemiology of Tuberculosis among Homeless Persons, Pereira, Risaralda, Colombia, 2007-2010[☆]

Epidemiología de la tuberculosis entre personas sin techo de Pereira, Risaralda, Colombia, 2007-2010

Tuberculosis (TB) is one of the most important infectious diseases globally and its occurrence among risk groups –those with HIV/AIDS, indigenous populations, and individuals living in crowded conditions and/or extreme poverty such as homeless persons all the more¹⁻⁴. Within the world morbidity and mortality burden of TB, special consideration has been recently given to disease incidence among homeless persons, not only because they can be a risk, a source of transmission and a hub for persistent presence of the pathogen, but also for their varying clinical evolution and outcome⁵⁻⁸. Despite the above, there are few published studies regarding TB among homeless persons in Latin America. Some isolated references can be found at Medline and Scopus from Brazil and Puerto Rico^{9,10}, but none from Colombia. In this country, only approaches to the social aspects¹¹ and public policies regarding TB among this population¹², have been reported.

For these reasons, the objective of this epidemiological study was to assess the occurrence of TB among registered HLP in the Pereira metropolitan area, Colombia, 2007-2010, and design a proposal for improvement of our approach to the disease and its evolution. Pereira is the major city among the Coffee-Triangle (also called Coffee Axis). Morbidity and mortality, as well as clinical-therapeutic evolution of the registered homeless population in the Pereira metropolitan area, Department of Risaralda, Colombia, 2007-2010, at the TB program were analyzed. Clinical, epidemiological and bacteriological diagnoses were made for each case. Cumulated incidence and mortality rates were also analyzed in comparison to those of the general population within the same

municipality. Data was obtained through active and passive collection as part of the TB program, including multiple-source verification supported by the Epidemiological Surveillance System (SIVIGILA), TB trimester program reports and TB treatment cohort reports. Collected data was included in an Excel[®] 2007 for Windows Vista[®] spreadsheet and analyzed with SPSS v.17.0[®].

During the study period, 74 homeless persons with TB were diagnosed, from a total of 1,470 evaluated individuals (registered at the Social Development Secretary of Pereira) (cumulated incidence=50.3 cases/1,000pop., range 5.44 to 22.44; year 2010=16.32). Outcome condition was: treatment self-withdrawal in 43.2% (range 36.4-55.6%), therapeutic failure, 4.1% (0.0-9.1%), deaths, 2.7% (0.0-8.3%), cured, 39.2% (33.3-42.4%) and completed treatments, 10.8% (4.2-25.0%) (fig. 1).

Comparing the 2010 TB incidence among homeless persons with that of the general population (0.6585/1,000pop.) is 76.45 times higher in that risk population. In the same way, case fatality rate in 2010 was 2.5 times higher than in the general population (3.3%). These results have important implications in public health as well in the management and evolution of TB in these individuals, persistence of infection, drug-resistance and potential transmission to those in proximity to these infected persons¹⁻¹². For these reasons, secondary to this report a project was proposed for a Center to attend homeless persons with TB, in order to address the issue through integrated social, epidemiological, clinical, diagnostic and therapeutic management. In the future, we expect to analyze the potential impacts of such interventions on the reduction and better control TB incidence among homeless persons in Pereira, Risaralda, Colombia.

Funding

Cooperativa de Entidades de Salud de Risaralda (COODESURIS) and Asociación Colombiana de Infectología (ACIN). A. J. Rodríguez-Morales and D. M. Castañeda-Hernández were supported from ACIN to participate at the meetings where this work was presented.

[☆] This study was previously presented in part at the XVIII International Congress for Tropical Medicine and Malaria, Rio de Janeiro, Brazil, 23-27 September 2012 (Poster Tb011) and at the VIII National Meeting of Research in Infectious Diseases, Armenia, Colombia, 24-26 May 2012 (Oral presentation #143).




Figure 1 Outcomes for the cohort of homeless persons, Municipality Pereira, Tuberculosis Control Program, 2007-2010.

Conflict of Interests

We declare no conflict of interests.

Acknowledgments

Thanks to the Social Development Office for its cooperation in this study. This study was previously presented in part at the XVIII International Congress for Tropical Medicine and Malaria, Rio de Janeiro, Brazil, 23-27 September 2012 (Poster Tb011) and at the VIII National Meeting for Research in Infectious Diseases, Armenia, Colombia, 24-26 May 2012 (Oral presentation #143).

References

- Centers for Disease Control and Prevention. Tuberculosis outbreak associated with a homeless shelter - Kane county, Illinois, 2007-2011. *MMWR Morb Mortal Wkly Rep.* 2012;61:186-9.
- Abel EK. Taking the cure to the poor: patients' responses to New York City's tuberculosis program, 1894 to 1918. *Am J Public Health.* 1997;87:1808-15.
- Araujo Z, Gonzalez N, de CL, Ziegler RC, de Waard JH, Giampietro F, et al. Levels of complement C3 and C4 components in Amerindians living in an area with high prevalence of tuberculosis. *Mem Inst Oswaldo Cruz.* 2006;101:359-64.
- Chaves NJ, Gibney KB, Leder K, O'Brien DP, Marshall C, Biggs BA. Screening practices for infectious diseases among Burmese refugees in Australia. *Emerg Infect Dis.* 2009;15:1769-72.
- Tan de BJ, Rossi C, Rivest P, Zwertling A, Thibert L, McIntosh F, et al. Tuberculosis and homelessness in Montreal: a retrospective cohort study. *BMC Public Health.* 2011;11:833.
- Khan K, Rea E, McDermaid C, Stuart R, Chambers C, Wang J, et al. Active tuberculosis among homeless persons, Toronto, Ontario, Canada, 1998-2007. *Emerg Infect Dis.* 2011;17:357-65.
- Tabuchi T, Takatorige T, Hirayama Y, Nakata N, Harihara S, Shimouchi A, et al. Tuberculosis infection among homeless persons and caregivers in a high-tuberculosis-prevalence area in Japan: a cross-sectional study. *BMC Infect Dis.* 2011;11:22.
- McAdam JM, Bucher SJ, Brickner PW, Vincent RL, Lascher S. Latent tuberculosis and active tuberculosis disease rates among the homeless, New York, New York, USA, 1992-2006. *Emerg Infect Dis.* 2009;15:1109-11.
- Paton NI, Castello-Branco LR, Jennings G, Ortigao-de-Sampaio MB, Elia M, Costa S, et al. Impact of tuberculosis on the body composition of HIV-infected men in Brazil. *J Acquir Immune Defic Syndr Hum Retrovirol.* 1999;20:265-71.
- Robles RR, Marrero CA, Reyes JC, Colon HM, Matos TD, Finlison HA, et al. Risk behaviors, HIV seropositivity, and tuberculosis infection in injecting drug users who operate shooting galleries in Puerto Rico. *J Acquir Immune Defic Syndr Hum Retrovirol.* 1998;17:477-83.
- Álvarez Gómez M, Correa M, Uribe Ríos M. Percepciones sociales frente a la comunicación del tratamiento y apoyo recibido por la población habitante de calle con tuberculosis. *Medicina UPB.* 2009;20:22-32.
- Tirado Otálvaro AF, Correa Arango ME. Accesibilidad de la población habitante de calle a los programas de promoción y prevención establecidos por la resolución 412 de 2000. *Investig Andina.* 2009;11:23-36.

Diana Milena Castañeda-Hernández^{a,b,d}, Adrián Bolívar-Mejía^c and Alfonso J. Rodríguez-Morales^{d,e,f,*}

^aTuberculosis and Leprosy Program, Secretaría de Salud y Seguridad Social de Pereira, Pereira, Risaralda, Colombia

^bFundación Universitaria del Área Andina, Seccional Pereira, Pereira, Risaralda, Colombia

^cFaculty of Health Sciences, Universidad Industrial de Santander, Bucaramanga, Santander, Colombia

^dFundación para la Consultoría en Salud, Información, Desarrollo Social, Científico y Tecnológico Cenit Colombia, Pereira, Risaralda, Colombia

^eInfection and Immunity Research Group, Faculty of Health Sciences, Universidad Tecnológica de Pereira, Pereira, Risaralda, Colombia

^fOffice of Scientific Research, Cooperativa de Entidades de Salud de Risaralda, Pereira, Risaralda, Colombia

* Corresponding author.

E-mail: arodriguez@utp.edu.co (A.J. Rodríguez-Morales).