

REDOX POTENTIALS OF ELECTRON ACCEPTORS IN PHOTOSYSTEM II $_{\alpha}$ AND II $_{\beta}$

A. P. G. M. THIELEN and H. J. van GORKOM

Department of Biophysics, Huygens Laboratory of the State University, PO Box 9504, 2300 RA Leiden, The Netherlands

Received 30 March 1981; revised version received 19 May 1981

1. Introduction

Redox titrations of the chlorophyll fluorescence yield usually showed two transitions which were ascribed to reduction of the primary acceptor of photosystem II [1–5]. The source of this heterogeneity remained obscure. The two components titrated near 0 mV and near –300 mV and were called Q_H and Q_L, respectively. In [6,7] we have shown that in chloroplasts two different types of system II antennae exist, connected to the α and β centers postulated in [8]. Different electron transport properties of the systems II $_{\alpha}$ and II $_{\beta}$ have also been reported [9,10] and Q $_{\beta}$ was found to have a much higher midpoint potential than Q $_{\alpha}$ [3,11,10]. However, in [12] Q_H was reported not to be related to Q $_{\beta}$, because the latter is reduced already at +120 mV.

We have reinvestigated the problem by measurement and analysis of the light induced fluorescence rise curves of tobacco chloroplasts over a wide redox potential range, in the presence and absence of DCMU. The results confirm that Q $_{\beta}$ is reduced at +115 mV. Apart from a transition around +30 mV, due to some fluorescence quenching by the plastoquinone pool in its oxidized state, all other fluorescence changes were due to changes in the redox state of Q $_{\alpha}$. During titration in the presence of DCMU, Q $_{\alpha}$ was reduced in a single transition around –300 mV. In the absence of DCMU part of Q $_{\alpha}$ may be reduced at much higher potentials, depending on the redox state of the plasto-

quinone pool and of the secondary acceptor, Q₂. A consistent interpretation of all these phenomena is presented, which also explains some apparent contradictions in the literature.

Although most, if not all, of the redox reactions described involve protons directly or indirectly, we did not study and do not discuss their pH-dependence. In the reaction schemes the protons are left out, but all thermodynamic data refer to the molecular structures actually present under the conditions of measurement, at pH 8.3. An extensive discussion on the protonation reactions involved in the analogous part of bacterial photosynthesis may be found in [13].

2. Methods

Dark-adapted tobacco chloroplasts, isolated as in [14], were resuspended in 100 mM glycylglycine, 0.4 M sucrose, 10 mM KCl and 5 mM MgCl₂ (pH 8.3) at 50 μ M chl, determined according to [15]. The potential of the medium was measured by means of a platinum electrode with an Ag/AgCl electrode as reference. The titrations were started at about +300 mV; the potential was lowered by the addition of small amounts of a dithionite solution, followed by an equilibration period of 10–15 min. Except for ferricyanide (5 μ M) no redox mediators were added. The accumulated dithionite additions lowered the pH by at most 0.2 units. During the titration the chloroplasts were kept on ice, in complete darkness, under nitrogen. At each potential a small sample was taken for the measurement of a fluorescence induction curve. Chlorophyll fluorescence was excited by light transmitted by a heat reflecting filter and Corning CS 4-96 and CS 3-67 filters (550–600 nm), of sufficient intensity to reduce all Q $_{\alpha}$ and Q $_{\beta}$ within 2 s, when DCMU (10 μ M) was added. Fluorescence was detected at 685 nm (bandwidth 10 nm).

Abbreviations and symbols: chl, chlorophyll; DCMU, 3-(3',4'-dichlorophenyl)-1,1-dimethylurea; Q, all plastoquinone molecules are denoted by the symbol Q, with a suffix indicating their functional position at a given moment as follows: Q $_{\alpha}$, primary electron acceptor of system II $_{\alpha}$; Q $_{\beta}$, primary electron acceptor of system II $_{\beta}$; Q₂, secondary electron acceptor, equivalent to B and R; Q_p, member of the plastoquinone pool; Q_z, plastoquinone bound to the cytochrome *b*-*f* complex

3. Results and discussion

In the range from +300 to -400 mV the maximum fluorescence yield, F_{\max} , obtained after complete photoreduction of Q_{α} and Q_{β} in the presence of DCMU showed a single increase around +30 mV (pH 8.3) of about 30% (fig.1A, ●). It coincides with the reduction of the plastoquinone pool (see below), which in the oxidized form is known to quench fluorescence to some extent [16]. The initial fluorescence yield, F_i , showed a broad transition around +65 mV and a further increase below -300 mV. The ratio F_{\max}/F_i showed transitions around +115 mV and near -300 mV. Apparently quenching by the plastoquinone pool did not strongly affect this ratio. Analysis of the area over the fluorescence induction curve [9] showed that the two transitions in the F_{\max}/F_i ratio corresponded to the reduction of Q_{α} and Q_{β} , respectively (fig.1B). Chemical reduction of Q_{β} took place in a one-electron reaction with a midpoint




Fig.1. (A) Redox titration of initial (X-X) and maximal (O-O) fluorescence level of chloroplasts in the presence of 10 μ M DCMU, at pH 8.3. The resulting F_{\max}/F_i ratio is given by the squares (right scale). (B) Redox titration of Q_{α} and Q_{β} (Δ , \blacktriangle and \circ , \bullet , respectively) and of k_{α} and initial rate constant of Q_{α} photoreduction (\bullet and \blacktriangle , respectively; right scale). The line accompanying the circles represents a one-electron transition with its midpoint at +115 mV. The circles below 75 mV were calculated assuming an invariant k_{β} .

potential of +115 mV at pH 8.3, in agreement with [12]. It may be noted that in the presence of hydroxylamine a lower midpoint potential was observed [10]. The photoreduction of Q_{β} had exponential kinetics with an invariant rate constant, k_{β} (\bullet). All Q_{α} reduction took place in a broad transition around -300 mV (fig.1B, Δ). Due to energy transfer between the system II_{α} antennae, the absorption cross section and hence the observed rate constant of Q_{α} photoreduction increased with the fraction of Q_{α} reduced. Its initial value (k_{α} \blacktriangle) correspondingly increased upon chemical reduction of Q_{α} . Normal sigmoidal kinetics of fluorescence induction in system II_{α} were observed (in contrast to [17]), except that at negative potentials a small 'tail' occurred. Its kinetics were too fast to be due to Q_{β} and the amplitude and rate constant were apparently proportional to those of the main kinetic phase of Q_{α} photoreduction down to -350 mV. The change in $k_{\alpha,i}$ around +30 mV was not caused by a change in the amount of Q_{α} (fig.1B), but due to quenching by oxidized Q_p , which quenching lowered the quantum yields of Q_{α} photoreduction and fluorescence emission similarly (cf. fig.1A, \bullet).

Without DCMU, F_{\max} did not change with the redox potential (fig.2A, \circ , \bullet) but above +100 mV F_{\max} could only be reached by strong illumination. During the light-induced fluorescence rise the plastoquinone pool, too, was reduced as evidenced by the large area over the induction curve; this extra area titrated around +30 mV (not shown), in agreement with [5].

In the absence of DCMU the high potential transition in the initial fluorescence yield (fig.2A, \square) was not only due to the reduction of Q_{β} and Q_p , as in the titration with DCMU (\blacksquare), but also contained about 20% of the variable fluorescence of system II_{α} , corresponding to about 50% reduction of Q_{α} . The value of 50% was obtained from analysis of the induction curve in the presence of DCMU, and can also be estimated from the data in fig.1. If DCMU was added after setting the redox potential to values between -100 and -200 mV, it caused a dark oxidation of all Q_{α} ($X-X$). The addition of DCMU not only led to a lower F_i but also to a faster light-induced fluorescence rise and in fact the area over the induction curve was essentially unaffected (fig.2B). This shows that in the absence of DCMU, although half of the α centers were closed, the total amount of electrons that could be stored was equal to the total amount of α centers. The similar shape of the kinetics suggests that the




Fig.2. (A) Redox titration of initial (\square, \blacksquare) and maximal (\circ, \bullet) fluorescence yield of chloroplasts in presence (closed symbols) or absence (open symbols) of DCMU, at pH 8.3; ($\times - \times$) addition of DCMU after poisoning the potential of the medium resulted in the initial fluorescence yields; (Δ) titration of chloroplasts with most PS II $_{\alpha}$ centers in the state $Q_{\alpha}Q_2^-$. (B) Kinetics of the area over the fluorescence induction curve at -175 mV, before and after addition of DCMU. In both cases the areas and hence also the electron storage capacities of the reducing side of system II $_{\alpha}$ are of equal size.

fluorescence rise did not require two successive photoreactions but only one, proceeding with a 20% decreased quantum efficiency. Thus we conclude that each system II $_{\alpha}$ center accepted one electron and that initially Q_{α} in each center was in the reduced state half of the time, in equilibrium with the reduced secondary acceptor:



Apparently DCMU strongly shifts the equilibrium to the right. Comparison of the two titrations of F_i in fig.2A shows that reduction of Q_2 takes place at or somewhat below the midpoint potential of Q_p , in agreement with [5,18].

At higher potentials, where Q_p is oxidized, DCMU addition may lead to reduction of Q_{α} . This effect depends on preillumination, being larger after an uneven than after an even number of short saturating flashes, as first described in [19]. Subsequent fluorescence induction revealed no oscillations in the amplitude of the β phase, but a fixed amount of 20% reduced Q_{β} after one or more flashes preillumination (fig.3). The fluorescence oscillation with flash number was totally due to Q_{α} reduction and $k_{\alpha,i}$ (inset) oscillated correspondingly. The reduction of Q_{α} is ascribed to a DCMU-induced return of the electron which is stored every other photoreaction on a secondary acceptor, Q_2 , which is required for the two-electron transport via Q_p [19].

Since the midpoint potential of Q_{β} seems too high for electron transport via Q_p , the absence of a two-electron gate in system II $_{\beta}$ is not surprising. In support of earlier indications [20,21], we have recently shown [22] that the ultraviolet absorption difference spectrum of the reduction of Q_2 to Q_2^- is nearly identical to that of Q_1 ($\alpha + \beta$) and shows the reduction of a plastoquinone to the semiquinone anion. The absorbance measurements also showed that the amount of Q_2 corresponds only to the amount of Q_{α} and, in combination with fluorescence measurements, indicated a quantitative oxidation of Q_2^- by Q_{α} upon addition of DCMU.

Most of the reaction centers of system II $_{\alpha}$ can be brought in the state $Q_{\alpha}Q_2^-$ by illuminating the chloroplasts in the presence of *o*-phenanthroline and NH_2OH and subsequently adding Zn^{2+} , to relieve the inhibitory effect of *o*-phenanthroline on Q_{α}^- oxidation [23]. Titration of chloroplasts pretreated this way (fig.2A, Δ) showed a large increase of F_i around a midpoint potential of 30 mV. Of the about 90% Q_{α}^- produced in this transition only a small part could be reoxidized by adding DCMU, indicating that the fluorescence rise was caused by reduction of $Q_{\alpha}Q_2^-$ to $Q_{\alpha}^-Q_2^{2-}$.

The reduction of Q_{α} and oxidation of Q_{α}^- by Q_2^- upon addition of DCMU suggest that the semiquinone form of Q_2 is unstable in the presence of the inhibitor. This instability is most easily explained in terms of a









Fig. 3. (A) The addition of 10 μM DCMU to chloroplasts, which, with NH₂OH added to prevent back reaction, had been illuminated with a number of saturating flashes (*t*_{1/2} = 3 μs), caused an increase of fluorescence from the (x-x) level to (o-o) level. Thereafter a fluorescence rise curve was recorded; (o-o) the maximum fluorescence level reached. Without preillumination the *F*_{max}/*F*_i ratio was 4.8. The redox potential was +150 mV. (B) Fractions Q_α and Q_β (Δ,▲ and ○,●, respectively) found in the reduced state, plotted against the number of flashes given before addition of DCMU. The behavior of *k*_β (dots) and the initial rate constant of Q_α photoreduction (triangles) are shown in the inset.

competition of DCMU with Q₂ for a common binding site on the reaction center. In [24] evidence is presented for a rapid exchange between Q_p and Q₂ or Q₂²⁻, whereas Q₂⁻ is firmly bound to the reaction center (see also [22]). The reduction of Q_α by Q₂⁻ upon addition of DCMU can then be written as:



and the DCMU-induced oxidation of Q_α⁻ by Q₂⁻ as:




Unless Q₂⁻ is present, deactivation of the S-states is not accelerated by reduction of Q_p [25] and probably does not significantly proceed via back reaction. This finding was ascribed to a large value of *K*₂, but we believe that it is not incompatible with the low value

of *K*₂ found in [5] and by us. Our data show that the semiquinone pair Q_α⁻Q₂⁻ is as stable as Q_αQ₂²⁻ and the activation energy for its oxidation to Q_αQ₂⁻ by back reaction with the S-states may be prohibitively high.

4. Conclusions

At pH 8.3 in the presence of DCMU all Q_α titrates in a single transition around -300 mV. In the absence of DCMU Q_α is reduced together with Q₂ near the midpoint potential of the plastoquinone pool, +30 mV, yielding an equilibrium with equal amounts of Q_αQ₂²⁻ and Q_α⁻Q₂⁻. This mixed state was further reduced to Q_α⁻Q₂²⁻ at -300 mV. The reduction of Q_αQ₂⁻ to Q_α⁻Q₂²⁻ was found to occur also near +30 mV. The reactions at the acceptor side of system II_α upon reductive titration can be summarized as follows (leaving out the protons):



The reductions indicated by the dashed arrows did not occur during titration, but their midpoint potentials follow from $x + y = 2 \times 30 \text{ mV}$ and $y - 300 \text{ mV} = 2 \times 30 \text{ mV}$, so that $x = -300 \text{ mV}$ and $y = +360 \text{ mV}$. Q_α and Q_2 should not be considered as independent components in redox titrations.

In the presence, and presumably also in the absence of DCMU the midpoint potential of Q_β/Q_β^- is $+115 \text{ mV}$ (pH 8.3). In agreement with the high potential of its primary acceptor PS II $_\beta$ electron transport does not proceed via the two electron accumulating plastoquinone Q_2 . Possibly Q_β is in close contact with Q_2 , which has a similar midpoint potential [26], implying that oxidation of Q_β^- does not drive proton translocation across the membrane. The absence of a transition at $+115 \text{ mV}$ in the flash-induced electrochromic change [27] supports this tentative conclusion.

Acknowledgements

Thanks are due to Dr B. R. Velthuys (Baltimore) and Dr J. Amesz for stimulating discussions. This investigation was supported by the Netherlands Foundation for Chemical Research (SON), financed by the Netherlands Organization for the Advancement of Pure Research (ZWO).

References

- [1] Duysens, L. N. M. and Sweers, H. E. (1963) in: *Studies on Microalgae and Photosynthetic Bacteria* (Japanese Society of Plant Physiologists, eds) pp. 353–372, University of Tokyo Press, Tokyo.
- [2] Cramer, W. A. and Butler, W. L. (1969) *Biochim. Biophys. Acta* 172, 503–510.
- [3] Horton, P. H. and Croze, E. (1979) *Biochim. Biophys. Acta* 545, 188–201.
- [4] Malkin, R. and Barber, J. (1979) *Arch. Biochim. Biophys.* 193, 169–178.
- [5] Golbeck, J. H. and Kok, B. (1979) *Biochim. Biophys. Acta* 547, 347–360.
- [6] Thielen, A. P. G. M. and Van Gorkom, H. J. (1981) *Biochim. Biophys. Acta* 635, 111–120.
- [7] Thielen, A. P. G. M., Van Gorkom, H. J. and Rijgersberg, C. P. (1981) *Biochim. Biophys. Acta* 635, 121–131.
- [8] Melis, A. and Homann, P. H. (1975) *Photochem. Photobiol.* 21, 431–437.
- [9] Melis, A. and Homann, P. H. (1976) *Photochem. Photobiol.* 23, 343–350.
- [10] Thielen, A. P. G. M. and Van Gorkom, H. J. (1981) *Proc. 5th Int. Congr. Photosynth.* 1980 (Akoyunoglou, G. ed) Intl. Science Services, Jerusalem, in press.
- [11] Melis, A. (1978) *FEBS Lett.* 95, 202–206.
- [12] Horton, P. H. (1981) *Biochim. Biophys. Acta* 635, 105–110.
- [13] Wraight, C. A. (1979) *Biochim. Biophys. Acta* 548, 309–327.
- [14] Amesz, J., Pulles, M. P. J. and Velthuys, B. R. (1973) *Biochim. Biophys. Acta* 325, 472–482.
- [15] Arnon, D. I. (1949) *Plant Physiol.* 24, 1–15.
- [16] Vernotte, C., Etienne, A.-L. and Briantais, J.-M. (1979) *Biochim. Biophys. Acta* 545, 519–527.
- [17] Horton, P. H. (1981) *Proc. 5th Int. Congr. Photosynth.* 1980 (Akoyunoglou, G. ed) Intl. Science Services, Jerusalem, in press.
- [18] Diner, B. A. (1974) in: *Proc. 3rd Int. Congr. Photosynth.* 1974 (Avron, M. ed) pp. 589–601, Elsevier/North-Holland, Amsterdam, New York.
- [19] Velthuys, B. R. and Amesz, J. (1974) *Biochim. Biophys. Acta* 333, 85–94.
- [20] Pulles, M. P. J., Van Gorkom, H. J. and Willemsen, J. G. (1976) *Biochim. Biophys. Acta* 449, 536–540.
- [21] Mathis, P. and Haveman, J. (1977) *Biochim. Biophys. Acta* 461, 167–184.
- [22] Van Gorkom, H. J., Thielen, A. P. G. M. and Gorren, A. C. F. (1981) in: *Quinones in Energy Conserving Systems* (Trumpower, B. L. ed), Academic Press, New York, in press.
- [23] Bennoun, P. and Li, Y.-S. (1973) *Biochim. Biophys. Acta* 292, 162–168.
- [24] Velthuys, B. R. (1981) in: *Quinones in Energy Conserving Systems* (Trumpower, B. L. ed) Academic Press, New York, in press.
- [25] Diner, B. A. (1977) *Biochim. Biophys. Acta* 460, 247–258.
- [26] Bouges-Bocquet, B. (1980) *FEBS Lett.* 117, 54–58.
- [27] Malkin, R. (1978) *FEBS Lett.* 87, 329–333.