

Mutations in the *JARID1C* Gene, Which Is Involved in Transcriptional Regulation and Chromatin Remodeling, Cause X-Linked Mental Retardation

Lars Riff Jensen,¹ Marion Amende,¹ Ulf Gurok,¹ Bettina Moser,¹ Verena Gimmel,¹ Andreas Tzschach,¹ Andreas R. Janecke,² Gholamali Tariverdian,³ Jamel Chelly,⁴ Jean-Pierre Fryns,⁵ Hilde Van Esch,⁵ Tjitske Kleefstra,⁶ Ben Hamel,⁶ Claude Moraine,⁷ Jozef Gécz,⁸ Gillian Turner,⁹ Richard Reinhardt,¹ Vera M. Kalscheuer,¹ Hans-Hilger Ropers,¹ and Steffen Lenzner¹

¹Max Planck Institute for Molecular Genetics, Berlin; ²Department of Medical Biology and Human Genetics, Innsbruck Medical University, Innsbruck; ³Institute for Human Genetics, University Heidelberg, Heidelberg; ⁴Institut Cochin de Génétique Moléculaire, CNRS/INSERM, CHU Cochin, Paris; ⁵Center for Human Genetics, Clinical Genetics Unit, Leuven, Belgium; ⁶Department of Human Genetics, University Medical Centre, Nijmegen, The Netherlands; ⁷Services de Génétique-INSERM U316, CHU Bretonneau, Tours, France; ⁸Women's and Children's Hospital and the University of Adelaide, Adelaide; and ⁹Genetics of Learning Disability (GOLD) Service, Hunter Genetics, University of Newcastle, New South Wales, Australia

In families with nonsyndromic X-linked mental retardation (NS-XLMR), >30% of mutations seem to cluster on proximal Xp and in the pericentric region. In a systematic screen of brain-expressed genes from this region in 210 families with XLMR, we identified seven different mutations in *JARID1C*, including one frameshift mutation and two nonsense mutations that introduce premature stop codons, as well as four missense mutations that alter evolutionarily conserved amino acids. In two of these families, expression studies revealed the almost complete absence of the mutated *JARID1C* transcript, suggesting that the phenotype in these families results from functional loss of the *JARID1C* protein. *JARID1C* (Jumonji AT-rich interactive domain 1C), formerly known as “SMCX,” is highly similar to the Y-chromosomal gene *JARID1D/SMCY*, which encodes the H-Y antigen. The *JARID1C* protein belongs to the highly conserved ARID protein family. It contains several DNA-binding motifs that link it to transcriptional regulation and chromatin remodeling, processes that are defective in various other forms of mental retardation. Our results suggest that *JARID1C* mutations are a relatively common cause of XLMR and that this gene might play an important role in human brain function.

Introduction

Mental retardation (MR), usually defined as cognitive impairment with an IQ <70, is estimated to affect 2%–3% of the population in industrialized countries (Roelvelde et al. 1997). Etiologically, MR is a very heterogeneous condition that involves environmental, stochastic, and/or genetic factors. Numerous studies of large families containing males with MR have highlighted the importance of the X chromosome. Prevalence estimates for X-linked mental retardation (XLMR) range from 0.56/1,000 males for severe forms to 1.83/1,000 for mild and severe forms (Herbst and Miller 1980; Fishburn et al. 1983). Approximately two-thirds of male patients with XLMR exhibit a nonsyndromic

form of XLMR (NS-XLMR) in which MR is the only clinical manifestation (Fishburn et al. 1983). NS-XLMR is a very heterogeneous condition. With one exception, all of the relevant genes identified until 2002 (*FMR2* [MIM 309548], *OPHN1* [MIM 300127], *PAK3* [MIM 300142], *GDI1* [MIM 300104], *IL1RAPL1* [MIM 300206], *TM4SF2* [MIM 300096], *ARHGEF6* [MIM 300267], *MECP2* [MIM 300005], *FACL4* [MIM 300147], and *ARX* [MIM 300382]; see review by Chelly and Hamel [2003]) showed mutations only in a very small proportion of families with XLMR, suggesting that up to 100 different genes might be involved in NS-XLMR. In an analysis of linkage intervals found in families examined in published and unpublished studies, we have recently shown that mutations involved in NS-XLMR seem to cluster in three different regions of the human X chromosome (Ropers et al. 2003). Two clusters, one at Xq28 and the other at Xp22.1-p21.3, coincide with known MR genes. However, the presence of a broad peak on proximal Xp argued for the existence of unknown genes for NS-XLMR in this region. Therefore, we chose a 7.4-Mb interval in Xp11, flanked by

Received September 7, 2004; accepted for publication November 17, 2004; electronically published December 7, 2004.

Address for correspondence and reprints: Dr. Steffen Lenzner, Max Planck Institute for Molecular Genetics, Ihnestr. 73, D-14195 Berlin, Germany. E-mail: lenzner@molgen.mpg.de

© 2004 by The American Society of Human Genetics. All rights reserved. 0002-9297/2005/7602-0010\$15.00

the genes *ELK1* (MIM 311040) and *ALAS2* (MIM 301300), to systematically screen 47 candidate genes for mutations in families with overlapping linkage intervals. This led to the identification of *PQBP1* (MIM 300463), which encodes the polyglutamine binding protein 1, and *FTSJ1* (MIM 300499), a human homolog of the *Escherichia coli* 2'-O-rRNA methyltransferase *FtsJ/RrmJ* gene, the first two XLMR genes found in Xp11 (Kalscheuer et al. 2003; Freude et al. 2004). In the present study, we report the identification of seven different mutations, in a novel brain-expressed gene, that are associated with familial XLMR.

Subjects and Methods

Patients and Controls

Mutation screening was performed in 31 families with XLMR and overlapping linkage intervals and in 179 unlinked families from the European XLMR Consortium. All patients had a normal karyotype, and fragile X syndrome was ruled out. The control panel included 312 unrelated healthy male controls. Samples from patients and their relatives as well as from controls were obtained after receiving informed consent. A summary of clinical data from all patients with mutations in *JARID1C* is given in table A1 (online only).

In family N017, with four affected males (fig. 1A), linkage studies had mapped the gene defect to the DXS1003-PGK1P1 interval. The patients in this family present with MR and microcephaly. The index patient, III:14, was born on time with a weight of 1,750 g. He was able to walk at the age of 14 mo and spoke his first words at the age of 3 years. At the age of 41 years, at his last clinical examination, his height was 165 cm (<3rd percentile), his weight was 58.8 kg, and his head circumference was 53.8 cm (~3rd percentile). He showed mild dysmorphic features, including small forehead, small eyelashes, prognathism, and small testes. His brother, patient III:2, was examined at the age of 53 years: his height was 154 cm (<3rd percentile), and his head circumference was 53.5 cm (~3rd percentile). He has mild dysmorphic features, including flat philtrum, thin upper lip, prognathism, and a small penis. The third patient, IV:4, was born on time with a weight of 2,500 g. He was able to walk at the age of 2 years and spoke his first words at the age of 3.5 years. His examination at the age of 25 years (height 171 cm [~25th percentile], weight 65 kg, head circumference 54 cm [~3rd percentile]) also showed mild dysmorphic features: small, deep-set eyes; upslanting palpebral fissures; and micrognathia. However, his genitals were normal. At that time, his total IQ was measured to be 70 (verbal 60, performance 86). Clinical data were not available for patient III:16.

Linkage data and clinical findings for family L017 (fig.

1B) have been described elsewhere (Claes et al. 2000). In this family, which has a syndromic form of XLMR, four males in two generations present with severe MR, slowly progressive spastic paraplegia, facial hypotonia, and maxillary hypoplasia. Additional features are aggressive behavior and strabismus.

In family D034 (fig. 1C), the two affected brothers exhibit a severe form of MR. The index patient, II:4, was born after an uneventful pregnancy. Developmental delay was noticed at the age of 1 year. The patient shows strabismus divergens. He understands spoken language well, but the only word he speaks is "yes." At the age of 5 years, electroencephalogram examination revealed abnormalities (groups of sharp theta/delta waves), but epileptic seizures have never been observed. His behavior is characterized by frequent unmotivated aggressive outbursts along with indolence. He has good memory skills and can perform simple tasks such as shopping at the local bakery. His older brother shows an almost identical phenotype. In addition, he exhibits abundant body hair and diastema of the teeth. Because of gallstones, the gallbladders of both brothers had been removed at the age of 20 years and 31 years, respectively. No other medical abnormalities have been noticed.

The two boys in family P018 (fig. 1D) have MR and short stature (patient II:1, body height 105 cm at the age of 6.5 years [<3rd percentile]; patient II:2, body height 91 cm at the age of 4.5 years [<3rd percentile]). Apart from strabismus and myopia in the younger brother (II:2), no other clinical abnormalities have been reported. The nonconsanguineous parents are clinically inconspicuous.

In family N063 (fig. 1E), two brothers and their sister have MR. They share a conspicuous phenotype of severe MR, spasticity, epileptic seizures, short stature, microcephaly, hypermetropia, and small feet. Both males have cryptorchism. Additional symptoms include alopecia areata in patient II:1. The sister (II:3) was described to have an unstable mood disorder. The parents of these children are distantly related.

In family D029, with two affected brothers (fig. 1F), the index patient (II:3) has severe MR. His height (169 cm [~3rd percentile]) and head circumference (59 cm [~97th percentile]) at the age of 22 years were in the normal range. The testis volume was 25 ml (>97th percentile), but there was no expansion of the CGG repeat in the *FMR1* gene. His older brother (II:1) is also affected with a severe form of MR. Height (169 cm [~3rd percentile]), head circumference (58 cm [~97th percentile]), and testis volume at the age of 25 years are normal, and he has no additional abnormalities. The son of a half sister (III:2) is said to suffer from learning impairment, but no clinical details are available.

In family A034 (fig. 1G), four affected males have been reported. Little is known about the early lives of the three




Figure 1 Mutations identified in the *JARID1C* gene in families N017 (A), L017 (B), D034 (C), P018 (D), N063 (E), D029 (F), and A034 (G). The sequence chromatograms of the index patients (*asterisks*) are shown beside the wild-type sequences. The corresponding pedigrees are presented above the chromatograms. Family members who have been analyzed for sequence changes are numbered. For patients with numbers in parentheses, only clinical data were available (no DNA was available for mutation analysis).

affected brothers (III:1, III:2, and III:4) except that development was slow and all were institutionalized in early childhood. Patient III:4 died as a result of accidental drowning at the age of 12 years. The two other brothers (III:1 and III:2), now aged 61 years and 56 years, respectively, reside in a group home. Both had short stature (154 cm [<3 rd percentile] and 156 cm [<3 rd percentile], respectively), with normal head circumferences (57 cm [~ 75 th percentile] and 55 cm [25th–50th percentile], respectively). Both were severely handicapped, and neither spoke more than a few single words. Occasional short outbursts of aggression were reported. In his 30s, patient III:2 started to have generalized seizures about once a year, which sometimes led to hospitalization. Their uncle (II:4) is mildly handicapped and lives at home. Recent re-examination of this patient revealed a very different phenotype, strongly suggesting a different disorder. He was previously known as the “sunny boy” of the family and had a “big head at birth, which had to be carried on a pillow.” He did not walk until the age of 6 years, and, later on, his walk was unsteady and awkward. He had “plenty of speech, which was unclear.” Now 74 years of age, he sits and smiles and is very unsteady on his feet. He is of normal height, but he has a large head circumference of 62 cm (>97 th percentile).

Mutation Analysis by Denaturing High-Performance Liquid Chromatography (DHPLC)

For each patient, the complete *JARID1C* (Jumonji AT-rich interactive domain 1C [MIM 314690]) coding sequence and ~ 40 nucleotides of the neighboring intronic sequences were amplified in 34 independent PCRs from genomic DNA. All primer sequences and PCR product sizes are given in table A2 (online only). In general, PCR amplifications were performed in 20- μ l reaction volumes that contained 100 ng genomic DNA, 1 \times buffer (50 mM KCl, 15 mM Tris-HCl [pH 9.0], 2.5 mM MgCl₂, 0.01% Tween 20), 0.2 μ M of each primer, 100 μ M dNTPs, 0.5 M betaine (Sigma), 1 U *Taq* polymerase, and 0.025 U Pfu polymerase. The PCR profile consists of a touchdown PCR with an initial denaturing step at 96°C for 3 min and then 20 cycles at 95°C for 30 s, 65°C for 30 s (lowered 0.5°C at each cycle), and 72°C for 30 s, followed by 30 cycles at 95°C for 30 s, 55°C for 30 s, and 72°C for 30 s. The PCR was finalized by a 5-min extension at 72°C. PCR products were checked by 1.5% agarose gel electrophoresis before they were subjected to DHPLC by the WAVE nucleic-acid-fragment analysis system (Transgenomic). For DHPLC, we pooled PCR products from two patients. Melting profiles and resolution temperatures were predicted by the Transgenomic WAVEMAKER software, version 4.1. Conditions are available on request. PCR products that gave rise to abnormal elution profiles either were gel purified by use

of the QIAquick gel extraction kit (Qiagen) and then subjected to sequencing or were directly sequenced in both directions by use of BigDye Terminator chemistry (PE Biosystems). Separation was performed on an Applied Biosystems 3730xl DNA Analyzer.

RNA Isolation and Northern Blot Analysis

Total RNA was isolated from patient and control lymphoblastoid cell lines by use of TRIzol (Invitrogen), in accordance with the manufacturer’s protocol. Poly-A⁺ RNAs, which were each obtained from 100 μ g of total RNA by use of Dynabeads (Dyna), were separated in a formaldehyde-containing gel in 1 \times MOPS buffer, transferred to a Hybond N+ membrane, and crosslinked by UV. A human multiple-tissue northern blot was obtained from BioChain. The *JARID1C* (GenBank accession number L25270) and *JARID1D* (GenBank accession number NM_004653) probes corresponding to nucleotides c.4563 to c.5292 (730 bp) and c.4400 to c.5128 (729 bp), respectively, were synthesized using the following primer pairs: *JARID1C* (forward): 5'-GAACCAGAATG-GCTTGGAAC-3' and *JARID1C* (reverse): 5'-GGACTG-AGGTGGTTGTGAGG-3'; *JARID1D* (forward): 5'-CTC-GGAGCTCAGGGATTATG-3' and *JARID1D* (reverse): 5'-GAGTCATGAAATTCTTGTTC-3'. The probes were purified, and their sequences were verified. Subsequently, probes were labeled with ³²P]dCTP and were hybridized to the blots in ULTRAhyb buffer (Ambion). The blots were washed in accordance with the manufacturer’s protocol and were exposed to Fuji medical X-ray films at -80°C for 6 h for up to 8 d by use of intensifying screens. Autoradiograms were scanned using an Epson Expression 1680 Pro scanner. To control for RNA loading, blots were reprobbed with a β -actin probe that was provided by BioChain.

Results

Identification of JARID1C Mutations in Families with XLMR

In a recent analysis of linkage data from published and unpublished studies of families with NS-XLMR, we showed that mutations involved in XLMR seem to cluster in three distinct regions on the X chromosome (Ropers et al. 2003). Therefore, we chose a 7.4-Mb interval on Xp11.23-p11.21, flanked by the genes *ELK1* and *ALAS2*, to systematically screen 47 brain-expressed candidate genes for mutations in families with overlapping linkage intervals. One of these genes, *JARID1C*, was found to be mutated in 2 of 31 families investigated. The *JARID1C* gene contains 26 exons and encodes a member of a conserved protein family, which carries several DNA-binding motifs potentially involved in transcriptional regulation and chromatin remodeling. In family N017, we found a

Table 1**Summary of Sequence Changes in JARID1C Found in Patients with MR**

Nucleotide Change	Amino Acid Variation	Position	Type of Sequence Change	Family Code	No. of Linked Families with Sequence Change (n = 31)	No. of Unlinked Families with Sequence Change (n = 179)	Frequency in Control X Chromosomes
c.202_203insC	R68fsX7	Exon 2	Frameshift mutation	D034	0	1	0/312
c.1162G→C	Ala388Pro	Exon 9	Missense mutation	N017	1	0	0/312
c.1204G→T	Asp402Tyr	Exon 9	Missense mutation	A034	0	1	0/312
c.2080C→T	Arg694Ter	Exon 15	Nonsense mutation	P018	0	1	0/312
c.2092G→A	Glu698Lys	Exon 15	Missense mutation	D029	0	1	0/312
c.2191C→T	Leu731Phe	Exon 15	Missense mutation	L017	1	0	0/312
c.3864G→A	Trp1288Ter	Exon 23	Nonsense mutation	N063	0	1	0/312
c.-239G→A		5' UTR	Intronic alteration		0	1	0/312
c.-90G→A		5' UTR	Intronic alteration		1	0	0/312
c.522+19G→A		Intron 4	Intronic alteration		3	3	9/96
c.564C→A	Lys188Lys	Exon 5	Silent mutation		0	3	3/94
c.1764G→A	Gln588Gln	Exon 13	Silent mutation		0	1	0/312
c.1794C→T	Pro598Pro	Exon 13	Silent mutation		0	2	0/312
c.2243+11G→T		Intron 15	Intronic alteration		0	5	2/96
c.2517-7_8insTAC		Intron 17	Intronic insertion		0	1	0/312




Figure 2 Northern blot analysis of poly-A⁺ RNA isolated from lymphoblastoid cell lines obtained from six patients and five controls. The blot was sequentially hybridized with a cDNA probe corresponding to nucleotides 4563–5292 of *JARID1C* and with a β -actin cDNA probe. Note the significant reduction of *JARID1C* transcript levels in patients from families N063 and P018.

missense mutation, c.1162G→C, that resulted in an alanine-to-proline exchange at amino acid 388 (fig. 1A). Another single-nucleotide substitution (c.2191C→T) was identified in family L017; this substitution changed the amino acid leucine into phenylalanine at position 731 (fig. 1B). Sequencing of PCR-amplified *JARID1C* transcripts from available family members showed that, in both families, the respective mutations cosegregate with the disease (data not shown). None of these changes was found in 312 male controls.

These findings prompted us to analyze 179 additional unrelated patients from smaller families with established or probable XLMR. In five of these families, we found additional mutations, including one frameshift mutation, two nonsense mutations, and two missense mutations. A single-nucleotide insertion of a cytidine (c.202_203insC) introduces a frameshift mutation in patients from family D034 (fig. 1C), eventually resulting in a premature stop codon at position 75 of *JARID1C*. In family P018, the single-nucleotide substitution c.2080C→T replaces the normal arginine by a termination codon at position 694 of the protein (fig. 1D). Another nonsense mutation (Trp1288Ter) was identified in family N063 and is caused by the sequence change c.3864G→A (fig. 1E). In family D029, a guanine-to-adenine mutation (c.2092G→A) results in a glutamic acid-to-lysine exchange at amino acid 698 (Glu698Lys) (fig. 1F). The second missense mutation found in the patient cohort that is present in family A034 (c.1204G→T) changes an aspartic acid residue into tyrosine at position 402 (fig. 1G).

Reinvestigation of the respective sequences in all available affected and unaffected family members confirmed the cosegregation of the mutations with the phenotype

(data not shown). All the mutations that were identified in these families were absent in the remaining 205 analyzed patients with XLMR and also in 312 male controls. In addition to the seven most likely disease-causing mutations, we have found several other sequence variants that were not known before. However, these variants either were silent mutations or were found in intronic sequences and, therefore, for the time being, were not considered as disease causing. All sequence changes found in patients with MR are listed in table 1.

Expression Analysis of *JARID1C* in Patients

To check the stability of all mutated transcripts, we performed northern blot hybridizations of poly-A⁺ cell line RNA from one affected male in each family (except family A034) and five male controls. To avoid cross-hybridization with the highly similar Y-chromosomal gene *JARID1D*, probes were employed that did not show significant sequence similarity to each other upon BLAST2 alignment. *JARID1C* transcripts were almost undetectable in the patients from families P018 and N063, who had premature terminations at codon 694 and codon 1288 (fig. 2), respectively. Thus, the phenotype in these patients most likely results from functional loss of *JARID1C* protein. In contrast, no reduction was observed in the patient from family D034, in whom a frameshift mutation introduces a premature termination at codon 75 of *JARID1C* (fig. 2). In patients with missense mu-




Figure 3 Sequential hybridization of *JARID1C* (A), *JARID1D* (B), and β -actin (C) to a northern blot containing human poly-A⁺ RNAs from adult heart, brain, liver, pancreas, skeletal muscle, and lung. In panel A, probes correspond to nucleotides 4563–5292 of *JARID1C*; in panel B, probes correspond to nucleotides 4400–5128 of *JARID1D*; and, in panel C, probes correspond to a cDNA fragment of β -actin.




Figure 4 Schematic representation of the human *JARID1C* protein, showing the positions and types of mutations found. On the basis of the NCBI Conserved Domain Database, the positions of known domains and motifs are marked by different boxes, which are explained at the bottom of the figure.

tations, *JARID1C* RNA levels were not conspicuously changed.

JARID1C Is Mainly Expressed in Brain and Skeletal Muscle

A predominant transcript of ~6.0 kb was detected in almost all of the human adult tissues investigated, although at very different levels. The strongest expression was observed in brain and skeletal muscle, whereas the lowest expression was seen in heart and liver (fig. 3A). Hybridization of a *JARID1D*-specific probe to the same blot showed expression that was much lower than that of *JARID1C* in general, and expression was seen only in pancreas, lung, brain, and skeletal muscle (fig. 3B).

The *JARID1C* Protein Contains Several Conserved DNA-Binding Domains

JARID1C (GenBank accession number P41229) encodes a 1,560-aa protein that is homologous to three other proteins of the human *JARID1* family. On the basis of a BLAST2 alignment, the amino acid identity of *JARID1C* to *JARID1D* (GenBank accession number NP_004644), *JARID1A* (GenBank accession number NP_005047), and *JARID1B* (GenBank accession number NP_006609) is 85%, 51%, and 47%, respectively. Human *JARID1C* is evolutionarily conserved, with homologs in the mouse (*Jarid1c* [GenBank accession number NP_038696]), fruit fly (*Lid* [GenBank accession number NP_723140]), and nematode (ZK593.4 [GenBank accession number CAA93426]) that show 94%, 42%, and 34% amino acid identity, respectively. A Conserved Domain Database alignment (Marchler-Bauer et al. 2003) revealed several conserved domains between the *JARID* proteins, including a JmjN domain (smart00545), an Arid/Bright domain (smart00501), a JmjC domain (smart00558), a C5HC2 zinc-finger domain (pfam02928), and PHD zinc-finger domains (pfam00628) (fig. 4 and data not shown).

Discussion

We have shown that patients affected with XLMR carry mutations in a novel XLMR gene, *JARID1C*. Mutation screening in 210 unrelated families with XLMR revealed at least seven different sequence alterations that are most likely disease causing—including two nonsense mutations, one frameshift mutation, and four missense mutations. The three mutations that introduce premature termination codons are expected to remove >95%, >55%, and >17% of the normal protein, respectively. Moreover, in the two families with the longest residual transcripts, P018 and N063, *JARID1C* expression was almost undetectable, probably because of the mechanism of nonsense-mediated mRNA decay (NMD) (Holbrook et al. 2004), suggesting a complete loss of *JARID1C* function in these two families. However, the frameshift mutation in family D034, which truncates the *JARID1C* protein at position 75, seems to be insensitive to NMD. Similar findings have been reported elsewhere (Asselta et al. 2001), including intertissue (Bateman et al. 2003) and interindividual (Kerr et al. 2001) variations in NMD efficiency. In family D034, the aberrant phenotype may be a result of either extreme shortening of the protein and deletion of nearly all relevant domains or an abnormal protein produced by alternative usage of a methionine downstream of exon 2.

All missense changes affect amino acids, which are identical in the four different members of the human *JARID1* family and in homologous proteins of mouse and fruit fly, suggesting an important functional role for these amino acids (fig. 5). The missense change Leu731Phe (family L017), which is located in a predicted, evolutionarily conserved C5HC2 zinc-finger domain, might have a profound effect on that domain as a result of the differences in size and chemical properties of leucine and phenylalanine. Although the three other missense changes—Ala388Pro, Asp402Tyr, and Glu698Lys—are not located in a conserved domain,

	Ala388Pro	Asp402Tyr	Glu698Lys	Leu731Phe
Hs_JARID1C	385 FEQ A TREYTLQSFGE M AD S FK 405...695 EAF E LLPDDERQCIKCKTT C FLSALACYDCPDGLV C L S HIN 735			
Hs_JARID1D	375 FEQ A T O EY S LQSFGE M AD S FK 395...685 EAF E LLPDDERQCIKCKTT C FLSALACYDCPDGLV C L S HIN 725			
Hs_JARID1A	354 FEQ A VREYTLQSFGE M AD N FK 374...664 E V F E LV P DDERQ C S A C R TT C FLSAL T C S C N P E R L V C L Y H P T 704			
Hs_JARID1B	507 FEQ A A R DYTL R T F GE M AD A FK 527...817 M D F E LLPDDERQ C V K CKTT C F S A I S C S C K P G L L V C L H H V K 857			
Mm_Jarid1c	385 FEQ A TREYTLQSFGE M AD S FK 405...695 EAF E LLPDDERQCIKCKTT C FLSALACYDCPDGLV C L S HIN 735			
Dm_Lid	509 FEQ A E R EYTLQ F GE M AD O FK 529...818 R A F E L V N D D E R H C O E C N T T C FL S A V A C E C N D K L I V C L R H Y T 858			

Figure 5 Partial amino acid sequence alignment of the four different members of the human JARID1 family (Hs_JARID1A–Hs_JARID1D), the JARID1C orthologous protein of mouse (Mm_Jarid1c), and the homologous Lid protein of fruit fly (Dm_Lid). Amino acids that differ from the JARID1C sequence are highlighted. The four missense mutations and their positions are indicated by arrows and bold letters. The respective amino acids are identical in the six proteins.

they can cause considerable changes in the protein structure. The introduction of a proline residue causes the protein backbone to bend. The replacement of aspartic acid by tyrosine, as well as of glutamic acid by lysine, alters the charge of the protein and, presumably, its three-dimensional structure. Together, our data suggest that the seven *JARID1C* mutations detected affect the function of the protein and are responsible for the cognitive defects found in the patients studied.

Human JARID1C/SMCX protein—which contains several highly conserved domains, including Arid/Bright and Jumonji—belongs to the JARID1 subfamily of ARID DNA-binding proteins (Kortschak et al. 2000; Wilsker et al. 2002). Apart from JARID1C, three other proteins that are characterized by a core ARID of 81 aa—namely, JARID1D/SMCY, JARID1A/RBBP2, and JARID1B/PLU-1—comprise the JARID1 subfamily. JARID1A/RBBP2 was cloned in a search for retinoblastoma protein (pRB)–binding partners (Fattaey et al. 1993). pRB is a well-known and extensively characterized protein that regulates cell proliferation by inhibiting growth-promoting proteins. It can recruit histone deacetylase to chromatin (Luo et al. 1998), which represses the transcription of genes involved in cell-cycle regulation that contain sites for the E2F transcription factor in their promoters (Dyson 1998). The JARID1B protein, also known as “PLU-1,” is a transcriptional repressor. It binds brain factor 1 (BF1), also known as “FOXG1B,” and paired box 9 (PAX9), both of which are developmental transcription factors (Tan et al. 2003). In mouse embryos and human fetal brain, the expression of BF1/FOXG1B is restricted to the neuronal cells in the telencephalon, with strong expression in the developing dentate gyrus and hippocampus. In *Foxg1*-null mice, the generation of the earliest born neurons, the Cajal-Retzius cells, is suppressed (Hanashima et al. 2004).

JARID1C/SMCX and *JARID1D/SMCY* (MIM 426000) constitute an X-Y homologous gene pair, which probably evolved from an identical gene pair on the orig-

inal mammalian (proto-) sex chromosomes (Delbridge et al. 2004), implicating a functional equivalence of *JARID1C* and *JARID1D*. This probable similar function of X and Y copies is strengthened by the fact that the human (as well as the mouse) *JARID1C* gene was thought to escape X inactivation (Agulnik et al. 1994). However, the mouse *Jarid1c* gene exhibits partial X inactivation in embryos, in extraembryonic lineages, and in different adult tissues, showing an expression of 20%–70% from the inactive X allele, compared with the active X allele (Sheardown et al. 1996). Furthermore, *Jarid1c* and *Jarid1d* are expressed in mouse brain in a sex-specific fashion (i.e., in adulthood, *Jarid1c* is expressed at a level that is significantly higher in female brains than in male brains, and the expression of *Jarid1d* in males is not sufficient to compensate for the female bias in X-gene expression [Xu et al. 2002]). Hence, the X-Y gene pair *JARID1C/JARID1D* might not be functionally equivalent, suggesting an indispensable role of *JARID1C* for normal brain function. Together with other differentially regulated X-Y gene pairs (e.g., *USP9X/Y* [MIM 300072 and MIM 400005], *RPS4X/Y* [MIM 312760 and MIM 470000], and *UTX/Y* [MIM 300128 and MIM 400009]), *JARID1C/JARID1D* might also be involved in sex differences in brain development and function and in the prevalence of neuropsychiatric disorders (Vawter et al. 2004).

We have found mutations in the gene *JARID1C* in 7 of 210 families with XLMR, which corresponds to a mutation frequency of 3.3% (CI 0.009–0.057) for XLMR. This number is somewhat biased, since families with XLMR with linkage intervals outside the candidate region in Xp11 have been excluded from our mutation screen. However, the mutation frequency is still 2.8% (CI 0.004–0.052) if only the five changes found in 179 families with proven or probable XLMR are taken into account. Thus, *JARID1C* seems to be one of the more frequently mutated genes in patients with XLMR, and its mutation frequency is comparable to that described for the creatine transporter SLC6A8 (MIM 300036)

(Rosenberg et al. 2004). *JARID1C* mutations cause not only nonsyndromic but also syndromic XLMR, as seen in families N017, L017, and N063; this is in agreement with findings in other XLMR genes (Merienne et al. 1999; Meloni et al. 2000; Bienvenu et al. 2002; Strømme et al. 2002; Kalscheuer et al. 2003).

We conclude that mutations in *JARID1C* contribute significantly to the etiology of XLMR. Further studies will include the characterization of mice deficient in *Jarid1c* and expression profiling of the brain tissues. These investigations should shed more light on the mechanisms by which mutations in *JARID1C* result in dysfunction of the brain.

Acknowledgments

We thank the members of the families and their clinicians, for their participation in this study, and S. Freier and H. Madle, for help with cell cultures. This project was supported by Deutsches Humangenom-Programm (grant 01KW99087), Nationales Genomforschungsnetzwerk (grant 01GR0105), and the 5th European Union Framework (grant QLG3-CT-2002-01810).

Electronic-Database Information

Accession numbers and URLs for data presented herein are as follows:

BLAST2, <http://www.ncbi.nlm.nih.gov/blast/bl2seq/bl2.html>
 Conserved Domain Database, <http://www.ncbi.nlm.nih.gov/Structure/cdd/cdd.shtml>
 GenBank, <http://www.ncbi.nlm.nih.gov/Genbank/> (for *JARID1C* [accession number L25270], *JARID1C* [accession number P41229], *JARID1D* [accession number NM_004653], *JARID1D* [accession number NP_004644], *JARID1A* [accession number NP_005047], *JARID1B* [accession number NP_006609], *Jarid1c* [accession number NP_038696], *Lid* [accession number NP_723140], and *ZK593.4* [accession number CAA93426])
 Human Genome Browser Gateway, <http://genome.cse.ucsc.edu/cgi-bin/hgGateway?db=hg10>
 Online Mendelian Inheritance in Man (OMIM), <http://www.ncbi.nlm.nih.gov/entrez/Omim/> (for *FMR2*, *OPHN1*, *PAK3*, *GDI1*, *IL1RAPL1*, *TM4SF2*, *ARHGEF6*, *MECP2*, *FACL4*, *ARX*, *ELK1*, *ALAS2*, *PQB1*, *FTSJ1*, *JARID1C*, *JARID1D*, *SMCY*, *USP9X/Y*, *RPS4X/Y*, *UTX/Y*, and *SLC6A8*)

References

Agulnik AI, Mitchell MJ, Mattei MG, Borsani G, Avner PA, Lerner JL, Bishop CE (1994) A novel X gene with a widely transcribed Y-linked homologue escapes X-inactivation in mouse and human. *Hum Mol Genet* 3:879–884
 Asselta R, Duga S, Spena S, Santagostino E, Peyvandi F, Pisseddu G, Targhetta R, Malcovati M, Mannucci PM, Tenchini ML (2001) Congenital afibrinogenemia: mutations

leading to premature termination codons in fibrinogen A α -chain gene are not associated with the decay of the mutant mRNAs. *Blood* 98:3685–3692

- Bateman JF, Freddi S, Natrass G, Savarirayan R (2003) Tissue-specific RNA surveillance? Nonsense-mediated mRNA decay causes collagen X haploinsufficiency in Schmid metaphyseal chondrodysplasia cartilage. *Hum Mol Genet* 12:217–225
 Bienvenu T, Poirer K, Friocourt G, Bahi N, Beaumont D, Fauchereau F, Ben Jeema L, Zemni R, Vinet MC, Francis F, Couvert P, Gomot M, Moraine C, Van Bokhoven H, Kalscheuer V, Frints S, Géczy E, Ohzaki K, Chaabouni H, Fryns JP, Desportes V, Beldjord C, Chelly J (2002) ARX, a novel Prd-class-homeobox gene highly expressed in the telencephalon, is mutated in X-linked mental retardation. *Hum Mol Genet* 11:981–991
 Chelly J, Hamel BCJ (2003) Genetics of X-linked mental retardation. In: Fisch GS (ed) *Genetics and genomics of neurobehavioral disorders*. Humana Press, Totowa, NJ, pp 263–287
 Claes S, Devriendt K, Van Goethem G, Roelen L, Meireleire J, Raeymaekers P, Cassiman J-J, Fryns J-P (2000) Novel syndromic form of X-linked complicated spastic paraplegia. *Am J Med Genet* 94:1–4
 Delbridge ML, Longepied G, Depetris D, Mattei MG, Disteché CM, Marshall Graves JA, Mitchell MJ (2004) TSPY, the candidate gonadoblastoma gene on the human Y chromosome, has a widely expressed homologue on the X: implications for Y chromosome evolution. *Chromosome Res* 12:345–356
 Dyson N (1998) The regulation of E2F by pRB-family proteins. *Genes Dev* 12:2245–2262
 Fattaey AR, Helin K, Dembski MS, Dyson N, Harlow E, Vuocolo GA, Hanobik MG, Haskell KM, Oliff A, Defeo-Jones D, Jones RE (1993) Characterization of the retinoblastoma binding proteins RBP1 and RBP2. *Oncogene* 8:3149–3156
 Fishburn J, Turner G, Daniel A, Brookwell R (1983) The diagnosis and frequency of X-linked conditions in a cohort of moderately retarded males with affected brothers. *Am J Med Genet* 14:713–724
 Freude K, Hoffmann K, Jensen LR, Delatycki MB, Des Portes V, Moser B, Hamel B, Van Bokhoven H, Moraine C, Fryns J-P, Chelly J, Géczy J, Lenzner S, Kalscheuer VM, Ropers H-H (2004) Mutations in the FTSJ1 gene coding for a novel S-adenosyl-methionine-binding protein cause nonsyndromic X-linked mental retardation. *Am J Hum Genet* 75:305–309
 Hanashima C, Li SC, Shen L, Lai E, Fishell G (2004) Foxg1 suppresses early cortical cell fate. *Science* 303:56–59
 Herbst DS, Miller JR (1980) Nonspecific X-linked mental retardation II: the frequency in British Columbia. *Am J Med Genet* 7:461–469
 Holbrook JA, Neu-Yilik G, Hentze MW, Kulozik AE (2004) Nonsense-mediated decay approaches the clinic. *Nat Genet* 36:801–808
 Kalscheuer VM, Freude K, Musante L, Jensen LR, Yntema HG, Géczy J, Sefiani A, et al (2003) Mutations in the polyglutamine binding protein 1 gene cause X-linked mental retardation. *Nat Genet* 35:313–315
 Kerr TP, Sewry CA, Robb SA, Roberts RG (2001) Long mutant

- dystrophins and variable phenotypes: evasion of nonsense-mediated decay? *Hum Genet* 109:402–407
- Kortschak RD, Tucker PW, Saint R (2000) ARID proteins come in from the desert. *Trends Biochem Sci* 25:294–299
- Luo RX, Postigo AA, Dean DC (1998) Rb interacts with histone deacetylase to repress transcription. *Cell* 92:463–473
- Marchler-Bauer A, Anderson JB, DeWeese-Scott C, Fedorova ND, Geer LY, He S, Hurwitz DI, et al (2003) CDD: a curated Entrez database of conserved domain alignments. *Nucleic Acids Res* 31:383–387
- Meloni I, Bruttini M, Longo I, Mari F, Rizzolio F, D'Adamo P, Denvriendt K, Fryns JP, Toniolo D, Renieri A (2000) A mutation in the Rett syndrome gene, MECP2, causes X-linked mental retardation and progressive spasticity in males. *Am J Hum Genet* 67:982–985
- Merienne K, Jacquot S, Pannetier S, Zeniou M, Bankier A, Géczy J, Mandel JL, Mulley J, Sassone-Corsi P, Hanauer A (1999) A missense mutation in RPS6KA3 (RSK2) responsible for non-specific mental retardation. *Nat Genet* 22:13–14
- Roeleveld N, Zielhuis GA, Gabreels F (1997) The prevalence of mental retardation: a critical review of recent literature. *Dev Med Child Neurol* 39:125–132
- Ropers H-H, Hoeltzenbein M, Kalscheuer V, Yntema H, Hamel B, Fryns J-P, Chelly J, Partington M, Géczy J, Moraine C (2003) Nonsyndromic X-linked mental retardation: where are the missing mutations? *Trends Genet* 19:316–320
- Rosenberg EH, Almeida LS, Kleefstra T, deGrauw RS, Yntema HG, Bahi N, Moraine C, Ropers H-H, Fryns J-P, deGrauw TJ, Jakobs C, Salomons GS (2004) High prevalence of SLC6A8 deficiency in X-linked mental retardation. *Am J Hum Genet* 75:97–105
- Sheardown S, Norris D, Fisher A, Brockdorff N (1996) The mouse Smcx gene exhibits developmental and tissue specific variation in degree of escape from X inactivation. *Hum Mol Genet* 5:1355–1360
- Strømme P, Mangelsdorf ME, Shaw MA, Lower KM, Lewis SM, Bruyere H, Lutchera V, Gedeon AK, Wallace RH, Scheffer IE, Turner G, Partington M, Frints SG, Fryns JP, Sutherland GR, Mulley JC, Géczy J (2002) Mutations in the human ortholog of *Aristaless* cause X-linked mental retardation and epilepsy. *Nat Genet* 30:441–445
- Tan K, Shaw AL, Madsen B, Jensen K, Taylor-Papadimitriou J, Freemont PS (2003) Human PLU-1 has transcriptional repression properties and interacts with the developmental transcription factors BF-1 and PAX9. *J Biol Chem* 278:20507–20513
- Vawter MP, Evans S, Choudary P, Tomita H, Meador-Woodruff J, Molnar M, Li J, Lopez JF, Myers R, Cox D, Watson SJ, Akil H, Jones EG, Bunney WE (2004) Gender-specific gene expression in post-mortem human brain: localization to sex chromosomes. *Neuropsychopharmacology* 29:373–384
- Wilsker D, Patsialou A, Dallas PB, Moran E (2002) ARID proteins: a diverse family of DNA binding proteins implicated in the control of cell growth, differentiation, and development. *Cell Growth Differ* 13:95–106
- Xu J, Burgoyne P, Arnold AP (2002) Sex differences in sex chromosomes gene expression in mouse brain. *Hum Mol Genet* 11:1409–1419