

used to test this assumption and refine parameters in the model that are otherwise difficult to measure directly, such as the volume occupied by PEG inside the pore and the local concentration of electrolyte, found to be approximately half the bulk value. MD simulations are also used to test a central hypothesis in the theoretical model that PEG complexes cations to acquire a net positive charge. We confirm that this is indeed the case and that five PEG subunits participate in forming crown-ether like sub-structures with a single cation. The refined theoretical model is then fit to blockade depth and residence time values measured experimentally as a function of PEG size (varying from 1000 g/mol to 2000 g/mol). We find that the theoretical predictions of the model agree quantitatively with experiment thereby validating its assumptions.

#### 2679-Pos Board B698

##### Integration of Biogenic Nanopore Membranes on Prefabricated Fluidic Support Substrates

Michael Goryll, Kai-Chun Lin, Xiaofeng Wang, Shankar Ramakrishnan, Melissa Gosse, Sanwip K. Dey, B.L. Ramakrishna.  
Arizona State University, Tempe, AZ, USA.

Marine diatoms provide an alternative to machined silica nanopores, avoiding costly and slow throughput fabrication steps, while being able to achieve pore structures with diameters on the order of 40 nm. The hierarchical pore architecture makes these biogenic nanomembranes exceptionally mechanically stable, while maintaining a short pore length and a high porosity. The most prominent issue when replacing machined silica nanopore membranes with biogenic membranes is the initial random placement of the membranes on the solid substrate. This is also problematic when trying to accomplish a permanent fluidic seal around the membrane.

In our study, we demonstrate the ability to localize and immobilize a 200- $\mu$ m-diameter biomaterialized nanopore membrane structure from marine algae, *Coscinodiscus wailesii*, on pre-defined positions on micro-machined silicon substrates. The substrates feature micron-sized through-wafer channels that allow easy access to the nanopore membrane. Localization of the membrane structure is accomplished using patterning of 8  $\mu$ m thick hydrophobic resin. The addition of poly-L-lysine to the surface before solution-depositing the nanopore membranes results in a strong electrostatic binding force between the oxidized silicon platform and the diatom membranes. Lift-off of the photoresist in acetone removes randomly placed nanopore membranes on the resist-coated area, not affecting diatoms adhering to the silicon surface.


While poly-L-lysine provides an initial fluidic seal, permanent immobilization is accomplished by using UV-curable photoresist SU-8 and proximity photo lithography. Scanning electron micrographs after processing show intact diatoms without the presence of stress cracks. While initial electrochemical measurements indicate that some of the nanopores are clogged by residual epoxy resist after development, subsequent sulfuric-peroxide mixture (SPM) treatment removes the residual resist. Successful translocation experiments using polystyrene beads shows presence of unclogged pores, also indicating that the pore size of the biogenic silica nanomembranes can be modified by chemical treatment.

#### 2680-Pos Board B699

##### Mixed Company in a Protein Nanopore: Transient Double Occupancy Enables Direct Exchange of Pore Ligand in Nanopore-Based Single Molecule Sensing

Anne K. Schuler, Gerhard Baaken, Jan C. Behrends.

University of Freiburg, Institute of Physiology, Freiburg, Germany.  
Partitioning of poly(ethylene-glycol) (PEG) into an alpha-hemolysin nanopore gives rise to long-lasting blockades ( $\tau \approx 0.1$ -10 ms) of ionic current. The depth of blockade shows exquisite sensitivity for polymer length, yielding mass spectra with single monomer resolution in the range between MW  $\approx$  700-2200 or 15-50 repeat units (r.u.). Unexpectedly, high-resolution recordings of single PEG blocking events using a mixture of two monodisperse species unequivocally identified direct transitions between levels corresponding to 28 and 32 r.u. without an intervening unblocked interval. Closer analysis revealed that these occur by the intermediary of shorter, deeper blocks. Based on statistics and current amplitude distributions, we are able to identify three such deeply blocked states, each corresponding to the simultaneous presence in the pore of two ligands: either 2xPEG-28, 2xPEG-32 or PEG-28+PEG-32 (mixed double occupancy). Direct PEG28- $\rightarrow$ PEG32 transitions (or vice versa) are observed only with an intervening mixed occupancy block, which, however, can also result in return to the first blocked level. We conclude that the alpha-hemolysin pore is capable of accommodating two PEG oligomers, with ligand exchange occurring by displacement and translocation of the first blocker.



#### 2681-Pos Board B700

##### Nanopores with Fluid Walls for Determining the Shape, Dipole Moment, and Rotational Diffusion Coefficient of Non-Spherical Proteins

Erik C. Yusko<sup>1</sup>, Brandon R. Bruhn<sup>1</sup>, Ryan C. Rollings<sup>2</sup>, Jiali Li<sup>2</sup>, David Sept<sup>1</sup>, Michael Mayer<sup>1</sup>.

<sup>1</sup>University of Michigan, Ann Arbor, MI, USA, <sup>2</sup>University of Arkansas, Fayetteville, AR, USA.

Recording ionic current through electrolyte-filled nanopores during the passage of proteins is an emerging technique for characterizing unmodified proteins in their native, aqueous environment. By measuring the reduction in current,  $\Delta I$ , during the translocation of single proteins through an electrolyte-filled nanopore, this technique can characterize the size, charge, conformation, assembly and activity of hundreds of unlabeled proteins per second. For non-spherical proteins, however, broad distributions of  $\Delta I$  values make estimates of protein size unreliable. This work employs lipid-bilayer coated nanopores and describes quantitative procedures for determining the shape and volume of single spherical and non-spherical proteins from distributions of  $\Delta I$  values. Since the  $\Delta I(t)$  signal is related to the orientation of non-spherical proteins in the nanopore, individual resistive-pulses can be used to determine the rotational diffusion coefficient and dipole moment of non-spherical proteins while in the nanopore. Moreover, this method has the potential to detect transient changes in the conformation of flexible proteins (e.g. an IgG antibody). This work extends the power of nanopores for characterizing proteins by adding the parameters of shape, volume, rotational diffusion coefficient, and dipole moment of non-spherical proteins to those that can already be determined in a single experiment such as the volume of spherical proteins, charge, and affinity for a ligand.

#### 2682-Pos Board B701

##### Threading Immobilized DNA through a Solid-State Nanopore with a Tip

Changbae Hyun, Ryan Rollings, Harpreet Kaur, Jiali Li.

University of Arkansas, Fayetteville, AR, USA.

Nanopores have been of interest for scientific research in addition to medical applications since they have the ability to detect and characterize single biomolecules with potentially high throughput and low cost. The Scanning Probe Microscope (SPM) method has sub-nanometer spatial resolution. We have constructed a combined SPM-solid state nanopore apparatus to study the capture and release process of lambda-DNA by a voltage biased solid-state nanopore. By tethering the DNA to an fiber tip in ionic solution, we can control the position of one end of the DNA molecule precisely, allowing us to study the DNA capture and release distance from the nanopore. We also have detected DNA sticking to the nanopore mouth without translocation through, it produced small current blockage, and we can study this process with one DNA molecule repetitively. This tethered DNA nanopore sensing method will provide a means to slow DNA translocation, allowing more detailed features of single DNA molecules to be studied, and potentially can be used with all types of nanopores with single-biomolecule sensitivity at controlled translocation rates.

#### 2683-Pos Board B702

##### Brownian Dynamics Simulations of DNA Interaction with a Nanoporous Solid State Membrane

Maria Gracheva, Ining Amy Amy, Christopher McKinney,

Tetiana Timchenko, Zachery Hulings, Dmitriy Melnikov.

Clarkson University, Potsdam, NY, USA.

We have developed computational model that allows us to study the influence of a nanoporous solid state membrane on the dynamics of a biomolecule. We apply various electrolyte and membrane biases and monitor the effects on DNA translocation and extension. The translocation of DNA through a nanopore in a single layered doped semiconductor membrane is studied. With our single layered, electrically tunable membranes, the DNA translocation time can be varied by more than one order of magnitude. Nanopore functionalization is also studied by fixing one strand of DNA to the inner surface of the pore. Two different models of DNA molecule are developed. The first model represents each DNA nucleotide as a single bead, while in the second one we consider two beads per nucleotide: one bead representing the phosphate and sugar backbone, and the other being the base. This model is more realistic and allows us to better understand the principles of interaction between the semiconductor membrane and DNA nucleotides.

#### 2684-Pos Board B703

##### Ligand-Targeted Binding of a Novel Silicone Magnetic Microsphere

Julia C. Ronecker, Benjamin A. Evans.

Elon University, Elon, NC, USA.

In ligand-targeted drug delivery, a carrier particle conjugated with a ligand binds preferentially to an overexpressed receptor on the membrane of a specific cell type. A therapeutic agent is adsorbed onto or absorbed within the carrier, and its release is often triggered by magnetic stimulation or other means. In