

Scara5 Is a Ferritin Receptor Mediating Non-Transferrin Iron Delivery

Jau Yi Li,^{1,6} Neal Paragas,^{1,6} Renee M. Ned,^{2,3} Andong Qiu,¹ Melanie Viltard,¹ Thomas Leete,¹ Ian R. Drexler,¹ Xia Chen,¹ Simone Sanna-Cherchi,¹ Farah Mohammed,¹ David Williams,¹ Chyuan Sheng Lin,¹ Kai M. Schmidt-Ott,^{1,4} Nancy C. Andrews,⁵ and Jonathan Barasch^{1,*}

¹Renal Division, College of Physicians & Surgeons, Columbia University, New York, NY 10032, USA

²McKing Consulting Corporation, 2900 Chamblee Tucker Road, Building 10, Atlanta, Georgia 30341, USA

³National Office of Public Health Genomics, Center for Disease Control and Prevention, Atlanta, GA 30329, USA

⁴Department of Nephrology and Hypertensiology, Charité Berlin, Campus Buch, Max-Delbrück Center for Molecular Medicine, 13125 Berlin, Germany

⁵Department of Pediatrics and Pharmacology and Cancer Biology, Duke University School of Medicine, Durham, NC 27708, USA

⁶These authors contributed equally to this work.

*Correspondence: jmb4@columbia.edu

DOI 10.1016/j.devcel.2008.12.002

SUMMARY

Developing organs require iron for a myriad of functions, but embryos deleted of the major adult transport proteins, transferrin or its receptor transferrin receptor1 (TfR1^{-/-}), still initiate organogenesis, suggesting that non-transferrin pathways are important. To examine these pathways, we developed chimeras composed of fluorescence-tagged TfR1^{-/-} cells and untagged wild-type cells. In the kidney, TfR1^{-/-} cells populated capsule and stroma, mesenchyme and nephron, but were underrepresented in ureteric bud tips. Consistently, TfR1 provided transferrin to the ureteric bud, but not to the capsule or the stroma. Instead of transferrin, we found that the capsule internalized ferritin. Since the capsule expressed a novel receptor called Scara5, we tested its role in ferritin uptake and found that Scara5 bound serum ferritin and then stimulated its endocytosis from the cell surface with consequent iron delivery. These data implicate cell type-specific mechanisms of iron traffic in organogenesis, which alternatively utilize transferrin or non-transferrin iron delivery pathways.

INTRODUCTION

Iron travels from the placenta to embryonic organs bound to carrier proteins, such as transferrin, which is detectable in extracellular fluids by mid-gestation in rodents (Gustine and Zimmerman, 1973). Transferrin is then captured throughout the embryo by the transferrin receptor (TfR1). This pathway is essential for development, since deletion of *TfR1* resulted in embryonic death at E11-E12 (Levy et al., 1999). Indeed, during kidney development, transferrin was found to be both necessary and sufficient to stimulate tubulogenesis in vitro, suggesting a critical role for transferrin-mediated iron transport in nephrogenesis (Ekblom et al., 1983) and by extension perhaps to the development of other organs. The widespread expression of TfR1 (Aisen, 2004)

further reinforced the view that transferrin plays a critical role throughout the embryo.

A detailed analysis of specific embryonic lineages, however, suggests a more restricted view of the transferrin pathway. Transferrin is necessary for thymic maturation (Brekemans et al., 1994; Ned et al., 2003; Macedo et al., 2004) but this requirement is stage specific, since TfR1 neutralizing antibodies block the progression of double negative thymocytes, but fail to inhibit earlier stages of the lymphoid lineage, despite the fact that TfR1 is expressed at all stages. Likewise, while TfR1^{-/-} embryos (Levy et al., 1999) have specific defects in erythroid, lymphoid, and spinal cord development, most other organs appear to initiate development. Chimeras composed of TfR1^{-/-} and wild-type ES cells can live into adulthood, and TfR1^{-/-} cells can be detected by Southern blots in a variety of organs, confirming the notion that different cells can arise from TfR1^{-/-} ES cells in vivo (Ned et al., 2003). Similarly, apparently normal organogenesis occurs in hypotransferrinemic mice^{hpx/hpx} (Huggenvik et al., 1989; Trenor et al., 2000) and humans (Hayashi et al., 1993; Hamill et al., 1991), despite severe anemia. Hence, while specific stages of the hematopoietic lineage (Kanayasu-Toyoda et al., 1999; Sposi et al., 2000; Wingert et al., 2004), bone (Gentili et al., 1994), and nervous system (Siddappa et al., 2002; Levy et al., 1999) depend on TfR1, other cells in the embryo may capture iron by a non-transferrin mechanism. However, few non-transferrin iron trafficking molecules are known, and knockouts of non-transferrin carriers including *Lcn2* (Flo et al., 2004), and iron uptake proteins such as *DMT1* (Gunshin et al., 2005) and *Steap3* (Ohgami et al., 2005), do not grossly disrupt organogenesis. These data leave many fundamental questions unresolved, such as the identity of other iron carriers, their receptors that deliver iron into an appropriate intracellular compartment, and whether different pathways of iron delivery are cell-specific.

Ferritin is a major iron storage protein. Intracellular ferritin is composed of 24 subunits of L- and H-chains in varied proportion (Arosio and Levi, 2002). On the other hand, extracellular (serum) ferritin is composed mostly, but not exclusively of L-ferritin (Linder et al., 1996; Ghosh et al., 2004). Secretion of L-ferritin has been carefully documented in cell cultures (Tran et al., 1997; Ghosh et al., 2004) and in syndromes which overexpress




Figure 1. Initial Stages of Renal Development Were Preserved in $Tfr1^{-/-}$ Mice

(A and B) (A) $Pax2^{+}$ Wolffian ducts and mesonephros and (B) $Pax2^{+}$ ureteric buds surrounded by mesenchymal cells were detected in $Tfr1^{-/-}$ E11 embryos. Crosses with $Hox\beta7$ -GFP mice distinguished the ureteric bud ($GFP^{+}Pax-2^{+}$) from mesenchyme ($GFP^{-}Pax-2^{+}$). Bar = 10 μm .

L-chain, such as heterozygous H chain-deficient mice (Ferreira et al., 2001), *hpx/hpx* mice (Simpson et al., 1991; Dickinson and Connor, 1995), and humans with hyperferritinemia and cataracts (Beaumont et al., 1995). Cytokine stimulation of some cell types also induces H-ferritin secretion (Tran et al., 1997), particularly from macrophages (Chen et al., 2005; Recalcati et al., 2008). While the two chains of ferritin have complementary functions [H-ferritin is a ferroxidase (Levi et al., 1988) and L-ferritin induces polynucleation of iron (Levi et al., 1994)], both ferritins can independently incorporate iron into a core. For example, while serum-ferritin is iron-poor compared to intracellular ferritin, it contains 7 iron atoms/molecule (Watanabe et al., 2001), similar to bacterially expressed L-ferritin, which contains 5–15 iron atoms/molecule (Levi et al., 1989, 1992). In the presence of even a single H chain, up to 4500 iron atoms can be incorporated into L-ferritin cages.

Extracellular ferritin can deliver iron to a variety of cells. Ferritin can bind cell surface receptors, undergo endocytosis, and transfer of iron in sufficient quantities to alter iron-dependent gene expression (Ponka et al., 1998). A receptor for H-ferritin has been identified, Tim2 (or Timd2; Chen et al., 2005), but its expression is highly restricted in the embryo. In contrast, receptors for L-ferritin have not been documented, but may in fact be particularly relevant in the embryo, whose serum ferritin concentration is 3–8-fold greater than that of maternal blood (Bratliid and Moe, 1980; Hussain et al., 1977; MacPhail et al., 1980; Puolakka et al., 1980).

To clarify the mechanisms of iron delivery in embryogenesis, we developed an in vivo assay to evaluate the requirement for TfR1. We focused on the embryonic kidney where cell type progression is dependent on iron delivery (Ekblom et al., 1983). Using chimeric embryos composed of green fluorescent protein (GFP)-labeled $Tfr1^{-/-}$ cells ($GFP-Tfr1^{-/-}$ cells) and unlabeled wild-type cells, we found that mesenchymal, stromal, and capsule compartments were robustly populated by $GFP-Tfr1^{-/-}$ cells and accumulated iron independently of TfR1. Capsular cells expressed *Scara5* (Jiang et al., 2006), which we found bound extracellular ferritin, mediated its endocytosis,

and captured its iron. Tips of the branching ureteric bud on the other hand were poorly populated by $GFP-Tfr1^{-/-}$ cells, confirming a cell type-specific, rather than a global requirement, for transferrin-TfR1. These data imply that different mechanisms deliver iron to different cell types in the developing kidney.

RESULTS

Evidence for Transferrin and Non-Transferrin Iron Delivery in Kidney Development

The developing kidney is generated from three cellular lineages, epithelial (including the nephric duct, ureteric bud [UB], and collecting ducts), metanephric mesenchyme (composed of cap mesenchyme, pretubular aggregate, renal vesicle, C- and S-shaped bodies, and nascent nephron) and a stromal compartment (composed of the capsule, cortical and medullary stroma). The ureteric bud generates a branched collecting duct system. The mesenchymal lineage converts to epithelia and gives rise to both mesonephric and metanephric nephrons, containing glomeruli and epithelial tubules. Capsular and stromal cells form the outermost rings of cells in the kidney cortex and are composed of a variety of cells that have yet to be fully defined (Dressler, 2006; Hatini et al., 1996; Levinson and Mendelsohn, 2003; Schmidt-Ott et al., 2006).

To investigate pathways of iron delivery in the developing kidney, we first inspected $Tfr1^{+/-} \times Tfr1^{+/-}$ crosses (Levy et al., 1999; Ned et al., 2003). We examined 210 E11 embryos recovering $Tfr1^{+/+}$ (28%), $Tfr1^{+/-}$ (52%), and $Tfr1^{-/-}$ (20%) in approximate Mendelian ratios. $Tfr1^{-/-}$ embryos contained the Wolffian duct and associated mesonephric tubules (Figure 1A). The embryos also developed ureteric buds from the Wolffian duct (marked by creating $Tfr1^{-/-}; Hoxb7-GFP^{+}$ lines; Srinivas et al., 1999) as well as cap mesenchyme (Figure 1B). In rare, surviving embryos at E11.5, elongated T-shaped UBs and renal vesicles were found (Figure S1, see the Supplemental Data available online), but more advanced ureteric branching and mesenchymal-epithelial development could not be assayed due to the lethality of TfR1 knockout.




Figure 2. Development of GFP-TfR1^{-/-} Cells in E15 Kidney

(A) GFP-TfR1^{-/-} cells were readily detected in capsule and stroma, mesenchyme, renal vesicles, and S-shaped bodies, but detected with limited frequency in UB tips (bar = 10 μ m).

(B) GFP-TfR1^{-/-} cells (42%) reside in the capsule, interstitium, and podocalyxin⁺ vascular progenitors (confocal section; bar = 10 μ m).

(C and D) GFP-TfR1^{-/-} cells (58%) also resided in proximal and distal nephrons (Ecadherin⁺; dark blue), in podocytes (podocalyxin⁺; red, Glm = glomerulus) and in the UB stalk (Ecadherin⁺, dark blue; Troma⁺, pink). In some cases, entire segments of the S-shaped body were composed of GFP-TfR1^{-/-} cells (confocal section; bar = 10 μ m).

(E and F) Genes specific for UB tip or mesenchymal-stromal compartments were assayed in GFP-TfR1^{-/-} and wild-type cells and presented as the log₂ ratio of microarray values. UB tip-specific genes were under represented in GFP-TfR1^{-/-} cells, whereas mesenchymal-stromal genes were expressed.

(G) Both GFP-TfR1^{-/-} and wild-type cells (two sections are shown) contained cytoplasmic ferritin.

(H) Wild-type cells captured Alexa568-transferrin, but neighboring GFP-TfR1^{-/-} were unlabeled. Bar = 10 μ m.

To determine whether TfR1^{-/-} renal progenitors were competent to form organotypic structures, we used an in vitro culture system. As shown in Figure S2, when TfR1^{-/-} urogenital sheets (E11) were explanted and the media were supplemented with ferric ammonium citrate (20 μ g/ml), a TfR1-independent iron donor, the sheet consistently generated highly branched ureteric buds surmounted by cap mesenchyme (n = 21 urogenital sheets), whereas culture with transferrin resulted in regression of the tissue. This implied that urogenital progenitors, present in vivo when the sheet was explanted (E11) remained competent to form kidney-specific structures despite the deletion of TfR1.

To test the functional requirement for transferrin in vivo beyond the E11 stage, we created chimeric embryos by injecting TfR1^{-/-} ES cells into wild-type blastocysts. TfR1^{-/-} ES cells (Ned et al., 2003) were labeled by targeting the Rosa26 locus with GFP (Srinivas et al., 2001) and two different clones were utilized in subsequent experiments. The knock in of GFP was confirmed by Southern blotting, and the deletion of TfR1 was confirmed by a series of assays that included genotyping, measuring cell surface TfR1 protein with CD71 antibodies, and incubation with Alexa568-transferrin, which showed that GFP targeted TfR1^{-/-} ES cells neither expressed TfR1 nor captured transferrin (Figure S3).

The GFP-TfR1^{-/-} cells populated many embryonic organs including lung, liver, mid-gut, kidney, bladder, and gonad, and within these tissues, GFP-TfR1^{-/-} produced both epithelia (including branching and sheet structures) and mesenchymal precursors (Figure S4), demonstrating the multipotency of the GFP-TfR1^{-/-} cells.

In the kidney, we found extensive chimerism in the Wolffian ducts and the metanephric mesenchyme, as expected from our observation of early kidney structures in TfR1^{-/-} embryos at E11.5 (Figure S5). At the E15 stage, GFP-TfR1^{-/-} cells were also present throughout the kidney (Figure 2, Figure S5; n = 14 total foster mothers, using two clones of GFP-TfR1^{-/-} ES cells). To examine the contribution of the GFP-TfR1^{-/-} cells to individual compartments, we compared the percentage of GFP⁺ cells found in each compartment, to the relative size of the compartment, measured by planimetry. The GFP-TfR1^{-/-} cells populated distal tubules (identified by staining for E-cadherin; median ratio = 1.7; p = 0.0001), proximal tubules (which stained with lotus lectin; median ratio = 1.930; p = 0.0425) and the stroma and capsule, which contained as much as 42% \pm 17% of the GFP-TfR1^{-/-} cells (median ratio = 4.1; p < 0.001), even though these compartments account for only 6%–9% of the cells in the E15 kidney (as estimated by FACS analysis of kidneys expressing Foxd1-GFP). In contrast to these findings, cortical tips of the ureteric bud (highlighted by the expression of both E-cadherin and Troma1) contained the fewest GFP-TfR1^{-/-} cells (median ratio = 0.43; p = 0.003), even in kidneys with high overall degrees of chimerism (Figure 2, Figures S5 and S6). The exclusion of GFP-TfR1^{-/-} cells from the UB tip contrasted with chimeras generated with GFP-TfR1^{+/+} ES cells, which populated all compartments including UB tips (p = 0.78; Figure S7).

To confirm the anatomical distribution of GFP-TfR1^{-/-} cells, we used FACS to separate GFP-TfR1^{-/-} from wild-type cells using two different sets of kidneys (E15) with high degrees of




Figure 3. Ferritin Capture, Cell Proliferation, and Scara5 Expression in the Capsule of the E15 Kidney

(A and B) Both wild-type and GFP-TfR1^{-/-} capsular cells captured rhodamine coupled “Exogenous L-Ferritin.” Bars = 20 μ m.

(C and D) (C) Holo-ferritin (+HF; 1.25 μ g/ml) induced phosphohistone⁺ nuclei and expanded layers of capsular cells, whereas culture with apo-ferritin (-HF; 1.25 μ g/ml) failed to preserve the capsule (D).

(E) “Endogenous Ferritin” was present throughout the kidney, particularly in the capsule. E-cadherin marked the UB.

(F) Scara5 RNA was expressed by spindle-shaped cells encapsulating the kidney (arrowheads) and more faintly by cortical stroma.

detected by activated caspase-3 staining; $p = 0.26$). These findings imply that GFP-TfR1^{-/-} cells were not severely iron-starved. Yet, in situ, these cells failed to capture Alexa568-transferrin (Figure 2H), implying that they did not express

an alternative transferrin receptor. These data suggested that transferrin-independent mechanisms of iron capture were active in GFP-TfR1^{-/-} cells.

Endocytosis of Transferrin and Ferritin by Different Cellular Compartments in Developing Kidney

Extracellular ferritin is a non-transferrin iron carrier that is endocytosed by a variety of cells, including squamous-like cells of the adult kidney capsule (Kobayashi, 1978). To investigate ferritin capture by embryonic kidney, we incubated E15 kidneys with Alexa568-*apo*-ferritin, and found prominent endocytosis of ferritin by capsular cells. The capture of ferritin was independent of TfR1 because both wild-type and GFP-TfR1^{-/-} cells captured ferritin (Figures 3A and 3B). Ferritin L-chain was recognized by the cells, because the pattern of uptake was reproduced by L- but not by H-ferritin. To determine whether ferritin transferred its iron to the cell cytoplasm, we incubated explanted kidneys in the presence of either iron-loaded ferritin (holo-ferritin) or iron-depleted ferritin (*apo*-ferritin) and assayed proliferation in the capsular region. When holo-ferritin was administered to explanted kidneys, 49% of capsular nuclei expressed the proliferation marker phospho-histone H3, whereas *apo*-ferritin induced labeling in only 12% of nuclei ($p = 0.04$; Figures 3C and 3D), implying that ferritin-mediated iron delivery determined the cellularity of the renal capsule. Consistently, prominent staining for endogenous ferritin (Figure 3E) implied ongoing iron accumulation in the capsule in vivo.

Ferritin uptake was quite different from that of transferrin. TfR1 was largely restricted to the ureteric bud, cap mesenchyme, and primitive nephron (Figures 4A–4C), identical to sites where transferrin was captured (Figure 4D). In contrast, capsule and stroma demonstrated limited endocytosis of transferrin, even though they were capable of endocytosis of fluid phase fluorescent dextran (Figure 4E). These results demonstrate that ferritin and transferrin receptors are both present in developing kidney, but that they mediate iron transport to different cell types.

chimerism (~60%) and measured gene expression using Affymetrix arrays (Mouse Genome 430 2.0). In agreement with prior studies (Levy et al., 1999; Ned et al., 2003), GFP-TfR1^{-/-} cells were incompetent to participate in erythroid and granulocytic lineages: erythroid (*hemoglobin*, *aminolevulinic acid synthase 2*) and granulocytic (*lactoferrin*, *neutrophil granule protein*) genes were reduced on average 14-fold in GFP-TfR1^{-/-} compared to TfR1^{+/+} cells. Similarly, genes preferentially expressed by ureteric bud tips (*Wnt11*, *Wnt9B*, *Cytokine Receptor Like Factor1*, *Ros1*; Kanwar et al., 1995; Kispert et al., 1996; Marose et al., 2008; Schmidt-Ott et al., 2005) were decreased to 0.35 ± 0.27 fold (TfR1^{-/-} versus TfR1^{+/+}). On the contrary, genes expressed in renal stroma (*FGF-7*, *SFRP1*, *Semaphorin3A*, *Glypican3*, *FoxD1*, *Angiotensin II receptor, type 2*) and capsule (*Col6a1*, *Col5A2*, *Nfix*) were increased 2.9 ± 0.8 - and 2.7 ± 0.26 -fold, in GFP-TfR1^{-/-} versus TfR1^{+/+} cells, respectively (data not shown). These data were extended by analyzing a series of genes restricted to UB tip, stalk, or the mesenchyme-stroma compartment, that we identified by expression profiling (Schmidt-Ott et al., 2005). This comparison showed markedly reduced expression of the top 200 ureteric tip-specific genes in GFP-TfR1^{-/-} cells isolated from E15 chimeric kidneys, when compared with the TfR1^{+/+} population (0.42-fold; $p = 0.0000921$; Figure 2E). In contrast, the expression of the top 200 UB stalk-specific genes was only slightly reduced (0.71-fold; $p = 0.00265$), while the expression of the top 200 mesenchymal-stromal-specific genes was unchanged (1.2-fold; $p = 0.253$; Figure 2F). These data confirmed the preferential location of GFP-TfR1^{-/-} cells in mesenchyme and stroma, particularly compared to ureteric tip.

While it may be the case that the surviving GFP-TfR1^{-/-} cells were severely iron-deficient, this was unlikely because, as shown in Figure 2G, they displayed cytoplasmic iron stores (intracellular ferritin). In addition, GFP-TfR1^{-/-} cells demonstrated only slightly reduced proliferation (as detected by phospho-histone H3 staining median ratio = 0.69, $p = 0.0051$) compared to neighboring wild-type cells, and the same frequency of apoptosis (as




Figure 4. Localization of the Transferrin Pathway

(A–C) Tfr1 was initially localized to UB tips and cap mesenchyme. By E13, renal vesicles also expressed Tfr1 (*B, C). Tfr1 expression was reduced or absent in glomeruli (“G”), stroma, and capsule.

(D and E) (D) Rhodamine and (E) fluorescein transferrin endocytosis paralleled Tfr1 expression. Stroma and capsule were weakly labeled despite active fluid-phase endocytosis (rhodamine-dextran; [E]). (*Presumptive distal nephron; Bar = 10 μ m).

of 100-fold excess unlabelled ferritin blocked both binding and endocytosis (Figure 6B). Scara5 coimmunoprecipitated from clones D2 and H2 together with ferritin-avidin, but not with avidin itself (Figure 6C). Immunoprecipitation with ferritin-avidin coated biotin beads pulled down immunodetectable Scara5, but avidin-coated biotin beads did not (Figure 6D). These data indicate that Scara5 and ferritin interact at the cell surface.

Identification of a Novel Ferritin Receptor

The robust chimerism of GFP-Tfr1^{-/-} cells in the stroma and capsule suggested that they must capture non-transferrin iron, perhaps including ferritin. To examine this possibility, we isolated GFP-Tfr1^{-/-} cells by FACS and profiled their gene expression. We found that 156 genes were consistently upregulated ≥ 2 -fold in the GFP-Tfr1^{-/-} population compared with wild-type cells (data not shown), but only one of these, scavenger receptor, member 5 (Scara5) was markedly upregulated (7.1-fold in GFP-Tfr1^{-/-} versus Tfr1^{+/+}; $p = 0.0011$) and shown by in situ hybridization to be specifically located in a rim of cells where ferritin was captured in both embryonic (Figure 3F) and adult kidney (data not shown). Scara5 was also present in the airway, the developing aorta, and muscle bundles, and it was heavily expressed by gonadal epithelia (Figure S8; Jiang et al., 2006).

To directly test whether ferritin might be a ligand for Scara5, we expressed V5-tagged Scara5 in Trvb cells, a cell line that lacks endogenous expression Scara5 and Tfr1 (McGraw et al., 1987). Transiently transfected cells internalized apo-ferritin (Figure 5A), while Trvb cells transfected with a control vector did not (data not shown). Trvb clones (D2 and H2) that stably overexpressed Scara5 also demonstrated enhanced ferritin uptake compared to parental cells (Figure 5B) or clones (B2) that failed to express the protein. Western blots of these clones showed that Scara5 was present as 59 kDa and 70 kDa proteins (Figure 5C), suggesting posttranslational processing.

To demonstrate that Scara5 binds ferritin at the cell surface, we added native ferritin at 4°C to MSC-1 cells overexpressing V5-tagged Scara5. Using two-color immunofluorescence, we found immunodetectable patches of V5 and ferritin on the surface of nonpermeabilized cells (Figure 6A). Raising the temperature to 37°C resulted in endocytosis of the ferritin, while inclusion

of 100-fold excess unlabelled ferritin blocked both binding and endocytosis (Figure 6B). Scara5 coimmunoprecipitated from clones D2 and H2 together with ferritin-avidin, but not with avidin itself (Figure 6C). Immunoprecipitation with ferritin-avidin coated biotin beads pulled down immunodetectable Scara5, but avidin-coated biotin beads did not (Figure 6D). These data indicate that Scara5 and ferritin interact at the cell surface. It is well known that scavenger receptors bind more than one ligand, however to our knowledge iron-transporting proteins had not been shown to bind to this class of receptors. Using our stable Scara5-Trvb clones, we studied whether Scara5 recognized a variety of iron carriers. L-ferritin was the preferred ligand; addition of as little as 250 pmole of L-chain (equivalent to 10 pmoles of an L-chain complex) in the media led to detection of ferritin in cellular vesicles of H2 and D2 clones but not in parental Trvb cells nor in nonexpressing clone B2 (Figure 5D). Scavenger receptors bind to their ligands using electrostatic interaction. Because attachment of fluorescent probes to ferritin might have altered its net charge, we tested ferritin attached to two fluorophores, Alexa-568 which adds a negative charge and Alexa488 and Rhodamine which adds a positive charge. The uptake of these two types of ferritins resulted in similar levels of Scara5-dependent endocytosis. Haptoglobin-hemoglobin was also internalized by H2 and D2 clones, but the signal was less distinct than that of L-ferritin (Figure S9). H-ferritin, in contrast, was not captured, nor were transferrin, albumin, hemopexin-heme, or lipocalin2:catechololate siderophore. Further, Scara5 differed from other scavenger receptors since it did not capture fluorescent acetylated LDL (Jiang et al., 2006). These data suggest that among other potential ligands, Scara5 is a receptor or a component of a receptor for L-ferritin.

Scara5 Expressing Cells Can Obtain Iron from Ferritin

To examine whether Scara5 can deliver iron to the cell, we incubated Scara5 expressing D2 and H2 clones with ferritin loaded with radioactive iron and found that there was a time-dependent incorporation into the Scara5-expressing cells, but not into the parental Trvb cell (Figure 7A). Iron was likely obtained after endocytosis into an acidic compartment, because 4°C or inhibition of vacuolar H⁺ATPase with bafilomycin blocked iron capture and




Figure 5. Interaction between Scara5 and L-Ferritin

(A and B) (A) Transient transfection of Scara5-V5 (anti-V5, green) and (B) stable expression of Scara5-V5 resulted in the capture of Alexa568-ferritin ([A and B], right panel), whereas parental Trvb, which are Scara5⁻Tfr1⁻ cells, transfected with an empty vector were negative ([B] left panel). Tris-quenched Alexa568 dye (no ferritin) was an additional control (middle panel).

(C) Expression of Scara5-V5 (59 KDa, 70 KDa) was detected by V5 monoclonal antibody.

(D) Endocytosis of Alexa568-ferritin was analyzed in Scara5⁻ (Trvb, B2 clones) and Scara5⁺ (D2 and H2) stable clones. Nearly all D2 and H2 cells captured apo- and L-ferritins, but not H-ferritin. Bar = 10 μ m.

ferritin traffic to lysosomes (Figures 7A and 7B). Consistent data came from a MSC-1 reporter cell line which expressed endogenous Scara5 as well as fluorescent proteins driven by iron responsive elements (IRE, Li et al., 2004). Holo-ferritin upregulated CFP encoded by a 5'IRE reporter gene, and downregulated YFP encoded by a 3'IRE reporter gene, consistent with the transfer of

iron to the cytoplasm (Figure 7C). In the presence of holo-ferritin (2 μ g/ml), the 5'IRE/3'IRE fluorescence ratio (1.58 ± 0.31) was not significantly different (ratio = 3.8 ± 1.5 ; $p = 0.19$) from that of cells incubated in media containing ferric ammonium citrate (50 μ M) and holo-transferrin (20 μ g/ml). Cells incubated in media containing the iron chelator desferrioxamine (20 μ M) had




Figure 6. Interaction between Scara5 and Ferritin at the Cell Surface

(A) Scara5-V5 (anti-V5; Cy2; green) and ferritin (anti-ferritin; Cy3; red) colocalized at the surface (Z stack; inset: single confocal section) of Scara5-V5 transfected MSC-1 cells incubated with untagged ferritin (2 hr; 4°C). Toto3 marked nuclei. Bar = 10 μ m.

(B) Endocytosis of rhodamine-ferritin (30 hr; 37°C; top panel) was blocked by a 100-fold excess untagged ferritin (bottom panel). Bar = 20 μ m.

(C) V5 antibodies precipitated Scara5-V5 with avidin tetramer-ferritin monomer (85 kDa) or multimers (200 kDa > 250 kDa), but not with unconjugated avidin. Irrelevant isotype matched antibodies did not immunoprecipitate ferritin-avidin nor were ferritin-avidin complexes recovered without cell extracts.

(D) Scara5-V5 was detected in cell extracts (clone D2) and in pull-downs with ferritin-avidin-biotin-agarose beads, but not with avidin-biotin-agarose-beads.

a significantly lower ratio (0.53 ± 0.09 ; $p = 0.044$) than cells incubated with holo-ferritin. To test whether ferritin iron can enhance cell survival under iron-limiting conditions in a Scara5-dependent manner, we incubated clones (D2 or H2) in minimal iron-deficient media supplemented with either holo- or apo-ferritin. Whereas Trvb cells were unresponsive to either protein ($n = 8$), holo-ferritin, but not apo-ferritin, dose-dependently supported the survival of Scara5 expressing cells as measured by GAPDH blots (Figure 7D). Together, these data show that ferritin can deliver iron and regulate iron-dependent intracellular events in a Scara5-dependent manner.

Scara5 Is an Endogenous Ferritin Uptake Pathway

To determine whether native Scara5 acts as a ferritin receptor, we suppressed its message in MSC-1 cells by RNA interference. We derived stable clones expressing short hairpin RNA (shRNA) targeting Scara5 and then transfected these clones with Scara5 siRNA (Qiu et al., 2006). This protocol reduced Scara5 message to $29 \pm 3.0\%$. Control studies showed that LacZ shRNA plus LacZ siRNA did not restrict the expression of Scara5 ($110.5 \pm 0.26\%$ compared to nontransfected MSC-1 cells). The interference with Scara5 expression inhibited fluorescein-ferritin endo-

cytosis, measured by FACS ($61 \pm 0.1\%$ compared to LacZ targeted cells; $n = 5$; $p = 0.00093$).

Chemical blockade of Scara5 with polyanions further confirmed the interaction with ferritin. Polyinosinic acid (PI; 50 μ g/ml) nearly abolished the capture of Alexa488-ferritin by MSC-1 cells, whereas polyguanylic acid (PG) and polycytidylic acid (PC) were less effective (Figure 7E) as previously demonstrated for other ligands by Jiang et al. (2006).

Consistently, the transfer of iron from holo-ferritin to MSC-1 cells was inhibited by PI, suppressing the ratio of iron reporters 5'IRE-CFP/3'IRE-YFP (0.7 ± 0.06 ; holo-ferritin versus holo-ferritin+PI, $p = 0.05$) whereas PC was ineffective (1.2 ± 0.2 ; holo-ferritin versus holo-ferritin+PC, $p = 0.4$). In contrast, PI did not block the response of the iron reporters to ferric ammonium citrate (data not shown). To demonstrate that this behavior was not unique to the MSC-1 cells, we repeated the inhibition studies in clones D2 and H2 and in explanted kidneys and found similar results. In aggregate, our data demonstrate that Scara5 can act as a ferritin receptor delivering iron to the cytoplasm where it activates iron responsive elements.

DISCUSSION

While iron transport to developing organs is critical, the molecules that mediate this process remain poorly understood. Genes that deliver iron to some lineages do not seem to be needed throughout the embryo. For example, TfR1^{-/-} cells can populate developing organs (Ned et al., 2003), DMT1^{-/-} mice still undergo hepatic organogenesis (Gunshin et al., 2005), and mutations in Steap3 (Ohgami et al., 2005) are compatible with development. The candidate iron carrier Lcn2 is not essential in development




Figure 7. Scara5 Mediates Cellular Uptake of Ferritin-Bound Iron and Represents an Endogenous Pathway in MSC-1 Cells

(A) Capture of ^{59}Fe -ferritin by Scara5 $^{+}$ at 37°C (red line) but not at 4°C (black line) nor by parental Trvb cells (blue line). Error bars indicate mean \pm SD.
 (B) Lysosomes of Scara5 $^{+}$ Trvb cells were marked with fluorescein-dextran (green). They contained endocytosed Alexa568-ferritin (red). Original magnification: 40 \times .
 (C) Transfer of iron to the cytoplasm was detected using two IRE-based iron reporters and FACS analysis in MSC-1 cells. The maximal 5'IRE-CFP signal (designated 100%-black bars) was generated by iron loading with holotransferrin (20 $\mu\text{g}/\text{ml}$) and ferric ammonium citrate (20 μM) (HT+Fe) and the maximal 3'IRE-YFP signal (designated 100%-stippled bars) was generated by DFO (20 μM). These treatments produced a \sim 7-fold range in the ratio of 5'/3'IRE signals. Treatment with holoferritin (HF) increased 5'IRE-CFP but decreased the 3'IRE-YFP signal. Error bars indicate mean \pm SD.
 (D) Holoferritin induces cell growth. Scara5 $^{+}$ clone H responded to holo- but not to apo-ferritin, whereas Trvb cells failed to respond. Cell protein was detected by GAPDH immunoblots.
 (E) Alexa488-ferritin endocytosis was blocked by Poly I > Poly G > Poly C (50 μM). Similar data were obtained in Scara5 $^{+}$ Trvb and MSC1 cells. Bar = 20 μM .

(Flo et al., 2004) and was even downregulated in the GFP-TfR1 $^{-/-}$ cell population, suggesting that it cannot compensate for TfR1 $^{-/-}$.

To determine the mechanism of iron delivery in the embryonic kidney, we first examined the transferrin pathway. The most striking finding was a restricted requirement for TfR1. For example, thousands of differentiated kidney, gonad, bladder, mid-gut, liver, and lung cells were generated from only 3–5 GFP-TfR1 $^{-/-}$ ES cells, implying that expression of TfR1 was not required at any step of their lineage. In fact, we failed to visualize an interaction of transferrin with GFP-TfR1 $^{-/-}$ cells (Thorstensen et al., 1995), suggesting that they did not express an uncharacterized transferrin receptor, although chimeric embryos in the *hpx* background must be generated to formally test this idea. It also remained possible that iron was supplied from immediately adjacent TfR1 $^{+/+}$ mesenchyme or epithelia, but this idea was effectively ruled out by the large clusters of differentiated GFP-TfR1 $^{-/-}$ cells encompassing entire kidney compartments (Figures 2A, 2D, and 2H; Figure S5D, R, BB). In sum, we found

that TfR1 did not limit the development of mesenchyme, nephron, stroma, or capsule cells, most likely because of transferrin-independent pathways. In fact, when we introduced ^{55}Fe during pregnancy (E11.5 or E12), ^{55}Fe was uniformly distributed at E12.5 (Figure S10) despite variations in the expression of TfR1, again implicating one or more non-transferrin mechanisms of iron delivery in the kidney.

Extracellular ferritin may be a non-transferrin source of iron. First, while serum ferritin is not thought to be a major iron carrier in the adult, its concentration is nearly 3–8-fold higher in fetal blood than that of the mother (approximately 0.3 $\mu\text{g}/\text{ml}$; Puolakka et al., 1980; Siddappa et al., 2007) and it has the potential to carry a larger number of iron atoms/molecule than transferrin (Watanabe et al., 2001; Levi et al., 1989, 1992). Second, extracellular ferritin may be synthesized locally by macrophages (Leimberg et al., 2008; Yuan et al., 2004), which are abundant in metanephric mesenchyme (Rae et al., 2007). During kidney development, 25% of mesenchymal cells undergo apoptosis, and 3% of kidney cells

are apoptotic at steady state (Koseki et al., 1992). While apoptotic cells can be cleared by a number of cell types, there is evidence for macrophage involvement in the kidney (Camp and Martin, 1996). Hence, ferritin may derive from serum, but may also be obtained locally from macrophages, recycling iron from apoptotic cells.

The capture of ferritin was originally demonstrated in hepatocytes (Mack et al., 1983; Sibille et al., 1989), intestine (San Martin et al., 2008), white matter (Hulet et al., 1999), lymphoid cells (Moss et al., 1992), erythroid precursors (Meyron-Holtz et al., 1999), and in the capsule of the adult kidney (Kobayashi, 1978). In all of these cell types, ferritin used an endocytic pathway (Hulet et al., 2000; Gelvan et al., 1996; Bretscher and Thomson, 1983) and delivered iron to the cytoplasm (Radisky and Kaplan, 1998; San Martin et al., 2008). Our studies have added stroma of embryonic capsule to this long list of cells.

Scavenger receptors bind their ligands using a series of positively charged residues in the collagenous (Doi et al., 1993) or cysteine-rich domains (Ojala et al., 2007). Using these residues they bind lipopolysaccharides, bacteria, and polynucleotides, all of which share repetitive polyanionic motifs (Jiang et al., 2006). While iron carriers have not been studied as ligands for scavenger receptors, ferritin contains repeating chains of polyanionic monomers, and like classical polyanionic ligands, its binding was sensitive to polynucleotides. In this light, it is important to determine whether ferritin binds to more than one scavenger receptor, or is uniquely bound by Scara5, particularly given that Scara5 appears not to bind the typical ligands, such as acetylated or oxidized LDL. Moreover, Scara3 (Han et al., 1998) which was also expressed in renal stroma and capsule, did not capture ferritin (data not shown). Hence, while we present data demonstrating a Scara5-ferritin interaction, it is unresolved whether Scara5 is a unique ferritin receptor component, or rather simply an example of a common scavenger-ferritin interaction.

A number of functions may be served by a Scara5-ferritin interaction, in addition to supplying iron to the cell hosting the receptor. These include transporting iron across a cellular layer. The best studied model of iron transcytosis is the duodenal cell, which provides ferroportin-dependent iron export. In an analogous fashion, brain capillary cells (Fisher et al., 2007; Xu and Ling, 1994) and placenta can also transfer iron in fact from ferritin through ferroportin (Lamparelli et al., 1989; Bastin et al., 2006). The renal capsule includes c-kit⁺flk⁺podocalyxin⁺ nascent capillaries (Schmidt-Ott et al., 2006) which may enhance delivery of serum ferritin, while their expression of ferroportin may export iron to adjacent stromal and mesenchymal cells. Perhaps transcellular transport provides iron to developing organs at critical stages prior to the completion of the vascular tree. While such a mechanism remains speculative, it is supported by the coexpression of Scara5 and ferroportin in mesenchyme that surrounds proximal lung buds and by the expression of Scara5 and ferroportin in thin layers of cells that surround other developing organs.

In contrast to the widespread development of GFP-TfR1^{-/-} stroma, mesenchyme, and epithelia, we found limited development of GFP-TfR1^{-/-} in ureteric tips, suggesting their dependence on transferrin. Similarly, while the lung was heavily populated by GFP-TfR1^{-/-} cells, the tips of the airway had the fewest GFP-TfR1^{-/-} cells compared with stalks or mesenchyme (Figure S4). Because the expression of TfR1 has historically been linked to

stages of cell proliferation (Kanayasu-Toyoda et al., 1999; Sposi et al., 2000; Wingert et al., 2004), it is tempting to propose that the high proliferative rate of tip cells in kidney and in lung (Michael and Davies, 2004; Nogawa et al., 1998; Mollard and Dziadek, 1998) demands transferrin. While this model is compelling because TfR1 is prominent in branching epithelia, this hypothesis requires testing by measuring the relative proliferative rates of different cell types at different stages in vivo and direct correlation with the level of TfR1 expression.

In sum, we have utilized a deletion of TfR1 to study iron transport in embryonic kidney. This analysis was possible because the second transferrin receptor (TfR2) had limited or no expression in the kidney as demonstrated by our gene arrays and by *in situ* hybridization (data not shown). The data indicate that Tf-TfR1 targets specific compartments rather than every cell type. Transferrin is captured and required by the ureteric bud tip after E11, but it is not required by metanephric mesenchyme and is poorly internalized and not required by stromal and capsular cells. We identified a novel ferritin receptor and suggest that this pathway may support growth of kidney cells, not only in capsule, but potentially in adjacent mesenchymal cells as well, hence bypassing an overt requirement for TfR1.

EXPERIMENTAL PROCEDURES

TfR1 Knockout Embryos

TfR1^{+/-} mice (Levy et al., 1999) were bred with C57BL/6J for five generations and the alleles detected using radio- or ⁵Fam-labeled primers (Invitrogen) as described (Levy et al., 1999).

GFP-TfR^{-/-} ES Cells and Chimeric Embryos

The ROSA26-eGFP knockin construct (Srinivas et al., 2001) was reconstructed with a puromycin selectable marker and introduced into TfR1^{-/-} ES clones (H5, E12; Ned et al., 2003). One hundred and ninety-two puromycin-resistant (250 μg/ml) colonies were screened by Southern blot, by anti-CD71 (BD Pharmingen) and by Alexa-568-holo-transferrin (20 μg/ml, 1 hr) endocytosis.

Chimeras were generated by introducing 3–5 GFP-TfR1^{-/-} or TfR1^{+/-} ES cells into wild-type C57BL/6 blastocysts. Chimeric tissues were analyzed in permeabilized cryosections (0.1% Triton X-100/0.2% BSA in PBS for 15 min at RT) with antibodies to pan Cytokeratin (1:300; Sigma), Pax-2 (1:100; Zymed), Podocalyxin (1:300; R&D System), and Phospho-Histone3 (1:50; Cell Signaling Technology). Alternatively, permeabilized cryosections (0.1% Triton X-100 and 0.5% Saponin in PBS for 1 hr with 1% heat inactivated horse serum at RT) were immunoassayed with E-Cadherin (1:400; R&D Systems), Tenascin (1:400; Chemicon), TROMA-1 (1:20; Developmental Studies Hybridoma Bank), active Caspase-3 (1:250; Promega), and ferritin (1:250; Biogenesis) antibodies.

Microarray Analysis of GFP-TfR1^{-/-} Kidney Cells

Chimeric E15 kidneys from two independent litters were digested with collagenase (30 min) and trypsin (5 min), quenched with DMEM with 10% FCS, and 10⁶ GFP-TfR1^{-/-} or TfR1^{+/-} cells collected by FACS. Total RNA (50 ng) was biotin labeled using the Ovation Biotin System (NuGen) and cDNA (2.2 μg) was hybridized in duplicate to Mouse Genome 430 2.0 Genechip arrays (Affymetrix). Image files were analyzed by robust multichip analysis.

Real-time PCR amplification of Scara5 utilized forward (5'tggggctgtatcttc tggctc3') and reverse (5'atctctgaacctccacacct3') primers and iQ SYBR® Green Super Mix (Biorad), cycled 95°C/2 min, and 95°C/30 s and 60°C/30 s × 50 cycles. Specificity was checked by melting curve analysis and electrophoresis. Scara5 expression was normalized to β-actin expression (forward 5'ctaagg ccaaccgtgaaaag3' and reverse 5'tctcagctgtggtggaag3' primers) by ΔC_T method.

Generation of CMV-Scara5-V5 Expressing Cell Lines

Full-length scara5 cDNA was cloned from mouse E15 Clontech cDNA library by using a forward 5'caccatggacaacaagc3' and reverse 5' ggggacagtagca gtcac3' primers and pcDNA3.1 Directional TOPO expression plasmid. Scara5-V5 was then amplified with primers 5'gcggtatccatggacaacaagccat gtagc-3' that included a BamH1 site, and 5'gcggttaaaactcagtagaattcgagacc gag-3' that included a PmeI site followed by ligation to pcDNA3.1*.

Trvb cells (a kind gift of McGraw et al., 1987) and MSC-1 cells (a kind gift of McGuinness et al., 1994) were transfected with Scara5, selected with G418 (400 µg/ml) and authenticated by anti-V5 immunoblots (1:5,000 Invitrogen) or immunofluorescence (1:2,000). Ferritin mediated growth in serum free culture media was measured in 2×10^4 cells/well (60 hr) using GAPDH immunoblots (Abcam, 1:5,000).

Immunoprecipitation of Ferritin:Scara5 Complexes

Scara5⁺-Trvb clones were lysed (in 0.5% NP40, DNase I with protease inhibitors), sonicated (10 s, 4°C), sheared with an insulin syringe 3× and centrifuged (14,000 rpm; 20 min at 4°C). The supernatant was diluted 2-fold, filtered (0.45 µm) and avidin-ferritin (20 µg; Sigma) or equimolar avidin (2.5 µg; Sigma) added (3 hr). Immunoprecipitates were collected with Protein A and G Agarose Beads (10 µl each; Boehringer-Mannheim) coated with V5 or IgG2a isotype-matched antibodies (10 µg, Jackson ImmunoResearch) for 3 hr and washed in PBS with Ca, Mg, 0.5% NP40 (5 times). Avidin-ferritin was detected with biotin-HRP (1 µg/ml in 1% BSA in TBST, Jackson ImmunoResearch). The reciprocal immunoprecipitation was performed by coating Biotin-Agarose (10 µl) beads with Avidin or Avidin-ferritin and detecting Scara5 by anti-V5 blots.

Binding and Endocytosis of Ferritin

5,6 carboxytetramethyl-rhodamine, Alexa568, and Alexa488 succinimidyl ester (Molecular Probes) coupled apo-ferritin were authenticated by SDS-PAGE after clean-up on a 100 kDa cut-off microcon. Cells were incubated (37°C; 30 hr) with fluorescent ferritin (2 µg/ml) and polynucleotides (50 µg/ml) or untagged ferritin (100-fold excess) and fluorescence quantified by BD LSRiI flow cytometer and FloJo7.2.2 software. In some cases, cells were colabeled with fluorescein-dextran (1 mg/ml; 3 hr) followed by overnight washout. Surface binding utilized apo-ferritin (100 µg/ml; 4°C; 2hr) followed by immunodetection in unpermeabilized cells with anti-ferritin (Biogenesis, 1:200) and anti-V5. ⁵⁹Fe-ferritin was prepared by incubating apoferritin at a 1:10 ratio with ⁵⁹Fe in 10 mM HCO₃ at 37°C overnight, followed by clean-up on a 100 kDa cut-off microcon. ⁵⁹Fe-ferritin (3 × 10⁴cpm) was added to 24-well plates containing 2×10^4 cells.

Iron reporter probes were introduced into MSC1 cells by adenoviral mediated gene transfer. E1/E3 deleted type 5 adenoviral expression constructs contained either a 5'IRE upstream of destabilized ECFP or a 3'IRE complex downstream of destabilized EYFP. Virus was produced in 293A cells (ViraPower; Invitrogen) and purified on a CsCl step-gradient, followed by dialysis and infected with MOI = 10.

Knockdown of Scara5 in MSC-1 Cells

U6 pEntry Vector (Invitrogen) was used to express hairpin RNAs encoding two shRNA sequences unique to Scara5 (nucleotides 434–452 and 1222–1240; GenBank Accession Number: BC016096) from annealed ssDNAs (1 µg/µl; Invitrogen). MSC-1 were transfected with pU6SR5 or pU6LacZ control plasmid, selected with Blastidicin (2 µg/ml), and three anti-Scara5 siRNA oligos (Ambion) were then transfected to increase the interference. Knockdown was quantified by real-time PCR, incubation with FITC-ferritin (20 µg/ml; 2 hr at 37°C), and analysis by FACS.

In Situ Hybridization of Transferrin Receptor1 and Scara5

Digoxigenin-labeled (Roche) antisense riboprobes were generated from plasmids encoding TfR1 (1000 nt, exon 1–7) and Scara5 (1476 nt, exon 1–8). Sections were stained with anti-digoxigenin (Boehringer-Mannheim; 1:5000 dilution; Schmidt-Ott et al., 2005).

ACCESSION NUMBERS

Microarray data have been deposited in GEO DataSets with accession code GSE7396.

SUPPLEMENTAL DATA

Supplemental Data include ten figures and can be found at [http://www.cell.com/developmental-cell/supplemental/S1534-5807\(08\)00512-1/](http://www.cell.com/developmental-cell/supplemental/S1534-5807(08)00512-1/).

ACKNOWLEDGMENTS

We thank F.Costantini, C. Mendelsohn, and Q. Al-Awqati for key reagents and advice. We thank the reviewers for very insightful comments. This work was supported by grants from the Emerald Foundation, the March of Dimes, and the NIDDK (Grants DK-55388 and DK-58872) to J.B. and the NIHLB (R01-HL51057) to N.C. Andrews. K.M.S. is an Emmy Noether Fellow of the Deutsche Forschungsgemeinschaft, Germany. There are no conflicts of interest.

Received: June 27, 2008

Revised: October 13, 2008

Accepted: December 17, 2008

Published: January 19, 2009

REFERENCES

- Aisen, P. (2004). Transferrin receptor 1. *Int. J. Biochem. Cell Biol.* 36, 2137–2143.
- Arosio, P., and Levi, S. (2002). Ferritin, iron homeostasis, and oxidative damage. *Free Radic. Biol. Med.* 33, 457–463.
- Bastin, J., Drakesmith, H., Rees, M., Sargent, I., and Townsend, A. (2006). Localisation of proteins of iron metabolism in the human placenta and liver. *Br. J. Haematol.* 134, 532–543.
- Beaumont, C., Leneuve, P., Devaux, I., Scoazec, J.Y., Berthier, M., Loiseau, M.N., Grandchamp, B., and Bonneau, D. (1995). Mutation in the iron responsive element of the L ferritin mRNA in a family with dominant hyperferritinaemia and cataract. *Nat. Genet.* 11, 444–446.
- Bratlid, D., and Moe, P.J. (1980). Hemoglobin and serum ferritin levels in mothers and infants at birth. *Eur. J. Pediatr.* 134, 125–127.
- Brekelmans, P., van Soest, P., Llenen, P.J., and van Ewijk, W. (1994). Inhibition of proliferation and differentiation during early T cell development by anti-transferrin receptor antibody. *Eur. J. Immunol.* 24, 2896–2902.
- Bretscher, M.S., and Thomson, J.N. (1983). Distribution of ferritin receptors and coated pits on giant HeLa cells. *EMBO J.* 2, 599–603.
- Camp, V., and Martin, P. (1996). The role of macrophages in clearing programmed cell death in the developing kidney. *Anat. Embryol. (Berl.)* 194, 341–348.
- Chen, T.T., Li, L., Chung, D.H., Allen, C.D., Torti, S.V., Torti, F.M., Cyster, J.G., Chen, C.Y., Brodsky, F.M., Niemi, E.C., et al. (2005). TIM-2 is expressed on B cells and in liver and kidney and is a receptor for H-ferritin endocytosis. *J. Exp. Med.* 202, 955–965.
- Dickinson, T.K., and Connor, J.R. (1995). Cellular distribution of iron, transferrin, and ferritin in the hypotransferrinemic (Hp) mouse brain. *J. Comp. Neurol.* 355, 67–80.
- Doi, T., Higashino, K., Kurihara, Y., Wada, Y., Miyazaki, T., Nakamura, H., Uesugi, S., Imanishi, T., Kawabe, Y., Itakura, H., et al. (1993). Charged collagen structure mediates the recognition of negatively charged macromolecules by macrophage scavenger receptors. *J. Biol. Chem.* 268, 2126–2133.
- Dressler, G.R. (2006). The cellular basis of kidney development. *Annu. Rev. Cell Dev. Biol.* 22, 509–529.
- Eklom, P., Thesleff, I., Saxen, L., Miettinen, A., and Timpl, R. (1983). Transferrin as a fetal growth factor: acquisition of responsiveness related to embryonic induction. *Proc. Natl. Acad. Sci. USA* 80, 2651–2655.
- Ferreira, C., Santambrogio, P., Martin, M.E., Andrieu, V., Feldmann, G., Henin, D., and Beaumont, C. (2001). H ferritin knockout mice: a model of hyperferritinaemia in the absence of iron overload. *Blood* 98, 525–532.
- Fisher, J., Devraj, K., Ingram, J., Slagle-Webb, B., Madhankumar, A.B., Liu, X., Klingler, M., Simpson, I.A., and Connor, J.R. (2007). Ferritin: a novel mechanism for delivery of iron to the brain and other organs. *Am. J. Physiol. Cell Physiol.* 293, C641–C649.

- Flo, T.H., Smith, K.D., Sato, S., Rodriguez, D.J., Holmes, M.A., Strong, R.K., Akira, S., and Aderem, A. (2004). Lipocalin 2 mediates an innate immune response to bacterial infection by sequestering iron. *Nature* 432, 917–921.
- Gelvan, D., Fibach, E., Meyron-Holtz, E.G., and Konijn, A.M. (1996). Ferritin uptake by human erythroid precursors is a regulated iron uptake pathway. *Blood* 88, 3200–3207.
- Genilli, C., Doliana, R., Bet, P., Campanile, G., Colombatti, A., Cancedda, F.D., and Cancedda, R. (1994). Ovotransferrin and ovotransferrin receptor expression during chondrogenesis an endochondral bone formation in developing chick embryo. *J. Cell Biol.* 124, 579–588.
- Ghosh, S., Hevi, S., and Chuck, S.L. (2004). Regulated secretion of glycosylated human ferritin from hepatocytes. *Blood* 103, 2369–2376.
- Gunshin, H., Fujiwara, Y., Custodio, A.O., Drenzo, C., Robine, S., and Andrews, N.C. (2005). Slc11a2 is required for intestinal iron absorption and erythropoiesis but dispensable in placenta and liver. *J. Clin. Invest.* 115, 1258–1266.
- Gustine, D.L., and Zimmerman, E. (1973). Developmental changes in microheterogeneity of foetal plasma glycoproteins of mice. *Biochem. J.* 132, 541–551.
- Hamill, R.L., Woods, J.C., and Cook, B.A. (1991). Congenital atransferrinemia. A case report and review of the literature. *Am. J. Clin. Pathol.* 96, 215–218.
- Han, H.J., Tokino, T., and Nakamura, Y. (1998). CSR, a scavenger receptor-like protein with a protective role against cellular damage caused by UV irradiation and oxidative stress. *Hum. Mol. Genet.* 7, 1039–1046.
- Hatini, V., Huh, S.O., Herzlinger, D., Soares, V.C., and Lai, E. (1996). Essential role of stromal mesenchyme in kidney morphogenesis revealed by targeted disruption of Winged Helix transcription factor BF-2. *Genes Dev.* 10, 1467–1478.
- Hayashi, A., Wada, Y., Suzuki, T., and Shimizu, A. (1993). Studies on familial hypotransferrinemia: unique clinical course and molecular pathology. *Am. J. Hum. Genet.* 53, 201–213.
- Huggenvik, J.J., Craven, C.M., Idzerda, R.L., Bernstein, S., Kaplan, J., and McKnight, G.S. (1989). A splicing defect in the mouse transferrin gene leads to congenital atransferrinemia. *Blood* 74, 482–486.
- Hulet, S.W., Hess, E.J., Debinski, W., Arosio, P., Bruce, K., Powers, S., and Connor, J. (1999). Characterization and distribution of ferritin binding sites in the adult mouse brain. *J. Neurochem.* 72, 868–874.
- Hulet, S.W., Heyliger, S.O., Powers, S., and Connor, J.R. (2000). Oligodendrocyte progenitor cells internalize ferritin via clathrin-dependent receptor mediated endocytosis. *J. Neurosci. Res.* 61, 52–60.
- Hussain, M.A., Gaafar, T.H., Laulicht, M., and Hoffbrand, A.V. (1977). Relation of maternal and cord blood serum ferritin. *Arch. Dis. Child.* 52, 782–784.
- Jiang, Y., Oliver, P., Davies, K.E., and Platt, N. (2006). Identification and characterization of murine SCARA5, a novel class A scavenger receptor that is expressed by populations of epithelial cells. *J. Biol. Chem.* 281, 11834–11845.
- Kanayasu-Toyoda, T., Yamaguchi, T., Uchida, E., and Hayakawa, T. (1999). Commitment of neutrophilic differentiation and proliferation of HL-60 cells coincides with expression of transferrin receptor. *J. Biol. Chem.* 274, 25471–25480.
- Kanwar, Y.S., Liu, Z.Z., Kumar, A., Wada, J., and Carone, F.A. (1995). Cloning of mouse c-ras renal cDNA, its role in development and relationship to extracellular matrix glycoproteins. *Kidney Int.* 48, 1646–1659.
- Kispert, A., Vainio, S., Shen, L., Rowitch, D.H., and McMahon, A.P. (1996). Proteoglycans are required for maintenance of Wnt-11 expression in the ureter tips. *Development* 122, 3627–3637.
- Kobayashi, K. (1978). Fine structure of the mammalian renal capsule: the atypical smooth muscle cell and its functional meaning. *Cell Tissue Res.* 195, 381–394.
- Koseki, C., Herzlinger, D., and al-Awqati, Q. (1992). Apoptosis in metanephric development. *J. Cell Biol.* 119, 1327–1333.
- Lamparelli, R.D., Friedman, B.M., MacPhail, A.P., Bothwell, T.H., Phillips, J.I., and Baynes, R.D. (1989). The fate of intravenously injected tissue ferritin in pregnant guinea-pigs. *Br. J. Haematol.* 72, 100–105.
- Leimberg, M.J., Prus, E., Konijn, A.M., and Fibach, E. (2008). Macrophages function as a ferritin iron source for cultured human erythroid precursors. *J. Cell. Biochem.* 103, 1211–1218.
- Levinson, R., and Mendelsohn, C. (2003). Stromal progenitors are important for patterning epithelial and mesenchymal cell types in the embryonic kidney. *Semin. Cell Dev. Biol.* 14, 225–231.
- Levi, S., Luzzago, A., Cesareni, G., Cozzi, A., Franceschinelli, F., Albertini, A., and Arosio, P. (1988). Mechanism of ferritin iron uptake: activity of the H-chain and deletion mapping of the ferro-oxidase site. *J. Biol. Chem.* 263, 18086–18092.
- Levi, S., Salfeld, J., Franceschinelli, F., Cozzi, A., Dorner, M.H., and Arosio, P. (1989). Expression and structural and functional properties of human ferritin L-chain from *Escherichia coli*. *Biochemistry* 28, 5179–5184.
- Levi, S., Yewdall, S.J., Harrison, P.M., Santambrogio, P., Cozzi, A., Rovida, E., Albertini, A., and Arosio, P. (1992). Evidence of H- and L-chains have co-operative roles in the iron-uptake mechanism of human ferritin. *Biochem. J.* 288, 591–596.
- Levi, S., Santambrogio, P., Cozzi, A., Rovida, E., Corsi, B., Tamborini, E., Spada, S., Albertini, A., and Arosio, P. (1994). The role of the L-chain in ferritin iron incorporation. Studies of homo and heteropolymers. *J. Mol. Biol.* 238, 649–654.
- Levy, J.E., Jin, O., Fujiwara, Y., Kuo, F., and Andrews, N.C. (1999). Transferrin receptor is necessary for development of erythrocytes and the nervous system. *Nat. Genet.* 21, 396–399.
- Li, J.Y., Ram, G., Gast, K., Chen, X., Barasch, K., Mori, K., Schmidt-Ott, K., Wang, J., Kuo, H.C., Savage-Dunn, C., et al. (2004). Detection of intracellular iron by its regulatory effect. *Am. J. Physiol. Cell Physiol.* 287, C1547–C1559.
- Linder, M.C., Schaffer, K.J., Hazegh-Azam, M., Zhou, C.Y., Tran, T.N., and Nagel, G.M. (1996). Serum ferritin: does it differ from tissue ferritin? *J. Gastroenterol. Hepatol.* 11, 1033–1036.
- Macedo, M.F., de Sousa, M., Ned, R.M., Mascarenhas, C., Andrews, N.C., and Correia-Neves, M. (2004). Transferrin is required for early T-cell differentiation. *Immunology* 112, 543–549.
- Mack, U., Powell, L.W., and Halliday, J.W. (1983). Detection and isolation of a hepatic membrane receptor for ferritin. *J. Biol. Chem.* 258, 4672–4675.
- MacPhail, A.P., Charlton, R.W., Bothwell, T.H., and Torrance, J.D. (1980). The relationship between maternal and infant iron status. *Scand. J. Haematol.* 25, 141–150.
- Marose, T.D., Merkel, C.E., McMahon, A.P., and Carroll, T.J. (2008). Beta-catenin is necessary to keep cells of ureteric bud/Wolffian duct epithelium in a precursor state. *Dev. Biol.* 314, 112–126.
- McGraw, T.E., Greenfield, L., and Maxfield, F.R. (1987). Functional expression of the human transferrin receptor cDNA in Chinese hamster ovary cells deficient in endogenous transferrin receptor. *J. Cell Biol.* 105, 207–214.
- McGuinness, M.P., Linder, C.C., Morales, C.R., Heckert, L.L., Pikus, J., and Griswold, M.D. (1994). Relationship of a mouse Sertoli cell line (MSC-1) to normal Sertoli cells. *Biol. Reprod.* 51, 116–124.
- Meyron-Holtz, E.G., Vaisman, B., Cabantchik, Z.I., Fibach, E., Rouault, T.A., Hershko, C., and Konijn, A.M. (1999). Regulation of intracellular iron metabolism in human erythroid precursors by internalized extracellular ferritin. *Blood* 94, 3205–3211.
- Michael, L., and Davies, J.A. (2004). Pattern and regulation of cell proliferation during murine ureteric bud development. *J. Anat.* 204, 241–255.
- Mollard, R., and Dziadek, M. (1998). A correlation between epithelial proliferation rates, basement membrane component localization patterns, and morphogenetic potential in the embryonic mouse lung. *Am. J. Respir. Cell Mol. Biol.* 19, 71–82.
- Moss, D., Powell, L.W., Arosio, P., and Halliday, J.W. (1992). Characterization of the ferritin receptors of human T lymphoid (MOLT-4) cells. *J. Lab. Clin. Med.* 119, 273–279.
- Ned, R.M., Swat, W., and Andrews, N.C. (2003). Transferrin receptor 1 is differentially required in lymphocyte development. *Blood* 102, 3711–3718.

- Nogawa, H., Morita, K., and Cardoso, W.V. (1998). Bud formation precedes the appearance of differential cell proliferation during branching morphogenesis of mouse lung epithelium in vitro. *Dev. Dyn.* 213, 228–235.
- Ohgami, R.S., Campagna, D.R., Greer, E.L., Antiochos, B., McDonald, A., Chen, J., Sharp, J.J., Fujiwara, Y., Barker, J.E., and Fleming, M.D. (2005). Identification of a ferrireductase required for efficient transferrin-dependent iron uptake in erythroid cells. *Nat. Genet.* 37, 1264–1269.
- Ojala, J.R., Pikkarainen, T., Tuuttila, A., Sandalova, T., and Tryggvason, K. (2007). Crystal structure of the cysteine-rich domain of scavenger receptor MARCO reveals the presence of a basic and an acidic cluster that both contribute to ligand recognition. *J. Biol. Chem.* 282, 16654–16666.
- Ponka, P., Beaumont, C., and Richardson, D.R. (1998). Function and regulation of transferrin and ferritin. *Semin. Hematol.* 35, 35–54.
- Puolakka, J., Janne, O., Pakarinen, A., and Vihko, R. (1980). Serum ferritin in the diagnosis of anemia during pregnancy. *Acta Obstet. Gynecol. Scand. Suppl.* 95, 57–63.
- Qiu, A., Jansen, M., Sakaris, A., Min, S.H., Chattopadhyay, S., Tsai, E., Sandoval, C., Zhao, R., Akabas, M.H., and Goldman, I.D. (2006). Identification of an intestinal folate transporter and the molecular basis for hereditary folate malabsorption. *Cell* 127, 917–928.
- Radisky, D.C., and Kaplan, J. (1998). Iron in cytosolic ferritin can be recycled through lysosomal degradation in human fibroblasts. *Biochem. J.* 336, 201–205.
- Rae, F., Woods, K., Sasmono, T., Campanale, N., Taylor, D., Ovchinnikov, D.A., Grimmond, S.M., Hume, D.A., Ricardo, S.D., and Little, M.H. (2007). Characterisation and trophic functions of murine embryonic macrophages based upon the use of a Csf1r-EGFP transgene reporter. *Dev. Biol.* 308, 232–246.
- Recalcati, S., Invernizzi, P., Arosio, P., and Cairo, G. (2008). New functions for an iron storage protein: the role of ferritin in immunity and autoimmunity. *J. Autoimmun.* 30, 84–89.
- San Martin, C.D., Garri, C., Pizarro, F., Walter, T., Theil, E.C., and Núñez, M.T. (2008). Caco-2 intestinal epithelial cells absorb soybean ferritin by mu2 (AP2)-dependent endocytosis. *J. Nutr.* 138, 659–666.
- Schmidt-Ott, K.M., Yang, J., Chen, X., Wang, H., Paragas, N., Mori, K., Li, J.Y., Lu, B., Costantini, F., Schiffer, M., et al. (2005). Novel regulators of kidney development from the tips of the ureteric bud. *J. Am. Soc. Nephrol.* 16, 1993–2002.
- Schmidt-Ott, K.M., Chen, X., Paragas, N., Levinson, R.S., Mendelsohn, C.L., and Barasch, J. (2006). c-kit delineates a distinct domain of progenitors in the developing kidney. *Dev. Biol.* 299, 238–249.
- Sibille, J.C., Ciriolo, M., Kondo, H., Crichton, R.R., and Aisen, P. (1989). Subcellular localization of ferritin and iron taken up by rat hepatocytes. *Biochem. J.* 262, 685–688.
- Siddappa, A.J., Rao, R.B., Wobken, J.D., Leibold, E.A., Connor, J.R., and Georgieff, M.K. (2002). Developmental changes in the expression of iron regulatory proteins and iron transport proteins in the perinatal rat brain. *J. Neurosci. Res.* 68, 761–775.
- Siddappa, A.M., Rao, R., Long, J.D., Widness, J.A., and Georgieff, M.K. (2007). The assessment of newborn iron stores at birth: a review of the literature and standards for ferritin concentrations. *Neonatology* 92, 73–82.
- Simpson, R.J., Raja, K.B., Halliwell, B., Evans, P.J., Aruoma, O.I., Konijn, A.M., and Peters, T.J. (1991). Iron speciation in hypotransferrinaemic mouse serum. *Biochem. Soc. Trans.* 19, 317S.
- Sposi, N.M., Cianetti, L., Tritarelli, E., Pelosi, E., Militi, S., Barberi, T., Gabbianelli, M., Saulle, E., Kuhn, L., Peschle, C., and Testa, U. (2000). Mechanisms of differential transferrin receptor expression in normal hematopoiesis. *Eur. J. Biochem.* 267, 6762–6774.
- Srinivas, S., Goldberg, M.R., Watanabe, T., D'Agati, V., al-Awqati, Q., and Costantini, F. (1999). Expression of green fluorescent protein in the ureteric bud of transgenic mice: a new tool for the analysis of ureteric bud morphogenesis. *Dev. Genet.* 24, 241–251.
- Srinivas, S., Watanabe, T., Lin, C.S., William, C.M., Tanabe, Y., Jessell, T.M., and Costantini, F. (2001). Cre reporter strains produced by targeted insertion of EYFP and ECFP into the ROSA26 locus. *BMC Dev. Biol.* 1, 4.
- Thorstensen, K., Trinder, D., Zak, O., and Aisen, P. (1995). Uptake of iron from N-terminal half-transferrin by isolated rat hepatocytes. Evidence of transferrin-receptor-independent iron uptake. *Eur. J. Biochem.* 232, 129–133.
- Tran, T.N., Eubanks, S.K., Schaffer, K.J., Zhou, C.Y., and Linder, M.C. (1997). Secretion of ferritin by rat hepatoma cells and its regulation by inflammatory cytokines and iron. *Blood* 90, 4979–4986.
- Trenor, C.C., 3rd, Campagna, D.R., Sellers, V.M., Andrews, N.C., and Fleming, M.D. (2000). The molecular defect in hypotransferrinemic mice. *Blood* 96, 1113–1118.
- Watanabe, K., Yamashita, Y., Ohgawara, H., Sekiguchi, M., Satake, N., Orino, K., and Yamamoto, S. (2001). Iron content of rat serum ferritin. *J. Vet. Med. Sci.* 63, 587–589.
- Wingert, R.A., Brownlie, A., Galloway, J.L., Dooley, K., Fraenkel, P., Axe, J.L., Davidson, A.J., Barut, B., Noriega, L., Sheng, X., et al. (2004). The chianti zebrafish mutant provides a model for erythroid-specific disruption of transferrin receptor1. *Development* 131, 6225–6235.
- Xu, J., and Ling, E.A. (1994). Studies of the ultrastructure and permeability of the blood-brain barrier in the developing corpus callosum in postnatal rat brain using electron dense tracers. *J. Anat.* 184, 227–237.
- Yuan, X.M., Li, W., Baird, S.K., Carlsson, M., and Meleforts, O. (2004). Secretion of ferritin by iron-laden macrophages and influence of lipoproteins. *Free Radic. Res.* 38, 1133–1142.