


Contents lists available at ScienceDirect

International Journal of Infectious Diseases

journal homepage: www.elsevier.com/locate/ijid

Aerosolized amphotericin B as prophylaxis for invasive pulmonary aspergillosis: a meta-analysis



Di Xia^{a,1}, Wen-Kui Sun^{a,1}, Ming-Ming Tan^a, Ming Zhang^a, Yuan Ding^a, Zhi-Cheng Liu^b, Xin Su^{a,*}, Yi Shi^{a,*}

^a Department of Respiratory and Critical Care Medicine, Jinling Hospital, Medical School of Nanjing University, Nanjing, People's Republic of China

^b Southern Medical University, Guangdong, People's Republic of China

ARTICLE INFO

Article history:

Received 6 October 2014

Received in revised form 3 November 2014

Accepted 4 November 2014

Corresponding Editor: Eskild Petersen, Aarhus, Denmark

Keywords:

Aerosolized amphotericin B
Invasive pulmonary aspergillosis
Prophylaxis
Meta-analysis
Immunosuppression

SUMMARY

Objectives: Invasive pulmonary aspergillosis (IPA) is associated with high mortality in high-risk (immunosuppressed) patients. Many studies have investigated whether prophylactic inhalation of amphotericin B (AMB) reduces the incidence of IPA, but no definitive conclusions have been reached. The present meta-analysis was performed to evaluate the efficacy of prophylactic inhalation of AMB for the prevention of IPA.

Methods: MEDLINE and other databases were searched for relevant articles published until December 2013. Randomized controlled trials that compared aerosolized AMB with placebo were included. Two reviewers independently assessed and extracted the data of all trials.

Results: Six animal studies and two clinical trials involving 768 high-risk patients were eligible. The animal studies showed lower overall mortality rate among animals that underwent aerosolized AMB prophylaxis (odds ratio (OR) 0.13, 95% confidence interval (CI) 0.08–0.21). Similarly, the clinical trials showed a lower incidence of IPA among patients who underwent aerosolized AMB prophylaxis (OR 0.42, 95% CI 0.22–0.79).

Conclusions: This analysis provides evidence supporting the notion that the prophylactic use of aerosolized AMB effectively reduces the incidence of IPA among high-risk patients.

© 2014 The Authors. Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

1. Introduction

Invasive aspergillosis (IA) is an increasingly frequent cause of morbidity and mortality in immunosuppressed patients, especially those undergoing solid organ or hematopoietic stem cell transplantation and those with prolonged neutropenia.¹ Invasive pulmonary aspergillosis (IPA) is the most common form of IA. Despite the fact that new non-invasive laboratory methods have been developed to improve the diagnostic yield, including the *Aspergillus* galactomannan assay, the (1,3)- β -D-glucan assay, and PCR techniques, IPA remains associated with a high fatality rate. In one systematic review, 70% of 1941 patients with aspergillosis exhibited pulmonary involvement, and the case-fatality rate was >60% despite the administration of intensive antifungal therapy.² Therefore, prophylactic therapy is important in high-risk patients. However, there is no consensus on the optimal agent or administration route.

Amphotericin B (AMB) was the first commercially significant antifungal drug. It has a broad spectrum of activity against many different fungal species and has been the standard IA treatment for decades.³ Although new agents such as voriconazole and itraconazole have been recommended for patients with IPA, AMB is still considered to be the primary therapeutic agent for some patients and is included in many prophylactic regimens for fungal infection.⁴ One study showed that the prophylactic administration of intravenous AMB to patients undergoing bone marrow transplantation was associated with fewer fungal microorganisms and higher survival rates compared to the placebo group; however, significantly greater numbers of infusion-related side effects occurred.⁵ Therefore, aerosolized AMB represents an attractive alternative for the prevention of IPA because the administration of drugs by inhalation ensures a high drug concentration in the respiratory tract and a lower incidence of side effects.

Since the 1990s, many studies have been conducted to elucidate the feasibility, tolerability, and effectiveness of aerosolized AMB for the prevention of *Aspergillus* infection.^{6–11} A retrospective study of 99 patients who underwent heart transplantation with

* Corresponding authors. Tel.: +86 25 80860148; fax: +86 25 80860148.

E-mail addresses: suxinjs@163.com (X. Su), shiyi56@126.com (Y. Shi).

¹ Both authors contributed equally to this article and were co-first authors.

no prophylaxis and 120 patients who underwent prophylactic inhalation of AMB demonstrated a significant difference between the two groups; prophylaxis with AMB effectively prevented IPA.¹² Another retrospective study evaluated the impact of prophylactic AMB inhalation on IA in 611 recipients of allogeneic stem cell transplantation and examined the recipients' tolerance of the inhalation therapy. The incidence of IA was lower in the prophylactic AMB inhalation group than in the placebo group, and the inhalation therapy was well tolerated.¹³ However, other studies have reached different conclusions. In another study that investigated the effectiveness of aerosolized AMB as prophylaxis against IPA, 28% of the patients developed proven or possible infections. Inhalation of AMB does not appear to be useful in preventing IPA in patients with granulocytopenia.¹⁴

The present meta-analysis was performed to assess the prophylactic effect of aerosolized AMB against IPA by examining the IPA-associated mortality among immunocompromised animals and the incidence of IPA among high-risk patients.

2. Materials and methods

2.1. Search strategy

Two separate electronic searches were conducted to identify eligible studies. MEDLINE, Embase, the Chinese Biomedical Literature Database, and the Cochrane Library were searched for relevant articles published until December 25, 2013. The following search terms were used: “inhaled” or “inhalational” or “aerosol” or “aerosolized” or “nebulized” or “nebulization” and “amphotericin”. No limitations were placed on language or year. The reference lists of related reviews and original papers were also checked for relevant trials.

2.2. Study selection

The following inclusion criteria were established before article collection. Animal studies were required to (1) be randomized controlled trials, (2) compare aerosolized AMB with placebo, (3) administer aerosolized AMB before exposure to *Aspergillus fumigatus* conidia, and (4) provide the number of animals sacrificed. Human studies were required to (1) be randomized controlled trials, (2) include adult patients (aged >18 years) scheduled to receive chemotherapy with an anticipated duration of neutropenia $<0.5 \times 10^9$ cells/l of ≥ 10 days, (3) compare aerosolized AMB with placebo, and (4) administer aerosolized AMB before any signs of proven or probable IPA. When an individual author published several articles involving the same patient population, only the most complete article was included. Studies that did not meet the above-described inclusion criteria were excluded from the meta-analysis.

2.3. Quality assessment

Clinical randomized controlled trials were assessed using the Jadad scale.¹⁵ This scale is used to assess trials according to the following three questions: (1) Was the study described as randomized (i.e., did it use the terms ‘randomly’, ‘random’, or ‘randomization’)? (0–2 points); (2) Was the study described as double-blind? (0–2 points); (3) Was there a description of withdrawals and dropouts? (0–1 point). A study can receive a maximum Jadad score of 5 points.

2.4. Data extraction

Two reviewers (DX and WKS) independently carried out the data extraction and validity assessment, and any discrepancies

were resolved by discussion. For the animal studies, a piloted data extraction form was used to collect information on the first author, year of publication, animal species, number of animals in each group, method of inducing immunosuppression, details of experimental drug and placebo treatments, follow-up duration, and final mortality rate. For the clinical trials, a data extraction form was used to collect information on the first author, year of publication, country of origin, Jadad score, number of patients in each group, and incidence of IPA.

2.5. Statistical analysis

The results of prophylaxis for dichotomous outcomes are expressed as odds ratios (ORs) with 95% confidence intervals (CIs) for both the animal studies and clinical trials. The I^2 statistic was used to determine the extent of inconsistency and thus assess the heterogeneity between trials. We considered an I^2 -value of >50% and a p -value of <0.1 to indicate heterogeneity. A fixed-effects model was used to estimate the effects of aerosolized AMB. However, if significant heterogeneity was present, a random-effects model was used to generate a more conservative estimate.

Publication bias among the randomized controlled trials involving animals was examined by visual inspection of a funnel plot. Publication bias was suspected when the funnel plot was asymmetrical; in such cases, Egger's test was performed for further analysis of bias.

Sensitivity analyses were conducted by comparing the estimates derived from the random- and fixed-effects models. One study that used AMB inhalation powder (ABIP) as the prophylactic drug was excluded from the sensitivity analyses because this drug is not widely used.

Subgroup analyses of the animal studies were performed to explore important differences that might be expected to alter the magnitude of the prophylactic effect.

3. Results

3.1. Study selection and characteristics

Figure 1 shows the study selection process. In total, 1362 potentially relevant citations were identified from the electronic search, 1348 of which were determined to be non-relevant after reading the titles and abstracts. The remaining 14 studies underwent full review by the two above-mentioned independent reviewers. Eight of these 14 studies met the inclusion criteria and were subjected to the meta-analysis.^{16–23} Six studies were initially thought to fulfill the inclusion criteria, but were excluded after detailed examination. One study was not a randomized controlled trial,²⁴ one evaluated the therapeutic rather than the prophylactic efficacy of aerosolized AMB,²⁵ one evaluated the beneficial effect of intravenous rather than aerosolized AMB,²⁶ one evaluated the beneficial effect of aerosolized AMB on the fungal burden rather than on mortality,²⁷ and two were duplicate publications.^{10,28} Of the eight remaining eligible studies, six were animal randomized controlled trials^{16–21} and two were human randomized controlled trials.^{22,23}

In all six animal studies, a systemic steroid and/or cyclophosphamide was used to induce immunosuppression. The fungal inoculation and drug administration methods were described in detail. The various formulations of aerosolized AMB were AMB desoxycholate (AMB-d), liposomal AMB (L-AMB), AMB lipid complex (ABLC), AMB colloidal dispersion (ABCD), and ABIP. Table 1 lists the details of the six animal studies included in this meta-analysis.

In both of the human studies, randomization was performed using a computer-generated blocked list. Both studies included a description of the patients who withdrew from or dropped out of




Figure 1. Flow diagram of study selection method
AMB: amphotericin B

the study, but only one trial was double-blind.²³ L-AMB was used in one trial and AMB-d in the other. The details of these two trials are given in Table 2. Both clinical trials mentioned the potential toxic effects of the therapy, and one described the toxic effects in detail.²² For the patients who received aerosolized L-AMB, the median serum creatinine levels after the last inhalation were not greater than the baseline levels, but coughing was observed more frequently than at baseline.²³ About two-thirds of patients who received aerosolized AMB-d reported at least one unpleasant sensation such as coughing, a bad taste, nausea, or others.²² No serious drug-related adverse events were reported.

3.2. Meta-analysis results

Six studies of immunosuppressed animals were eligible for inclusion in the meta-analysis. The overall mortality of animals treated with prophylactic inhalation of AMB was lower than that of animals treated with placebo. No heterogeneity was observed ($I^2 = 7\%$, $p = 0.36$) and a fixed-effects model was used. The combined OR for all six eligible studies was 0.13 (95% CI 0.08–0.21; $p < 0.00001$) (Figure 2), indicating that the prophylactic use of aerosolized AMB was effective in immunocompromised animals. No significant difference ($p = 0.28$) was observed between the

Table 1
Characteristics of the animal studies included in the meta-analysis

Author (year)	Animal	Method of inducing immunosuppression	Aspergillus inoculum dose	AMB administration dose and time prior to pulmonary inoculation	Time point of mortality measure
Schmitt (1988)	Rat	Steroid (100 mg/kg) administered 2 weeks before fungal inoculation and continued throughout the experiment	10^6 conidia	AMB-d: 1.6 mg/kg 2 days	21 days after fungal inoculation
Niki (1991)	Rat	Steroid (150 mg/kg) administered three times weekly, 2 weeks before and 1 week after fungal inoculation	10^6 conidia	AMB-d: 1.6 mg/kg 48 h	4 weeks after fungal inoculation
Allen (1994)	Mouse	Steroid (150 mg/kg) administered 1 day before until 1 day after fungal inoculation	1.4×10^6 1.5×10^7 1.3×10^8 conidia	L-AMB: 6.05 mg/kg AMB-d: 6.73 mg/kg 1, 2, and 3 days	9 days after fungal inoculation
Cicogna (1997)	Rat	Steroid (150 mg/kg) administered for 2 weeks until the day of fungal inoculation or steroid (150 mg/kg) administered for 2 weeks before fungal inoculation and continued throughout the experiment	10^6 conidia	ABL: 0.4, 0.8, and 1.6 mg/kg AMB-d: 1.6 mg/kg ABL: 1.6 mg/kg 2 days	14 days after fungal inoculation
Ruijgrok (2005)	Rat	Cyclophosphamide (90 mg/kg) administered 5 days before and cyclophosphamide (60 mg/kg) administered every 4 days after fungal inoculation	1.5×10^5 conidia	AMB-d: 2 mg/ml L-AMB: 4 mg/ml ABL: 4 mg/ml 1, 2, and 6 weeks	12 days after fungal inoculation
Kirkpatrick (2012)	Guinea pig	Cyclophosphamide (250 mg/kg) and steroid (250 mg/kg) administered 2 days before and 3 days after fungal inoculation	1×10^8 conidia	ABIP: 0.05, 0.50, 4.00, and 10.00 mg/kg 24 h	11 days after fungal inoculation

AMB, amphotericin B; AMB-d, amphotericin B desoxycholate; L-AMB, liposomal amphotericin B; ABL, amphotericin B lipid complex; ABIP, amphotericin B inhalation powder.

Table 2
Characteristics of clinical trials included in the meta-analysis

Author (year)	Study period	Study region	Patients	Jadad quality score	Treatment group, IPA/total	Placebo group, IPA/total
Schwartz (1999)	1993/03–1996/04	Germany	Adult patients with hematological disease and neutropenia	3	10/227	11/155
Rijnders (2008)	2000/11–2006/01	Netherlands	Adult patients with hematological disease or solid tumors and neutropenia	5	6/139	18/132

IPA, invasive pulmonary aspergillosis.

effectiveness of AMB-d (OR 0.07, 95% CI 0.03–0.18; $p < 0.00001$) (Figure 3A) and that of lipid-associated AMB formulations (OR 0.06, 95% CI 0.03–0.14; $p < 0.00001$) (Figure 3B).

Two human trials involving 768 high-risk patients were eligible for inclusion in the meta-analysis. The incidence of IPA in patients who underwent administration of aerosolized AMB during neutropenic episodes was lower than that of patients who underwent administration of placebo (4.4% vs. 10.4%, respectively). No evidence of heterogeneity was observed ($I^2 = 21\%$, $p = 0.26$), and a fixed-effects model was used. Aerosolized AMB demonstrated a significant preventive advantage over placebo in terms of a lower incidence of IPA (OR 0.42, 95% CI 0.22–0.79; $p = 0.007$) (Figure 4).

3.3. Sensitivity analyses

Despite the absence of statistical heterogeneity, significant trial heterogeneity was present across the analyzed studies (different animal types, AMB formulations, AMB doses, etc.). Therefore, we

performed a sensitivity analysis of the six animal studies by repeating the main computations using a random-effects model. The random-effects model did not significantly change the results of our meta-analysis. Similarly, other sensitivity analyses showed no changes in the results after exclusion of specific studies (Table 3). Therefore, the results of this meta-analysis are stable.

3.4. Publication bias

The funnel plot of the animal studies was asymmetrical, suggesting possible publication bias (Figure 5). Egger's test was then performed to check for bias; the result was significant ($p = 0.01$), again suggesting possible publication bias.

4. Discussion

This meta-analysis has shown that aerosolized AMB can help to prevent IA in both immunocompromised animals and high-risk





Figure 2. Forest plot showing effect of prophylactic aerosolized amphotericin B on mortality of immunosuppressed animals
M-H: Mantel–Haenszel analysis, CI: confidence interval

A



B




Figure 3. Forest plot showing effect of prophylactic aerosolized amphotericin B desoxycholate (A) and lipid-associated amphotericin B (B) on mortality of immunosuppressed animals.

M-H: Mantel-Haenszel analysis, CI: confidence interval




Figure 4. Forest plot showing effect of prophylactic aerosolized amphotericin B on incidence of invasive pulmonary aspergillosis in high-risk patients

M-H: Mantel-Haenszel analysis, CI: confidence interval

patients. Such high-risk patients include recipients of hematopoietic stem cell or solid organ transplantation, patients with malignancies undergoing intensive chemotherapy, and patients with other causes of immunosuppression. Asymptomatic patients with *Aspergillus galactomannan* in the bronchoalveolar lavage

fluid or serum may especially benefit from the prophylactic use of aerosolized AMB.

Aerosolized AMB is relatively safe. To the best of our knowledge, no serious drug-related adverse events have been reported in association with its prophylactic use. In one study, patients who

Table 3
Sensitivity analyses for the primary outcome

	Number of studies	OR (95% CI)	p-Value
Random-effects model	6	0.13 (0.07–0.22)	<0.00001
Exclusion of one study that used ABIP	5	0.06 (0.03–0.12)	<0.00001

OR, Odds ratio; 95% CI, 95% confidence interval; ABIP, amphotericin B inhalation powder.

underwent administration of aerosolized AMB as antifungal prophylaxis considered the inhalations to be unpleasant mostly because of a bad taste or the development of coughing. However, the inhalation therapy was feasible and safe, and no severe side effects occurred.²⁹ Monforte et al. found that nebulized L-AMB for prophylactic treatment of *Aspergillus* infection exhibits neither significant systemic absorption nor adverse effects on respiratory function.³⁰ They also found that nebulized L-AMB does not change the lipid content of pulmonary surfactant. This agent safely and effectively prevents *Aspergillus spp* infection in lung transplant recipients.³¹

Different AMB formulations may have different clinical effects because each has a distinct pharmacological profile. Drew et al. concluded that patients who received AMB-d are more likely to experience adverse events.³² However, our subgroup analysis of animal studies showed no significant difference between AMB-d and lipid-associated AMB formulations. This result is consistent with that obtained in a previous observational study in which 104 consecutive patients who underwent prophylaxis with aerosolized L-AMB were compared with 49 historical control subjects who received aerosolized AMB-d. The two groups exhibited similar rates of *Aspergillus* infection and side effects such as transitory breathing difficulty, nausea, and bronchospasm.³³ One systematic review and meta-analysis showed no difference between the adverse events associated with inhaled AMB-d and those associated with lipid formulations of inhaled AMB.³⁴ A worldwide survey on antifungal prophylaxis in patients undergoing lung transplantation also revealed that inhaled lipid formulations of AMB are effective and being used with increased frequency.³⁵ Therefore, lipid formulations may be more effective than AMB-d in preventing IPA.

New formulations of aerosolized AMB have recently been developed. Lipid nanoemulsions may serve as successful nano-carriers for the delivery of AMB to the peripheral airways.³⁶ Nonionic surfactant vesicles that deliver AMB to the lungs

reportedly enhance pulmonary delivery while minimizing systemic exposure and toxicity.³⁷ Further studies comparing the effects of these various formulations on the prevention of IPA are also needed.

New antifungal agents in the azole group have been developed in recent years. They are also prescribed as prophylaxis against IPA. Neoh et al.³⁸ conducted a retrospective cohort study to explore the effect of prophylactic voriconazole in lung transplant recipients. They concluded that preemptive voriconazole treatment resulted in a lower incidence of IA and a lower IA-related mortality rate. Another retrospective study suggested that the routine use of prophylactic voriconazole against *Aspergillus* infection in lung transplant recipients did not appear to be warranted.³⁹ Additionally, single-agent itraconazole treatment in heart or lung transplant recipients did not affect the incidence of fungal infection as compared with a control group.⁴⁰ To the best of our knowledge, no definitive guidelines on the prophylactic use of azole agents in IPA have been established, and few studies comparing aerosolized AMB and azole agents for the prevention of IPA have been performed. One study assessed the efficacy of an inhaled aqueous solution of voriconazole as prophylaxis against IPA in a murine model.⁴¹ Rodents with IPA that underwent treatment with inhaled voriconazole demonstrated significantly higher survival than those treated with AMB. However, AMB-d was administered intraperitoneally. Therefore it remains unknown whether inhaled voriconazole is superior to aerosolized AMB. Clinical trials comparing azoles and inhaled AMB are essential to shed light on the question of which agent and administration method is optimal for IPA prophylaxis. Meanwhile, cost should be taken into consideration when comparing different antifungal agents.

Some limitations of this meta-analysis should be noted. First, because the analysis was limited to the published scientific literature, the potential impact of publication bias cannot be ignored. Publication bias is a known threat to the validity of all forms of meta-analysis. Journals tend to accept positive results, while negative results are often rejected or not even submitted by authors. Second, none of the animal studies in the present meta-analysis explicitly described the blinding or allocation concealment methods used. Future studies should clearly explain the details of their blinding and allocation concealment methods. Third, all placebo groups in the present studies were used more than once to compare the effects of the different interventions. Both the placebo and experimental subgroups in some eligible studies exhibited a 100% mortality rate. However, this high mortality rate was not included in the overall estimate of this meta-analysis, making the combined OR appear to be much stronger. Fourth, the number of eligible clinical trials was small. Thus, the relatively small number of participants might not allow for a reliable conclusion. Fifth, of the two clinical studies, one did not mention the performance of an intention-to-treat analysis and was conducted in an unblinded fashion,²² which may have resulted in high performance, measurement, and selection biases. Finally, a cost-effectiveness analysis was not performed; this may have given rise to a potential preference for the use of aerosolized AMB as prophylaxis for fungal infections in an era when many other antifungal drugs are available.

In conclusion, aerosolized AMB effectively reduces the incidence of IPA in high-risk patients and has proved to be useful clinically when used as prophylaxis. However, its effects should be confirmed in large sample-size, multicenter, randomized controlled trials. Analysis of cost-effectiveness and adverse effects between various aerosolized AMB formulations and newer antifungal agents should be included in future clinical trials.




Figure 5. Funnel plot showing absence of small negative studies suggestive of small publication bias.

Acknowledgements

This study was funded by the National Natural Science Foundation of China (grant numbers 81270064 and 81200063) and the Hospital Foundation of Jinling Hospital (grant number 2013021).

Conflict of interest: The authors declare no conflicts of interest. Ethical approval was not required.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.ijid.2014.11.004>.

References

- Segal BH. Aspergillosis. *N Engl J Med* 2009;**360**:1870–84.
- Lin SJ, Schranz J, Teutsch SM. Aspergillosis case-fatality rate: systematic review of the literature. *Clin Infect Dis* 2001;**32**:358–66.
- Garbati MA, Alasmari FA, Al-Tannir MA, Tleyjeh IM. The role of combination antifungal therapy in the treatment of invasive aspergillosis: a systematic review. *Int J Infect Dis* 2012;**16**:e76–81.
- Walsh TJ, Anaissie EJ, Denning DW, Herbrecht R, Kontoyiannis DP, Marr KA, et al. Treatment of aspergillosis: clinical practice guidelines of the Infectious Diseases Society of America. *Clin Infect Dis* 2008;**46**:327–60.
- Perfect JR, Klotman ME, Gilbert CC, Crawford DD, Rosner GL, Wright KA, et al. Prophylactic intravenous amphotericin B in neutropenic autologous bone marrow transplant recipients. *J Infect Dis* 1992;**165**:891–7.
- Conneally E, Cafferkey MT, Daly PA, Keane CT, McCann SR. Nebulized amphotericin B as prophylaxis against invasive aspergillosis in granulocytopenic patients. *Bone Marrow Transplant* 1990;**5**:403–6.
- Myers SE, Devine SM, Topper RL, Ondrey M, Chandler C, O'Toole K, et al. A pilot study of prophylactic aerosolized amphotericin B in patients at risk for prolonged neutropenia. *Leuk Lymphoma* 1992;**8**:229–33.
- Beyer J, Barzen G, Risse G, Weyer C, Miksits K, Dullenkopf K, et al. Aerosol amphotericin B for prevention of invasive pulmonary aspergillosis. *Antimicrob Agents Chemother* 1993;**37**:1367–9.
- Beyer J, Schwartz S, Barzen G, Risse G, Dullenkopf K, Weyer C, et al. Use of amphotericin B aerosols for the prevention of pulmonary aspergillosis. *Infection* 1994;**22**:143–8.
- Behre G, Schwartz S, Lenz K, Ludwig WD, Wandt H, Schilling E, et al. Aerosol amphotericin B inhalations for prevention of invasive pulmonary aspergillosis in neutropenic cancer patients. *Ann Hematol* 1995;**71**:287–91.
- Monforte V, Roman A, Gavalda J, Bravo C, Tenorio L, Ferrer A, et al. Nebulized amphotericin B prophylaxis for Aspergillus infection in lung transplantation: study of risk factors. *J Heart Lung Transplant* 2001;**20**:1274–81.
- Paniagua Martin MJ, Marzoa Rivas R, Barge Caballero E, Grille Cancela Z, Fernandez CJ, Solla M, et al. Efficacy and tolerance of different types of prophylaxis for prevention of early aspergillosis after heart transplantation. *Transplant Proc* 2010;**42**:3014–6.
- Nihtinen A, Anttila VJ, Ruutu T, Juvonen E, Volin L. Low incidence of invasive aspergillosis in allogeneic stem cell transplant recipients receiving amphotericin B inhalation prophylaxis. *Transpl Infect Dis* 2012;**14**:24–32.
- Erjavec Z, Woolthuis GM, de Vries-Hospers HG, Sluiter WJ, Daenen SM, de Pauw B, et al. Tolerance and efficacy of amphotericin B inhalations for prevention of invasive pulmonary aspergillosis in haematological patients. *Eur J Clin Microbiol Infect Dis* 1997;**16**:364–8.
- Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJ, Gavaghan DJ, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Controlled Clinical Trials* 1996;**17**:1–12.
- Schmitt HJ, Bernard EM, Hauser M, Armstrong D. Aerosol amphotericin B is effective for prophylaxis and therapy in a rat model of pulmonary aspergillosis. *Antimicrob Agents Chemother* 1988;**32**:1676–9.
- Niki Y, Bernard EM, Edwards FF, Schmitt HJ, Yu B, Armstrong D. Model of recurrent pulmonary aspergillosis in rats. *J Clin Microbiol* 1991;**29**:1317–22.
- Allen SD, Sorensen KN, Nejd MJ, Durrant C, Proffitt RT. Prophylactic efficacy of aerosolized liposomal (AmBisome) and non-liposomal (Fungizone) amphotericin B in murine pulmonary aspergillosis. *J Antimicrob Chemother* 1994;**34**:1001–13.
- Cicogna CE, White MH, Bernard EM, Ishimura T, Sun M, Tong WP, et al. Efficacy of prophylactic aerosol amphotericin B lipid complex in a rat model of pulmonary aspergillosis. *Antimicrob Agents Chemother* 1997;**41**:259–61.
- Ruijgrok EJ, Fens MH, Bakker-Woudenberg IA, van Etten EW, Vulto AG. Nebulization of four commercially available amphotericin B formulations in persistently granulocytopenic rats with invasive pulmonary aspergillosis: evidence for long-term biological activity. *J Pharm Pharmacol* 2005;**57**:1289–95.
- Kirkpatrick WR, Najvar LK, Vallor AC, Wiederhold NP, Bocanegra R, Pfeiffer J, et al. Prophylactic efficacy of single dose pulmonary administration of amphotericin B inhalation powder in a guinea pig model of invasive pulmonary aspergillosis. *J Antimicrob Chemother* 2012;**67**:970–6.
- Schwartz S, Behre G, Heinemann V, Wandt H, Schilling E, Arning M, et al. Aerosolized amphotericin B inhalations as prophylaxis of invasive Aspergillus infections during prolonged neutropenia: results of a prospective randomized multicenter trial. *Blood* 1999;**93**:3654–61.
- Rijnders BJ, Cornelissen JJ, Slobbe L, Becker MJ, Doorduyn JK, Hop WC, et al. Aerosolized liposomal amphotericin B for the prevention of invasive pulmonary aspergillosis during prolonged neutropenia: a randomized, placebo-controlled trial. *Clin Infect Dis* 2008;**46**:1401–8.
- Reichenspurner H, Gamberg P, Nitschke M, Valantine H, Hunt S, Oyer PE, et al. Significant reduction in the number of fungal infections after lung-, heart–lung, and heart transplantation using aerosolized amphotericin B prophylaxis. *Transplant Proc* 1997;**29**:627–8.
- Sheppard DC, Rieg G, Chiang LY, Filler SG, Edwards Jr JE, Ibrahim AS. Novel inhalational murine model of invasive pulmonary aspergillosis. *Antimicrob Agents Chemother* 2004;**48**:1908–11.
- Olson JA, Adler-Moore JP, Schwartz J, Jensen GM, Proffitt RT. Comparative efficacies, toxicities, and tissue concentrations of amphotericin B lipid formulations in a murine pulmonary aspergillosis model. *Antimicrob Agents Chemother* 2006;**50**:2122–31.
- Leleu C, Menotti J, Meneceur P, Choukri F, Sulahian A, Garin YJ, et al. Efficacy of liposomal amphotericin B for prophylaxis of acute or reactivation models of invasive pulmonary aspergillosis. *Mycoses* 2013;**56**:241–9.
- Slobbe L, Boersma E, Rijnders BJ. Tolerability of prophylactic aerosolized liposomal amphotericin-B and impact on pulmonary function: data from a randomized placebo-controlled trial. *Pulm Pharmacol Ther* 2008;**21**:855–9.
- Hullard-Pulstinger A, Holler E, Hahn J, Andreesen R, Krause SW. Prophylactic application of nebulized liposomal amphotericin B in hematologic patients with neutropenia. *Onkologie* 2011;**34**:254–8.
- Monforte V, Ussetti P, Lopez R, Gavalda J, Bravo C, de Pablo A, et al. Nebulized liposomal amphotericin B prophylaxis for Aspergillus infection in lung transplantation: pharmacokinetics and safety. *J Heart Lung Transplant* 2009;**28**:170–5.
- Monforte V, Lopez-Sanchez A, Zurbano F, Ussetti P, Sole A, Casals C, et al. Prophylaxis with nebulized liposomal amphotericin B for Aspergillus infection in lung transplant patients does not cause changes in the lipid content of pulmonary surfactant. *J Heart Lung Transplant* 2013;**32**:313–9.
- Drew RH, Dods Ashley E, Benjamin DK, Duane Davis R, Palmer SM, Perfect JR. Comparative safety of amphotericin B lipid complex and amphotericin B deoxycholate as aerosolized antifungal prophylaxis in lung-transplant recipients. *Transplantation* 2004;**77**:232–7.
- Monforte V, Ussetti P, Gavalda J, Bravo C, Laporta R, Len O, et al. Feasibility, tolerability, and outcomes of nebulized liposomal amphotericin B for Aspergillus infection prevention in lung transplantation. *J Heart Lung Transplant* 2010;**29**:523–30.
- Bhaskaran A, Mumtaz K, Husain S. Anti-Aspergillus prophylaxis in lung transplantation: a systematic review and meta-analysis. *Curr Infect Dis Rep* 2010;**12**:251–5.
- Neoh CF, Snell GI, Kotsimbos T, Levvey B, Morrissey CO, Slavin MA, et al. Antifungal prophylaxis in lung transplantation—a world-wide survey. *Am J Transplant* 2011;**11**:361–6.
- Nasr M, Nawaz S, Elhissi A. Amphotericin B lipid nanoemulsion aerosols for targeting peripheral respiratory airways via nebulization. *Int J Pharm* 2012;**436**:611–6.
- Alsaadi M, Italia JL, Mullen AB, Ravi Kumar MN, Candlish AA, Williams RA, et al. The efficacy of aerosol treatment with non-ionic surfactant vesicles containing amphotericin B in rodent models of leishmaniasis and pulmonary aspergillosis infection. *J Control Release* 2012;**160**:685–91.
- Neoh CF, Snell GI, Levvey B, Kotsimbos T, Morrissey CO, Slavin MA, et al. Preemptive treatment with voriconazole in lung transplant recipients. *Transpl Infect Dis* 2013;**15**:344–53.
- Tofte N, Jensen C, Tvede M, Andersen CB, Carlsen J, Iversen M. Use of prophylactic voriconazole for three months after lung transplantation does not reduce infection with Aspergillus: a retrospective study of 147 patients. *Scand J Infect Dis* 2012;**44**:835–41.
- Hayes Jr D, Ball AM, Mansour HM, Martin CA, Flynn JD. Fungal infection in heart–lung transplant recipients receiving single-agent prophylaxis with itraconazole. *Exp Clin Transplant* 2011;**9**:399–404.
- Tolman JA, Wiederhold NP, McConville JT, Najvar LK, Bocanegra R, Peters JI, et al. Inhaled voriconazole for prevention of invasive pulmonary aspergillosis. *Antimicrob Agents Chemother* 2009;**53**:2613–5.