

## Amino acid sequence of a heat-stable enterotoxin isolated from enterotoxigenic *Escherichia coli* strain 18D

Toshifumi Takao<sup>+</sup>, Tetsuo Hitouji<sup>+</sup>, Saburo Aimoto<sup>+</sup>, Yasutsugu Shimonishi<sup>+,\*</sup>, Saburo Hara<sup>†</sup>, Tae Takeda<sup>++</sup>, Yoshifumi Takeda<sup>++</sup> and Toshio Miwatani<sup>++</sup>

<sup>+</sup>*Institute for Protein Research*, <sup>†</sup>*Department of Chemistry, Faculty of Science* and <sup>++</sup>*Research Institute for Microbial Diseases, Osaka University, Suita, Osaka 565, Japan*

Received 8 November 1982

A heat-stable enterotoxin produced by a strain of enterotoxigenic *Escherichia coli* 18D was purified by ion-exchange and reversed-phase high-pressure liquid chromatography. The amino acid sequence of the purified toxin was determined by Edman-degradation and a combination of fast atom bombardment mass spectrometry and carboxypeptidase digestion to be Asn-Thr-Phe-Tyr-Cys-Cys-Glu-Leu-Cys-Cys-Asn-Pro-Ala-Cys-Ala-Gly-Cys-Tyr.

*Enterotoxin*      *Isolation*      *Amino acid sequence*      *FAB mass spectrometry*

### 1. INTRODUCTION

Enterotoxigenic *Escherichia coli* (ETEC) produces at least two types of enterotoxins that cause diarrheal disease in man and various domestic animals: low- $M_r$ , heat-stable enterotoxin (ST), and high- $M_r$ , heat-labile enterotoxin (LT) [1]. It was recently confirmed [2,3] that there are two genetically distinct ST's, which were named STIa and STIb. The DNA sequences encoding STIa and STIb were determined in [3] and [4], respectively. Since STIa and STIb originated from porcine and human strains of ETEC, we here designate them as ST<sub>p</sub> and ST<sub>h</sub>, respectively.

Recently, we isolated and purified ST<sub>h</sub> from ETEC of human patients [5,6] and determined its amino acid sequence of 19 amino acid residues [6]. The sequence was different from that deduced from the DNA sequence encoding ST<sub>p</sub> but identical to that deduced from the DNA sequence of ST<sub>h</sub>, except for the C-terminal amino acid residue. In [7] the amino acid sequence of an ST isolated from an ETEC strain 18D, was not similar to that

of ST<sub>h</sub> but was similar to that deduced from the DNA sequence of ST<sub>p</sub>, although the strain was derived from human patients. However, strain 18D was found by the DNA hybridization technique to be a strain producing ST<sub>p</sub> (P. Echeverria, personal communication).

Here, we describe the isolation of an ST (= ST<sub>p</sub>) from ETEC strain 18D and its sequence determination by Edman-degradation and a combination of fast atom bombardment (FAB) mass spectrometry and carboxypeptidase digestion [8,9].

### 2. MATERIALS AND METHODS

#### 2.1. Bacterial strain and culture conditions

*E. coli* 18D, kindly given by Dr Giannella (University of Cincinnati, OH), was used throughout. The cells were grown in CAYE medium [10] at pH 8.6 without glucose at 37°C for 24 h with vigorous shaking, as in [5].

#### 2.2. Isolation of the toxin

Crude toxin was extracted from the culture filtrate (20 L), as in [5]. The crude toxin was chromatographed successively on SP-Sephadex C-50 (H<sup>+</sup> form) and DEAE-Sephadex A-25 (acetate

\* To whom correspondence should be addressed

form) as in [6]. In each case, the main fraction with biological activity was collected for further purification. Final purification was achieved by reversed-phase HPLC on a LiChrosorb RP-8 column under the conditions described in the legend to fig.3.

### 2.3. Biological assay

ST activity was assayed in suckling mice of 2–4 days old, as in [5]. The fluid accumulation ratio of each animal was calculated as the ratio of the weight of the entire intestine to that of the rest of the body. The minimal amount of ST giving a fluid accumulation ratio of over 0.09 was designated as 1 mouse unit (MU), as in [5].

### 2.4. Amino acid analysis

Purified toxin and its derivatives were hydrolyzed in 4 M methane-sulfonic acid for 24 h at 105°C in vacuum-sealed tubes and the hydrolysates were analyzed in a Hitachi type 835 analyzer.

### 2.5. Aminoethylation and carboxymethylation

Samples of 685  $\mu\text{g}$  ( $\sim 348$  nmol) and 550  $\mu\text{g}$  ( $\sim 280$  nmol) of native ST<sub>p</sub> were reduced and then aminoethylated and carboxymethylated, respectively, as in [6]. The recoveries of aminoethylated ST<sub>p</sub> and carboxymethylated ST<sub>p</sub> were 540  $\mu\text{g}$  and 310  $\mu\text{g}$ , respectively.

### 2.6. Enzymatic digestion

Aminoethylated ST<sub>p</sub> (540  $\mu\text{g}$ ,  $\sim 242$  nmol) was dissolved in 0.5% NH<sub>4</sub>HCO<sub>3</sub> (640  $\mu\text{l}$ ) with protease V8 (EC 3.4.21.19) (16  $\mu\text{g}$  in 8  $\mu\text{l}$ ). The mixture was deaerated in a stream of nitrogen and kept at 37°C for 18 h. The digest was lyophilized repeatedly, and then the lyophilizate (405  $\mu\text{g}$ ) was dissolved in 0.1 M pyridine acetate (pH 8.0; 45  $\mu\text{l}$ ) and treated with carboxypeptidase B (EC 3.4.17.2) (8  $\mu\text{g}$ ) at room temperature. Aliquots were withdrawn from the digest at appropriate times for mass measurement.

### 2.7. Edman degradation

Carboxymethylated ST<sub>p</sub> (155  $\mu\text{g}$ ,  $\sim 66.7$  nmol) was degraded manually [11].

### 2.8. FAB mass spectrometry

Mass spectra were recorded with a Jeol double-focusing mass spectrometer JMS-HX100 equipped

with a high-magnetic field and an FAB ion source. The sample (40–50  $\mu\text{g}$ ) was dissolved in 1 M HCl (1  $\mu\text{l}$ ), mixed with glycerol (3  $\mu\text{l}$ ), and applied to a stainless steel sample holder (3  $\times$  3 mm). Typical experimental conditions were as follows: xenon atom beam source, 3 KeV accelerating potential and 200  $\mu\text{A}$  emission current. Ion source, 5 KV accelerating potential.

## 3. RESULTS AND DISCUSSION

### 3.1. Purification

Crude toxin extracted from the culture supernatant of *E. coli* 18D was chromatographed on SP-Sephadex C-50 (fig.1). The main toxic activity was recovered in a peak eluted with 0.1 M AcONH<sub>4</sub> (pH 6.6). Since the peak fraction still contained a large amount of pigmented material, the area shown by a horizontal bar was then subjected to ion-exchange chromatography on DEAE-Sephadex A-25. The toxic activity emerged in the area shown by a horizontal bar in fig.2. Final purification was performed by HPLC on a reversed-phase column, as shown in fig.3. The main toxic fraction shown by a horizontal bar was confirmed to be pure by repeated HPLC (not shown). The purification of ST<sub>p</sub> is summarized in table 1. The amino acid composition was determined to be Asp<sub>2.01</sub> (2), Thr<sub>0.99</sub> (1), Glu<sub>1.16</sub> (1), Pro<sub>1.27</sub>




Fig.1. Chromatography on SP-Sephadex C-50 of crude toxin. Column size: 3  $\times$  46 cm, H<sup>+</sup> form. Elution: An aqueous solution of crude toxin was applied, the column was washed with dist. water and then material was eluted with 0.1 M AcONH<sub>4</sub> (pH 6.7). Fractionation size: 13.5 ml/tube; flow rate: 150 ml/h.




Fig.2. Chromatography on DEAE-Sephadex A-25 of the fraction shown by a horizontal bar in fig.1. Column size:  $1.2 \times 53$  cm, acetate form. Elution: a linear gradient from H<sub>2</sub>O to 0.32 M AcOH and then 1 M HCOOH. Fractionation size: 13.5 ml/tube; flow rate: 75 ml/h.




Fig.3. Reversed-phase HPLC of the toxic fraction obtained by DEAE-Sephadex A-25 chromatography (fig.2). Column: LiChrosorb RP-8 ( $8 \times 300$  mm,  $5 \mu\text{m}$ ). Elution: sample (tube no.54 in fig.2) was applied and material was eluted with 10% CH<sub>3</sub>CN in 0.01 M AcONH<sub>4</sub> (pH 5.7) for 10 min and then with a linear gradient (---) of 10–35% CH<sub>3</sub>CN in 0.01 M AcONH<sub>4</sub> (pH 5.7). Flow rate: 2 ml/min. The toxic activity was found in the peak marked by a horizontal bar.

(1), Gly<sub>1.16</sub> (1), Ala<sub>2.00</sub> (2),  $\frac{1}{2}$ Cys<sub>4.92</sub> (6), Leu<sub>1.13</sub> (1), Tyr<sub>1.92</sub> (2), and Phe<sub>0.99</sub> (1) (values were calculated as mol/mol of Ala; numbers in parentheses indicate nearest integer values). The values were identical to those of the ST in [12]. The minimum effective dose of purified ST<sub>p</sub> in suckling mice assay was 2.5 ng.

Table 1  
Summary of purification of ST<sub>p</sub> produced by ETEC 18D

	Total unit	Spec.act. (ng/MU)	Recovery (%)
Culture supernatant (20 L)	$3.2 \times 10^6$	17800	100
Crude toxin	$2.48 \times 10^6$	224	77.5
SP-Sephadex C-50	$1.89 \times 10^6$	172	59.1
DEAE-Sephadex A-25	$4.05 \times 10^5$		12.7
HPLC			2.5 (1.49 mg)



### 3.2. *M<sub>r</sub>*-value

Purified ST<sub>p</sub> was analyzed by FAB mass spectrometry, a method developed in [13] and the results are shown in fig.4. The observed signal at  $m/z = 1971$  indicated that the *M<sub>r</sub>* of ST<sub>p</sub> was 1970, because free peptides generally give protonated molecular ions ( $[M + H]^+$ , where M denotes the *M<sub>r</sub>*) in positive FAB mass spectra. This *M<sub>r</sub>*-value was compatible with the *M<sub>r</sub>* (1969–1972) calculated from the amino acid composition (Asp + Asn and Glu + Gln were analyzed as Asp and Glu, respectively, in the acid hydrolysate).

### 3.3. Amino acid sequence

Recently, we attempted to synthesize ST with the amino acid sequence reported in [7] and found that the synthetic peptide had a different retention time on HPLC from that of the native toxin isolated from the same strain ETEC 18D as that used in [12]. This finding suggested that the purified native toxin used in our work had a different amino acid sequence from that proposed in [7], although it had the same amino acid composition as that of their proposed sequence.

Some of us [8,9] recently developed a new method for sequence determination of peptide mixtures by a combination of field-desorption (FD) mass spectrometry and exopeptidase digestion. We used this method for determination of the sequence of ST<sub>p</sub>, but with FAB mass spectrometry instead of FD mass spectrometry [6,14]. Fig.5A shows the positive FAB mass spectrum of reduced and aminoethylated ST<sub>p</sub> digested with protease V8, giving intense signals at  $m/z = 965$  and 1289. The

Fig.4. Positive FAB mass spectrum of purified ST<sub>p</sub>.Fig.5. Positive FAB mass spectra of: (A) aminoethylated (AE) ST<sub>p</sub> digested with protease V8; (B) a mixture of samples of (A) treated with carboxypeptidase B for 30 s, 1 and 5 min.

results suggest that the resulting two peptide fragments had  $M_r$  of 964 and 1288. Fig.5B shows the positive FAB mass spectrum of the digest with carboxypeptidase B of the peptide fragments shown in fig.5A. From the observed signals at  $m/z = 980$  and 1126, the C-terminus of the peptide fragment with the mass value 1289 was -Cys(AE)(146)-Tyr(163)-OH (where AE means an aminoethyl group and numbers in parentheses denote residual weights of amino acids). The FD mass spectrum of the carboxypeptidase B digest gave intense signals at  $m/z = 165$  and 182 together with 1126 (not shown), indicating that aminoethyl-

cysteine (165) and tyrosine (182) were released from the peptide fragment (1289).

Reduced and carboxymethylated ST<sub>p</sub> was degraded manually by the Edman method, the results revealed the sequence of 16 amino acid residues from the N-terminus. At the C-terminal two residues were uncertain. The sequence was compatible with the data obtained from the mass spectra shown in fig.4,5A and 5B. The sequence data are summarized and compared with reported sequences for ST in fig.6.

In conclusion, the primary structure of ST<sub>p</sub> was established. As shown in fig.6, the sequence was




Fig.6. Amino acid sequence of ST: (a) present data: — Edman method, — a combination of FAB mass spectrometry and carboxypeptidase B digestion and  $M_r$  of aminoethylated  $ST_p$  digested with protease V8; (b) see [7]; (c) see [3]; (d) see [6].

identical to that deduced from the DNA sequence encoding  $ST_p$  [3], but different from that proposed in [7]. In addition, common and different structural features of  $ST_p$  and  $ST_h$  [6] were established.

#### ACKNOWLEDGEMENT

We express our sincere thanks to Jeol (Tokyo) for facilitating use of the mass instrument JMS-HX100.

#### REFERENCES

[1] Smith, H.W. and Gyles, C.L. (1970) *J. Med. Microbiol.* 3, 387–401.  
 [2] So, M., Atchison, R., Falkow, S., Moseley, S. and McCarthy, B.J. (1980) *Cold Spring Harbor Symposia on Quantitative Biology* 45, 53–58.

[3] So, M. and McCarthy, B.J. (1980) *Proc. Natl. Acad. Sci. USA* 77, 4011–4015.  
 [4] Moseley, S.L., Echeverria, P. and Falkow, S. (1981) in: *The 17th Joint Conference US–Japan Cooperative Medical Science Program, Cholera Panel, Baltimore.*  
 [5] Takeda, Y., Takeda, T., Yano, T., Yamamoto, K. and Miwatani, T. (1979) *Infect. Immun.* 25, 978–985.  
 [6] Aimoto, S., Takao, T., Shimonishi, Y., Hara, S., Takeda, T., Takeda, Y. and Miwatani, T. (1983) *Eur. J. Biochem.*, in press.  
 [7] Chan, S.-K. and Giannella, R.A. (1981) *J. Biol. Chem.* 256, 7744–7746.  
 [8] Shimonishi, Y., Hong, Y.-M., Takao, T., Aimoto, S., Matsuda, H. and Izumi, Y. (1981) *Proc. Japan Acad.* 57B, 304–308.  
 [9] Hong, Y.-M., Takao, T., Aimoto, S. and Shimonishi, Y. (1983) submitted.  
 [10] Mundell, D.H., Anselmo, S.R. and Wishnow, R.M. (1976) *Infect. Immun.* 14, 383–388.  
 [11] Edman, P. and Begg, G. (1967) *Eur. J. Biochem.* 1, 80–91.  
 [12] Staples, S.J., Asher, S.E. and Giannella, R.A. (1980) *J. Biol. Chem.* 255, 4716–4721.  
 [13] Barber, M., Bordori, R.S., Sedgwick, R.D. and Tyler, A.N. (1981) *J. Chem. Soc. Chem. Commun.* 325–327.  
 [14] Bradley, C.V., Williams, D.H. and Hanley, M.R. (1982) *Biochem. Biophys. Res. Commun.* 104, 1223–1230.