

of Diseases (ICD)-9 diagnosis codes 381 or 382. Incremental total expenditures associated with OM were estimated using a regression model adjusting for age, gender, race, ethnicity, education, geographic region, insurance status and number of medications used (proxy for comorbidity). Given the skewed distribution of expenditure variables, multiple model specifications including ordinary least squares regression, generalized linear model (GLM) with Poisson, gamma and negative binomial variance functions were evaluated. **RESULTS:** The prevalence of OM among children in the US was estimated at 11.32%, i.e., 8.47 million persons (95% CI: 10.30% to 12.33%). A majority of children with OM were male (51.7%), white (84.2%), and insured (97.3%) with mean age of 4.7±0.20 years and education of 0.9±0.11 years. In unadjusted analysis among patients with OM, physician office visits comprised the largest proportion of total expenditures at \$769.06 (SE: 54.99), followed by prescription medications at \$606.56 (SE: \$41.72), and inpatient visits at \$500.73 (SE: \$148.42). After controlling for covariates, children with OM had 48% higher total expenditures than those without OM (Estimate: 1.48; p=0.001). The annual adjusted mean incremental total expenditure associated with OM was \$543.10 (SE: \$192.0; p=0.005) per person. **CONCLUSIONS:** Given the prevalence of OM in children and its associated incremental expenditures, the annual direct medical expenditure for treating OM in children is estimated at approximately \$4.6 billion in 2012 US dollars.

PHS19

DIRECT MEDICAL COSTS ASSOCIATED WITH EXACERBATIONS RELATED TO NON-CYSTIC FIBROSIS BRONCHIECTASIS

Joish VN¹, Spilsbury-Cantalupo M², Kamalakar R³, Operschall E⁴, Luong B⁵
¹Bayer HealthCare Pharmaceuticals, Inc., Wayne, NJ, USA, ²Bayer HealthCare Pharmaceuticals, Inc., Wayne, NJ, USA, ³Bayer Pharmaceuticals, Wayne, NJ, USA, ⁴Bayer Schering Pharma, AG, Berlin, Berlin, Germany, ⁵Bayer Pharma AG, Berlin, Berlin, Germany

OBJECTIVES: To estimate direct medical costs in non-cystic fibrosis bronchiectasis (NCB) patients with exacerbations. **METHODS:** An observational, retrospective cohort study was conducted using data from MarketScan claims database. Bronchiectasis patients were identified as those with a medical claim with ICD-9 codes of 494.0 or 494.1 anytime between July 1, 2009 to June 30, 2010. The first bronchiectasis-related claim during this period was defined as the index event. All patients had to have no other bronchiectasis related medical claim 12-months before the index event and no cystic fibrosis related medical claim (ICD-9 code:277.0) 12-months before and after index event. NCB patients were divided into two cohorts. Each NCB patient with exacerbation (NCB-E) was matched to three NCB patients without exacerbations (NCB-W) on geographic region, health plan and gender. Medical resource use and expenditures were estimated for 12-months before and after index event and compared between the two cohorts using non-parametric univariate statistical tests. **RESULTS:** 897 NCB-E patients were matched to 2,691 NCB-W patients. NCB-E patients were older (67 vs. 64, p<0.01), had greater overall comorbidity as estimated by the Charlson's comorbidity score (1.83 vs. 1.66, p=0.01), and respiratory/chest-related symptoms (73% vs. 66%, p<0.001) at baseline compared to NCB-W patients. Total average costs per patient post index-event were significantly greater among NCB-E versus NCB-W patients (\$35,718 vs. \$26,868). After adjusting for baseline health care expenditures, the overall annual average incremental burden due to exacerbations were estimated to be \$7,643. Of this, inpatient hospital costs accounted for \$5,772 or 75% of this burden. Bronchiectasis-related total costs were also significantly (p<0.001) higher in the follow-up period among NCB-E patients compared to NCB-W patients (\$4,609 vs. \$1,210), even after adjusting for baseline consumption. **CONCLUSIONS:** NCB patients with exacerbations have greater medical expenditures than those without. Effective management of exacerbations may be the key in controlling economic burden in this patient population.

PHS20

DIRECT MEDICAL COSTS OF DIABETIC COMPLICATIONS IN THE UNITED STATES

Alvarez P¹, Ward A¹, Chow W², Vo L², Martin S²
¹United BioSource Corporation, Lexington, MA, USA, ²Janssen Scientific Affairs, LLC, Raritan, NJ, USA

OBJECTIVES: To estimate the direct medical costs associated with managing complications of type 2 diabetes in the United States. **METHODS:** A combination of direct data analysis and microcosting was used to estimate costs for an event leading to either a hospital admission or outpatient care, and the post-acute care associated with managing macrovascular, microvascular, and other complications. Data were obtained from many sources, including inpatient and emergency department care databases, national physician and laboratory fee schedules, government reports, and published literature. For each complication, the event year costs derived refer to those associated with the acute episode and, when appropriate, include subsequent ongoing care provided within the first year following the acute episode. Annual costs after the first year for the continued management of complications were also estimated. All costs are reported in 2012 USD. **RESULTS:** Annual costs accrued per patient in the year of the event were estimated for the following macrovascular complications: myocardial infarction \$56,445, ischemic stroke \$42,119, congestive heart failure \$23,758, ischemic heart disease \$21,406, and transient ischemic attack \$7,388. Event year costs per patient for microvascular complications were: chronic kidney disease \$71,714, lower extremity amputations \$9,041, blindness \$2,862, and diabetic foot ulcers \$2,147. Cost estimates of a medically managed hypoglycemic episode were: \$176, \$1,311, and \$16,478 for a mild, moderate, or severe (i.e., requiring hospitalization) episode. **CONCLUSIONS:** This study provides updated and additional estimates of the cost of diabetic complications in 2012 USD, following the methodology used to generate prior cost estimates in 2000 USD referenced in many diabetes economic models (O'Brien et al.

Clin Ther 2003;25(3):1017-1038). Managing complications results in substantial costs to the health care system, as macrovascular complications yield high inpatient care costs in the event year, and long-term management of complications such as chronic kidney disease lead to substantial ongoing annual costs.

PHS21

RENAL CELL CARCINOMA TREATMENT IN SÃO PAULO, BRAZIL: PATTERNS AND COSTS OF HOSPITALIZATION

Fernandes RA¹, Takemoto MMS¹, Takemoto MLS¹, Amaral LM¹, Vasconcellos JF¹, Haas LC¹, Silva MA¹, Fujii RK², Cortes M², Mould J³

¹ANOVA - Knowledge Translation, Rio de Janeiro, RJ, Brazil, ²Pfizer, Inc., São Paulo, Brazil, ³Pfizer, New York, NY, USA

OBJECTIVES: São Paulo is a Brazilian state with the largest number of cancer reference hospitals and hosts 21% of country population. This study aims to describe hospitalization patterns and costs for renal cell carcinoma (RCC) in public hospitals in São Paulo state, Brazil. **METHODS:** A retrospective analysis of São Paulo public hospital admissions for RCC was developed according to ICD-10 classification (C64) as reported in Brazilian Hospital Information System (SIH/DATASUS) database from January 2007 to December 2011. Admissions were categorized into surgical (containing any surgical procedures codes at hospitalization data eg: tumor excision, biopsy, etc.) or clinical (all non-surgical eg: chemotherapy, infection treatment, anemia, etc.). Costs represent federal reimbursement values for hospitalizations (includes medical procedures, exams, medications and taxes), presented in 2012 Brazilian Real (BRL). Average 2012 exchange rate was 1,9544BRL/USD. **RESULTS:** In the period, 4,138 hospital admissions were identified (951, 1,126, 1,207, and 854 for 2007, 2008, 2009, and 2011, respectively), and total cost was 5,472,534BRL, which represents 0.78% of all cancer in-hospital treatment in the state. Mean cost per patient was 1,326BRL, 675BRL, and 1,854BRL for all hospitalizations, clinical and surgical procedures, respectively. Surgical procedures were responsible for 55.07% of cases, and 77.29% of total costs, with partial or total nephrectomy representing the main cause (48.9%). Mean length of stay was 6.99 days. In-hospital mortality rate was 9.98% for all cause (17.7% and 3.69% for clinical and surgical procedures, respectively). When compared to the national cases, São Paulo represents 29.74% of them, while representing 33.18% of total national hospitalization costs related to RCC. **CONCLUSIONS:** RCC hospitalizations costs in São Paulo are responsible for less than 1% of all cancer in-hospital treatments in the state, which may suggest under-diagnosing when compared to international incidence rates. São Paulo is an important RCC treatment center in Brazil, capturing about one third of country resources.

PHS22

COSTS AND HEALTH CARE RESOURCE UTILIZATION ASSOCIATED WITH ASTHMA IN CANADA: A SYSTEMATIC REVIEW

Ismaila A¹, Sayani A¹, Marin M², Tacon C¹, Crater G¹, Su Z¹

¹GlaxoSmithKline, Mississauga, ON, Canada, ²Optuminsight, Life Sciences, Burlington, ON, Canada

OBJECTIVES: To determine the cost and health care resource utilization associated with asthma management in Canada. **METHODS:** A systematic literature search of PUBMED, EMBASE and EMCare electronic databases using relevant search strategy was conducted from 2000 until 2011 to identify articles, published in English, that reported the data on the clinical and economic burden of asthma in Canada. Two investigators independently screened citations, reviewed studies, assessed study quality and abstracted data. Heterogeneity in study design, data sources and asthma definition precluded meta-analysis of study outcomes. All costs were reported in Canadian dollars and inflated to 2011. **RESULTS:** Of the 570 citations identified, 67 full-text articles were reviewed, of which 33 studies met inclusion criteria. Both children and adults with asthma averaged less than one emergency department (ED) visit per patient per year. In children, frequency of visits was largely influenced by the presence of prior asthma exacerbations. In adults, increased ED visits were associated with inappropriate use of asthma medication. ED visit rates were significantly higher in women than in men and, increased with age. Hospitalization rates in children decreased with age and patients with persistent asthma were more than one and a half times admitted more often than those with non-symptomatic asthma. Average hospitalization costs per acute asthma episode ranged from \$458 to \$923 and the average costs for ED visit per acute asthma episode ranged from \$219.08 to \$276.76, excluding medication cost. The average annual cost of asthma medication was \$230.58 per patient. Annual productivity loss days in adult asthma patients varied from 12 in employed persons to 20 in students and 49 in disability pensioners. **CONCLUSIONS:** The available evidence shows that the asthma-related costs, both direct and indirect remain substantial. The overall burden varies based on whether studies reported costs from the perspective of an individual asthmatic or from the population level.

PHS23

MELANOMA HEALTH CARE PATHWAYS IN AN ITALIAN POPULATION

Ascierto P¹, Bernardi D², De Rosa M³, Martini N⁴, Rielli R⁵, Rossi F⁶

¹Istituto Nazionale Tumori Fondazione "G. Pascale", Napoli, Italy, ²CINECA, Casalecchio di Reno - Bologna, Italy, ³CINECA Interuniversity Consortium, Casalecchio di Reno, BO, Italy, ⁴National Academy of Medicine, Roma, Italy, ⁵CINECA, Casalecchio di Reno (BO), Italy

OBJECTIVES: To describe treatment patterns, direct costs and comorbidities in a population of patients with melanoma. **METHODS:** Starting from ARNO Observatory, an Italian population-based patient-centric system through record linkage of 11 million inhabitants, we identified a cohort of patients with melanoma (ICD-9 code: 172 and/or V10.82) during 2009. Discharge records, reimbursed prescription patterns, lab tests and diagnostic examinations were