

**BRIEF REPORT**

Registration of Human Embryonic Stem Cell Lines: Korea, 2010

Ji-Yoon Lee, Dae-Yeon Lee, Young-Sil Choi, Kyoung-Jae Lee, Yong-Ou Kim*

Division of Life Science Research Management, Korea National Institute of Health, Osong, Korea.

Received: April 20, 2011

Revised: May 5, 2011

Accepted: May 10, 2011

KEYWORDS:

embryo,
human embryonic stem
cell (hESC),
registry

Abstract

In an effort to increase the credibility of human embryonic stem cell (hESC) lines established in Korea, obligatory registration was introduced by the Bioethics and Safety Act 2008, effective as of January 1, 2010.

The DNA fingerprint, chromosome stability, expression of pluripotency markers, and contamination of mycoplasma of the submitted lines were analyzed by Korea Centers for Disease Control and Prevention (KCDC). The characterization data and ethical aspects, such as informed consent for donation of surplus embryos, were reviewed by a 10-member advisory review committee for stem cell registry. A total of 55 domestic hESC lines were submitted for registration in 2010; among them 51 were registered. Among these submitted lines, 26 were additionally characterized by KCDC, while 25 lines previously characterized by the Ministry of Education, Science and Technology were not additionally analyzed by KCDC.

Registration completed an oversight system for embryo research by registering the products of licensed embryo research projects, making embryo research more transparent in Korea. Information about hESC lines is available at the website of the Korea Stem Cell Registry (kscri.nih.go.kr).

1. Introduction

A recent clinical trial on a stem cell therapy was initiated using derivatives from embryonic stem cells (ESCs) [1]. Human ESCs (hESCs) have a wide variety of possibilities in regenerative medicine, because of their two advantageous characteristics that distinguish them from adult stem cells: (1) embryonic stem cells can differentiate into any type of cell or tissue; and (2)

embryonic stem cells have the capacity to proliferate indefinitely under optimal conditions. However, there are several obstacles in the use of hESCs that need to be overcome before they can be used in regenerative medicine. In addition to the scientific aspects, such as accurate control of proliferation and purity of differentiated cells, the reliability of the cell line is an ethical issue. This is because hESCs originate from human embryos. As the need for comprehensive information

*Corresponding author.

E-mail: youkim@nih.go.kr

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

about the characteristics of hESC lines and provenance increased, several countries and groups established hESC registries [2,3]. The European registry, hESCreg (<http://www.hescreg.eu/>), was founded as a collaborative and interdisciplinary platform where researchers and the general public can access information about available hESC lines. The international stem cell registry at the University of Massachusetts Medical School (<http://www.umassmed.edu/iscr/>) has been continuously expanding its database since its launch in 2008, and currently has information on >400 hESCs and 120 induced pluripotent stem cells (iPSCs). In accordance with an order from the President of the United States. The National Institute of Health (NIH) has registered 91 hESC lines eligible for federal funding since 2009 (<http://stemcells.nih.gov/research/registry>). These registries provide information through their websites with features relevant to their own purposes.

Several scientists have been active in the production of hESC lines in Korea [4,5]. Approximately 60 hESC lines have been known to be established in Korea. Although the Bioethics and Safety Act has been effective since 2005, the exact information about hESC lines was not available. To improve the credibility of hESC lines that have been established and used in Korea, an obligatory registration was initiated on January 1, 2010 according to the Bioethics and Safety Act 2008. Here, we report the first 1-year results of the registration.

2. Materials and Methods

2.1. Cell culture

The hESC line H9 was cultured on STO (ATCC Manassas, VA, USA) feeder in 80% DMEM/F12 medium (Gibco, Carlsbad, CA, USA) supplemented with 20% KO serum (Gibco), 0.1 mM nonessential amino acids (Gibco), 0.1 mM β -mercaptoethanol (Gibco), and 4 ng/mL bFGF (Invitrogen). The STO feeder cells were maintained in DMEM medium containing 10% FBS (Gibco), and used after mitotic inactivation by mitomycin C (Sigma, St. Louis, MO, USA). H9 cells were passaged every 5–7 days with 2 μ L/mL ROCK inhibitor (Sigma), and the culture medium was changed daily. The submitted hESC lines were minimally cultured in their own media. The human fibroblast cell line IMR-90 (ATCC) was maintained in DMEM medium containing 10% FBS and 1% penicillin/streptomycin (Gibco).

2.2. DNA fingerprinting

The identity of the hESC line was analyzed by short tandem repeat (STR) analysis. The hESC colonies were removed from the surface of the culture dish and collected in a 15-ml conical tube. The STR analyses were performed by DowGene (Seoul, Korea) and Kogene (Seoul, Korea); both companies used a Powerplex 16 system (Promega). The STR data of the

submitted hESC lines were compared with those of other lines with clustering software R and Microsoft Excel.

2.3. Karyotyping

Karyotyping of hESCs was conducted by GenDix (Seoul, Korea) and Samkwang (Seoul, Korea). For karyotyping, a 4-well dish of each cell line was analyzed by GTG-banding method.

2.4. RT-PCR

Total RNA was isolated using Trizol (Invitrogen) according to the manufacturer's protocol. Subsequently, cDNA was synthesized from 1 μ g of the total RNA by PrimeScript 1st strand cDNA synthesis kit (TAKARA). Quantitative real-time PCR (qRT-PCR) was performed with TaqMan assays (Applied Biosystems) and Gene Expression probes (Applied Biosystems; GAPDH, Hs99999905_m1; NANOG, Hs02387400_g1; OCT4, Hs00742896_s1; SOX2, Hs00602736_s1; TERT, Hs00162669_m1). qRT-PCR was performed using TaqMan Universal PCR Master Mix and ABI 7500 Real-Time PCR System (Applied Biosystems). Expression levels were analyzed by $\Delta\Delta C_t$ method. All reactions were duplicated.

2.5. Mycoplasma tests

Two methods were used to estimate mycoplasma contamination in the hESCs. Tests were performed immediately after receiving cells from applicants. For PCR detection, PCR Mycoplasma Detection Set (TAKARA) was used with 0.5 μ L ES culture media. For enzymatic detection, MycoAlert Mycoplasma Detection Kit (Lonza, Basel, Switzerland) was used with 100 μ L ES culture media. Fresh ES culture media and positive control contained in the kit were used as controls. The amplified PCR products were analyzed by gel electrophoresis with 1% agarose gels. Enzymatic reactions were analyzed by Sirius L tube Luminometer (Berthold, Pforzheim, Germany).

3. Results

3.1. Procedure for registration

The registration procedure is shown in Figure 1. A stem cell scientist submitted application forms with copies of informed consent for use of surplus embryos. After a pre-review of the submitted files by the advisory review committee, we requested and analyzed the hESC lines. We analyzed DNA fingerprints, karyotypes, pluripotency markers, and mycoplasma contamination. The submitted files and our analyzed data were reviewed by the advisory review committee, which consisted of seven stem cell scientists, one bioethics professor, and two government employees. We had four committee




Figure 1. Registration procedure for hESC lines in Korea. KCDC runs the registry and the advisory committee reviews the submitted data for the registry.

meetings and two document reviews for the registration in 2010.

3.2. Informed consent

The Bioethics and Safety Act has been in effect since 2005. According to this act, only surplus embryos should be used for research. Therefore according to the provisions of this act, fertility clinics need to obtain informed consent from patients before the surplus embryos can be used for research. The registration indicated that 29 hESC lines were established from 917 surplus frozen embryos donated from 151 patients since 2005. It is now confirmed that all hESC lines were established from surplus embryos that have been provided by patients, with their informed consent, since 2005.

3.3. Analyses of hESC lines

Among 51 registered hESC lines, we analyzed the DNA fingerprint, karyotype, expression of pluripotency markers, and mycoplasma contamination of 27 lines. We harvested and analyzed hESCs as soon as the cells were submitted, but the cells were maintained for 1–2 days if the amount of cells available was insufficient for analyses. Another 25 lines were verified by the Ministry of Education, Science and Technology (MOEST) and were not additionally analyzed by us. A summary of the verified data is shown in Table. DNA fingerprinting using short tandem repeat (STR) analyses showed that all except analyzed lines except for one line are independent lines (Table). Among the 27 lines analyzed by us, three lines showed abnormal karyotypes (Figure 2A–D). Expression of pluripotency marker genes was detected in all the tested lines, although the expression levels of several pluripotency markers of an aneuploid hESC line (CHA-hES 16) were relatively lower than those of other hESC lines (Figure 2E). Mycoplasma contamination tests were performed twice if the first analysis yielded positive results. Eight lines were mycoplasma-positive in the first analysis and were

resubmitted. Two lines were still mycoplasma-positive in the second analysis of submitted cells, while the other lines were negative in the second analysis of submitted cells, as shown in Table.

4. Discussion

The Korea hESC registry was founded to ensure that hESC research in Korea is ethically responsible and conducted in accordance with the Bioethics and Safety Act by providing reliable information about hESC lines. Our registry is enforced by the Bioethics and Safety Act 2008; under this act, registration of hESC lines is obligatory in Korea. The obligatory registration system is feasible in Korea because researchers who want to use human surplus embryos are required to obtain a license from the MOHW, and the hESC line is thus a product of licensed research. The Bioethics and Safety Act has been in effect since 2005, and obligatory registration started in 2010.

As a result of the registration, we confirmed that 29 hESC lines established after 2005 were from 917 surplus embryos obtained with informed consent. Among the 55 submitted lines, two lines could not be registered, and two other lines are under review. Dr. Hwang's Soom-hES1 (also known as NT-1) line was ineligible for registration because of the unethical aspects revealed by the report of the investigation committee of Seoul National University [6]. In addition, NT-1 was revealed to be a product of parthenogenesis [6,7], which is not allowed under the Bioethics and Safety Act. Another hESC line that could not be registered is AMC-1, which showed the same DNA fingerprint as a previously established hESC line (Miz-hES4). When we requested the researchers to submit early passage of AMC-1, we were informed that it failed to grow. Among the 51 registered lines, 10 lines have abnormal karyotypes, as shown in Table. The hESC lines with abnormal karyotypes might be used as models of aneuploid chromosomal syndrome [8–10].

Table. Summary of analyses of hESC lines submitted in 2010

a	Names of hESC lines ^b	Year	Submitted data							KCDC-analyzed data						Registered
			Karyotype	STR	Immuno- staining	RT- PCR	Mycoplasma	EB or teratoma	Copies of informed consents	List of surplus embryos	Karyotype	STR	Immuno- staining	RT- PCR	Mycoplasma	
1	SNUhES1	2001	46, XY or 47, XY,+12	√	+	+	—	Teratoma				√ ^c				Yes
2	SNUhES2	2001	46, XX	√	+	+	—	Teratoma				√ ^c				Yes
3	SNUhES3	2001	46, XY	√	+	+	—	Teratoma				√ ^c	+ ^c	+ ^c		Yes
4	SNUhES4	2003	46, XY	√	+	+	—	Teratoma				√ ^c	+ ^c	+ ^c		Yes
5	SNUhES5	2004	47, XY,+16	√	+	+	—	Teratoma				√ ^c				Yes
6	SNUhES6	2004	46, XY	√	+	+	—	EB				√ ^c				Yes
7	SNUhES7	2004	46, XY	√	+	+	—	Teratoma				√ ^c				Yes
8	SNUhES10	2004	69, XXX	√	+	+	—	EB				√ ^c				Yes
9	SNUhES11	2004	46, XY	√	+	+	—	Teratoma				√ ^c	+ ^c	+ ^c		Yes
10	SNUhES12	2004	46, XX	√	+	+	—	EB				√ ^c				Yes
11	SNUhES14	2004	46, XY	√	+	+	—	EB				√ ^c				Yes
12	SNUhES16	2004	46, XY	√	+	+	—	Teratoma				√ ^c	+ ^c	+ ^c		Yes
13	SNUhES18	2004	46, XY or 47, XY,+12	√	+	+	—	Teratoma				√ ^c				Yes
14	SNUhES19	2004	46, XY or 47, XY,+12	√	+	+	—	Teratoma				√ ^c				Yes
15	CHA-hES 3	2005	46, XY	√	+	+	—	Teratoma	√	√		√ ^c	+ ^c	+ ^c		Yes
16	CHA-hES 4	2004	46, XY	√	+	+	—	Teratoma	√	√		√ ^c	+ ^c	+ ^c		Yes
17	CHA-hES 5	2004	46, XY, inv(9)	√	+	+	—	Teratoma	√	√		√ ^c				Yes
18	CHA-hES 6	2004	46, XX	√	+	+	—	Teratoma	√	√		√ ^c				Yes
19	CHA-hES 7	2004	46, XX	√	+	+	—	Teratoma	√	√		√ ^c				Yes
20	CHA-hES 8	2004	46, XX	√	+	+	—	Teratoma	√	√		√ ^c				Yes
21	CHA-hES 9	2004	46, XX	√	+	+	—	Teratoma	√	√		√ ^c	+ ^c	+ ^c		Yes
22	CHA-hES 10	2005	46, XY	√	+	+	—	Teratoma	√	√		√ ^c				Yes
23	CHA-hES 11	2005	46, XY	√	+	+	—	Teratoma	√	√		√ ^c				Yes
24	CHA-hES 12	2005	46, XY	√	+	+	—	Teratoma	√	√		√ ^c				Yes
25	Miz-hES4	2003	46, XY	√	+	+		EB				√ ^c	+ ^c	+ ^c		Yes
26	CHA-hES 13	2007	46, XX	√	+	+	—	Teratoma	√	√	46, XX	√		+	— ^d	Yes
27	CHA-hES 14	2007	46, XY	√	+	+	—	Teratoma	√	√	46, XY	√		+	— ^d	Yes
28	CHA-hES 15	2007	46, XY	√	+	+	—	Teratoma	√	√	46, XY	√		+	—	Yes
29	CHA-hES 16	2007	71, XXY	√	+	+	—		√	√	70, XXY,+12	√		+	— ^d	Yes
30	CHA-hES 17	2007	46, XY	√	+	+	—	Teratoma	√	√	46, XY	√		+	— ^d	Yes
31	CHA-hES 19	2007	47, XX, der(2), -4,+5,+mar	√	+	+	—		√	√	47, XX, t(2;4) q(11.2;p15.2),+5	√		+	— ^d	Yes
32	CHA-hES 20	2007	46, XY	√	+	+	—	Teratoma	√	√	46, XY ^e	√		+	—	Yes
33	CHA-hES 21	2008	46, XX	√	+	+	—	Teratoma	√	√	46, XX	√		+	—	Yes

34	CHA-hES 23	2008	46, XX	✓	+		—	Teratoma	✓	✓	46, XX	✓	+	—	Yes
35	CHA-hES 24	2008	46, XY	✓	+	+	—	Teratoma	✓	✓	46, XY	✓	+	—	Yes
36	CHA-hES 26	2008	Tri- or tetraploidy	✓	+		—		✓	✓	90, XXXX,-6,-18,add(20)	✓	+	—	Yes
37	SNUhES31	2009	46, XY	✓	+	+		EB	✓	✓	46, XY	✓	+	— ^d	Yes
38	CHA-hES 18	2007	46, XX	✓	+	+	—	Teratoma	✓	✓	46, XX	✓	+	—	Yes
39	CHA-hES 22	2008	46, XX	✓	+	+	—	Teratoma	✓	✓	46, XX	✓	+	—	Yes
40	CHA-hES 25	2008	46, XX	✓	+		—	Teratoma	✓	✓	46, XX	✓	+	—	Yes
41	CHA-hES M1	2008	46, XX	✓	+	+	—	Teratoma	✓	✓	46, XX	✓	+	—	Yes
42	CHA-hES R1	2008	46, XY	✓	+	+	—	Teratoma	✓	✓	46, XY	✓	+	—	Yes
43	CHA-hES R2	2008	46, XX	✓	+	+	—	Teratoma	✓	✓	46, XX	✓	+	—	Yes
44	CHA-hES R3	2008	46, XX	✓	+	+	—	Teratoma	✓	✓	46, XX	✓	+	—	Yes
45	CHA-hES R4	2008	46, XY	✓	+	+	—	Teratoma	✓	✓	46, XY	✓	+	—	yes
46	CHA-hES B1	2009	46, XY	✓	+	+	—	Teratoma	✓	✓	46, XY	✓	+	—	Yes
47	CHA-hES B2	2009	46, XX	✓	+	+	—	Teratoma	✓	✓	46, XX	✓	+	—	Yes
48	CHA-hES B3	2009	46, XY	✓	+	+	—	Teratoma	✓	✓	46, XY	✓	+	—	Yes
49	AMC-1	2006	46, XY		+	+		Teratoma	✓	✓	46, XY	✓ ^f	+	—	No
50	JNU-hES-01	2009	46, XY	✓	+	+		Dif ^h	✓	✓	46, XY	✓	+	+ ^g	Yes
51	JNU-hES-02	2009	47, XY,+12	✓	+	+		Dif ^h	✓	✓	46, XY	✓	+	+ ^g	Yes
52	MB01	2002	46, XY		+	+									Under review
53	MB06	2004	46, XY		+	+									Under review
54	Sooam-hES1	2003	46, XX	✓	+	+	—	Teratoma							No
55	SNUhES32	2010	46, XY	✓	+	+	—	EB	✓	✓	46, XY	✓	+	—	Yes

^aIs the submitted number in the order of the review; ^bRegistered lines are indicated with bold; ^cAnalyzed by MOEST; ^dMycoplasma-positive in analysis of first submitted cells, but negative in analysis of second submitted cells;

^eAbnormal karyotype in analysis of first submitted cells, but normal in analysis of second submitted cells; ^fSame STR with Miz-hES4; ^gMycoplasma-positive in analyses of first and second submitted cells; ^hDifferentiation data.




Figure 2. Characteristics of three hESC lines with abnormal karyotypes. (A) Normal karyotype of H9. (B) Abnormal karyotype (70, XXY,+12) of CHA-hES 16. (C) Abnormal karyotype (47, XX, t(2;4)(q11.2;p15.2),+5) of CHA-hES 19. (D) Abnormal karyotype (90, XXXX,-6,-18,add(20)) of CHA-hES 26. (E) The relative quantities of pluripotency marker genes were detected by real-time PCR using Taqman probe. CHA-hES 16 showed lower levels of expression of several pluripotency markers, whereas other abnormal lines showed normal expression patterns. H9 was used as a positive control; mock and human fibroblasts were used as negative controls.

Recently, methods for establishing an iPSC line were developed [11,12]. iPSCs have characteristics similar to hESCs with respect to their pluripotency and self-renewal capacity. Human iPSCs have been recognized as a good model system for genetic diseases and drug screening [13]. We are considering the registration of human iPSC lines to provide information to the scientific community using the hESC registration system. The registration of human iPSC lines will be optional, not obligatory. We hope that this registry (ksr.nih.go.kr) promotes transparency of stem cell research in Korea.

Acknowledgements

We are grateful to SungBum Cho of the Division of Biomedical Informatics for his help with STR data analysis. We would also like to thank other members of the Division of Life Science Research Management for their assistance in running the stem cell registry.

References

1. Strauss S. Geron trial resumes, but standards for stem cell trials remain elusive. *Nat Biotechnol* 2010 Oct;28(10):989–90.
2. Luong MX, Smith KP, Stein GS. Human embryonic stem cell registries: value, challenges and opportunities. *J Cell Biochem* 2008 Oct;105(3):625–32.
3. Borstlap J, Luong MX, Rooke HM, et al. International stem cell registries. *In Vitro Cell Dev Biol Anim* 2010 Apr;46(3–4):242–6.
4. Oh SK, Kim HS, Ahn HJ, et al. Derivation and characterization of new human embryonic stem cell lines: SNUhES1, SNUhES2, and SNUhES3. *Stem Cells* 2005 Feb;23(2):211–9.
5. Lee JE, Kang MS, Park MH, et al. Evaluation of 28 human embryonic stem cell lines for use as unrelated donors in stem cell therapy: implications of HLA and ABO genotypes. *Cell Transplant* 2010;19(11):1382–95.
6. Chung MH. Investigation committee report. Seoul National University; 2006.
7. Kim K, Ng K, Rugg-Gunn PJ, et al. Recombination signatures distinguish embryonic stem cells derived by parthenogenesis and somatic cell nuclear transfer. *Cell Stem Cell* 2007 Sep 13;1(3):346–52.
8. Biancotti JC, Narwani K, Buehler N, et al. Human embryonic stem cells as models for aneuploid chromosomal syndromes. *Stem Cells* 2010 Sep;28(9):1530–40.
9. Yang S, Lin G, Tan YQ, et al. Differences between karyotypically normal and abnormal human embryonic stem cells. *Cell Prolif* 2010 Jun;43(3):195–206.
10. Urbach A, Benvenisty N. Studying early lethality of 45, XO (Turner's syndrome) embryos using human embryonic stem cells. *PLoS ONE* 2009;4(1):e4175.
11. Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 2006 Aug;126(4):663–76.
12. Yu J, Vodyanik MA, Smuga-Otto K, et al. Induced pluripotent stem cell lines derived from human somatic cells. *Science* 2007 Dec;318(5858):1917–20.
13. Park IH, Arora N, Huo H, et al. Disease-specific induced pluripotent stem cells. *Cell* 2008 Sep;134(5):877–86.