



Contents lists available at ScienceDirect

## Journal of Mathematical Analysis and Applications

[www.elsevier.com/locate/jmaa](http://www.elsevier.com/locate/jmaa)

## Homogenization of low-cost control problems on perforated domains

T. Muthukumar\*, A.K. Nandakumaran

Department of Mathematics, Indian Institute of Science, Bangalore-560012, India

## ARTICLE INFO

## Article history:

Received 16 August 2007

Available online 27 September 2008

Submitted by D.L. Russell

## Keywords:

Homogenization

Two-scale convergence

Optimal control

Perforated domain

Low-cost control

## ABSTRACT

The aim of this paper is to study the asymptotic behaviour of some low-cost control problems in periodically perforated domains with Neumann condition on the boundary of the holes. The optimal control problems considered here are governed by a second order elliptic boundary value problem with oscillating coefficients. It is assumed that the cost of the control is of the same order as that describing the oscillations of the coefficients. The asymptotic analysis of small cost problem is more delicate and need the  $H$ -convergence result for weak data. In this connection, an  $H$ -convergence result for weak data under some hypotheses is also proved.

© 2008 Elsevier Inc. All rights reserved.

## 1. Introduction

Let  $n \geq 2$ ,  $\Omega$  be a bounded open set in  $\mathbb{R}^n$  and  $Y = (0, 1)^n$  be the reference cell. Let  $S$ , the hole, be an open subset of  $Y$  and  $\bar{S}$  denote the closure of  $S$  in  $\mathbb{R}^n$ . Let  $Y' = Y \setminus \bar{S}$  be the material part. If  $S + \mathbb{Z}^n = \{x + k \mid x \in S \text{ and } k \in \mathbb{Z}^n\}$ , then, for a parameter  $\varepsilon > 0$  tending to zero, we set  $S_\varepsilon = \varepsilon(S + \mathbb{Z}^n)$ . Note that the open set  $\mathbb{R}^n \setminus \bar{S}_\varepsilon$  is periodically perforated with the reference periodic cell  $\varepsilon Y'$ . We define a periodically perforated domain  $\Omega_\varepsilon$  as,

$$\Omega_\varepsilon = \Omega \cap (\mathbb{R}^n \setminus \bar{S}_\varepsilon).$$

Let  $\chi$  be the characteristic function of  $Y'$  in  $Y$ , i.e.,

$$\chi(y) = \begin{cases} 1 & \text{if } y \in Y', \\ 0 & \text{if } y \in S. \end{cases}$$

We extend  $\chi$  periodically to whole of  $\mathbb{R}^n$  and shall denote this extension also by  $\chi$ . Observe that

$$\Omega_\varepsilon = \left\{ x \in \Omega \mid \chi\left(\frac{x}{\varepsilon}\right) = 1 \right\}.$$

Let  $\chi_0$  denote the ratio of the measure of  $Y'$  to the measure of  $Y$ , i.e.,

$$\chi_0 = \frac{|Y'|}{|Y|} = |Y'|.$$

Assume that the constant  $\chi_0$  is strictly positive. The holes  $S_\varepsilon$  may either be isolated or connected, and they may also meet the boundary  $\partial\Omega$ . We also assume that the open set  $\mathbb{R}^n \setminus \bar{S}_\varepsilon$  is connected and that the boundaries are all of Lipschitz type.

\* Corresponding author.

E-mail addresses: [tmk@math.iisc.ernet.in](mailto:tmk@math.iisc.ernet.in) (T. Muthukumar), [nands@math.iisc.ernet.in](mailto:nands@math.iisc.ernet.in) (A.K. Nandakumaran).

We remark that the hypothesis on the holes  $S_\varepsilon$  and the perforated domain  $\Omega_\varepsilon$  are similar to that of [2], and are stated here only for completeness sake.

We shall denote the norm in  $L^2(\Omega)$  by  $\|\cdot\|_2$  and the norm in  $L^2(\Omega_\varepsilon)$  by  $\|\cdot\|_{2,\Omega_\varepsilon}$ . For a function  $g$  defined on  $\Omega_\varepsilon$ , we shall denote by  $\tilde{g}$  its extension by zero on the holes  $\Omega \cap S_\varepsilon$ . The symbol  $C$  will always denote a generic positive constant independent of  $\varepsilon$ .

Let  $a, b, c$  and  $d$  be given constants such that  $0 < a \leq b$  and  $0 < c \leq d$ . Let  $A = A(x, y)$  be an  $n \times n$  matrix,  $Y$ -periodic in  $y$ , such that,

$$a|\xi|^2 \leq A(x, y)\xi \cdot \xi \leq b|\xi|^2 \quad \text{a.e. in } x \text{ and } y, \quad \forall \xi = (\xi_i) \in \mathbb{R}^n.$$

The above assumptions on  $A$  implies that the matrix is in  $[L^\infty(\Omega \times Y)]^{n \times n}$ . We shall, in addition, assume that  $A$  is in  $C[\Omega; L^\infty_{\text{per}}(Y)]^{n \times n}$ , i.e., the entries of the matrix are continuous functions with respect to  $x \in \Omega$  and with their values in the Banach space of essentially bounded measurable functions,  $Y$ -periodic in  $y$ . We remark that the results of this article remain valid if  $C[\Omega; L^\infty_{\text{per}}(Y)]^{n \times n}$  is replaced by either  $L^\infty[\Omega; C_{\text{per}}(Y)]^{n \times n}$  or  $L^\infty_{\text{per}}[Y; C(\bar{\Omega})]^{n \times n}$ . The weakest possible assumption on  $A(x, y) = (a_{ij}(x, y))$  is that  $a_{ij}(x, \frac{x}{\varepsilon})$  is measurable and satisfies

$$\int_{\Omega} \left| a_{ij} \left( x, \frac{x}{\varepsilon} \right) \right|^2 dx \rightarrow \int_{\Omega} \int_Y |a_{ij}(x, y)|^2 dy dx.$$

Let  $B = B(x, y)$  be a matrix with entries from  $L^\infty(\Omega \times Y)$ ,  $Y$ -periodic in  $y$  satisfying

$$c|\xi|^2 \leq B(x, y)\xi \cdot \xi \leq d|\xi|^2 \quad \text{a.e. in } x \text{ and } y, \quad \forall \xi = (\xi_i) \in \mathbb{R}^n.$$

Also,  $B$  satisfies the other hypotheses as that of  $A$ . In addition, we assume that  $B$  is symmetric. This assumption will not play any role in the homogenization process and is inherited from the optimal control problem.

Let  $U_\varepsilon$  be a closed convex subset of  $L^2(\Omega_\varepsilon)$ , called the admissible control set and  $f \in L^2(\Omega)$ . Given  $\theta_\varepsilon \in U_\varepsilon$ , let the cost functional  $J_\varepsilon$  be defined as,

$$J_\varepsilon(\theta_\varepsilon) = F_\varepsilon(u_\varepsilon, \theta_\varepsilon) + \frac{\varepsilon}{2} \|\theta_\varepsilon\|_{2,\Omega_\varepsilon}^2 \quad (1.1)$$

where  $u_\varepsilon = u_\varepsilon(\theta_\varepsilon) \in H^1(\Omega_\varepsilon)$  is the unique solution of

$$\begin{cases} -\operatorname{div} \left( A \left( x, \frac{x}{\varepsilon} \right) \nabla u_\varepsilon \right) = f + \theta_\varepsilon & \text{in } \Omega_\varepsilon, \\ A \left( x, \frac{x}{\varepsilon} \right) \nabla u_\varepsilon \cdot \nu = 0 & \text{on } \partial\Omega_\varepsilon \setminus \partial\Omega, \\ u_\varepsilon = 0 & \text{on } \partial\Omega \cap \partial\Omega_\varepsilon, \end{cases} \quad (1.2)$$

and  $\nu$  is the unit outward normal on  $\partial\Omega_\varepsilon \setminus \partial\Omega$ .

The aim of this article is to study the limiting behaviour of the system (1.1)–(1.2) in the following two situations:

(a) When  $F_\varepsilon(u_\varepsilon, \theta_\varepsilon)$  is a Dirichlet-type integral, i.e.,

$$F_\varepsilon(u_\varepsilon, \theta_\varepsilon) = \frac{1}{2} \int_{\Omega_\varepsilon} B \left( x, \frac{x}{\varepsilon} \right) \nabla u_\varepsilon(\theta_\varepsilon) \cdot \nabla u_\varepsilon(\theta_\varepsilon) dx. \quad (1.3)$$

(b) When  $F_\varepsilon(u_\varepsilon, \theta_\varepsilon)$  is the  $L^2$ -norm of the state, i.e.,

$$F_\varepsilon(u_\varepsilon) = \frac{1}{2} \|u_\varepsilon(\theta_\varepsilon)\|_{2,\Omega_\varepsilon}^2. \quad (1.4)$$

It is a classical result from the calculus of variations that there exists a unique  $\theta_\varepsilon^* \in U_\varepsilon$  such that

$$J_\varepsilon(\theta_\varepsilon^*) = \min_{\theta_\varepsilon \in U_\varepsilon} J_\varepsilon(\theta_\varepsilon). \quad (1.5)$$

The main results of this paper are: Theorem 2.1, Theorem 3.3, Theorem 3.7 and Theorem 4.4. In this article, we study the situation where the Neumann condition is imposed on the boundary of the holes and the Dirichlet condition on the exterior boundary (cf. (1.2)). In fact, the results will remain valid (locally inside  $\Omega$ ) when the Neumann condition is imposed both on the boundary of holes and on the exterior boundary (cf. [2]). However, the problem with Dirichlet condition on the holes is more delicate and is the topic of a future article.

The  $\varepsilon$  factor appearing in the second term of the right-hand side of the cost functional (1.1) is, usually, in literature called the *cost of the control*. The fact that this is of  $\varepsilon$  order motivates the name *low-cost control*. The low-cost control problems

were first introduced by J. L. Lions (who called it ‘cheap’ control) in [14,15]. Several variants of low-cost control problems have been studied by Kesavan and Saint Jean Paulin in [12], for non-perforated domains. The study is taken up further in [8,19] for both perforated and non-perforated domains. In the above articles, the problems are considered in the general (non-periodic) case using the method of oscillating test functions, introduced by L. Tartar (cf. [17,18,23]). In spite of these studies, the problem has not been completely settled in its full generality. One major drawback is the lack of information on the convergence of the optimal controls itself, except in some trivial cases (cf. [19]). The problem has only been settled for particular cases of admissible control set.

In [8,19], the state-adjoint system governing the optimal control problem was homogenized for the non-perforated domain with the assumption that the control set was the set of all positive functions (positive cone) of the control space. The perforated case was unsettled, even for the positive cone of the control space. In this paper, using the notion of two-scale convergence (cf. [1,16,20,21]), we show that the state-adjoint system governing the optimal control problem can be homogenized *without any assumptions on the control set* for both perforated and non-perforated domains.

However, we would like to note that the positive cone hypothesis on the admissible control set is not relaxed for the homogenization process of the optimal control problem. This is due to the lack of characterisation of the optimal controls, in general. The case of the positive cone gives a nice characterisation of the optimal controls in terms of the adjoint state which cancels out the product of two weak limits. Such a characterisation is not available, in general.

This article settles the problem, in non-perforated case, left open in [12] and the perforated case left open in [8]. Also, the positive cone hypothesis has been relaxed wherever possible. Moreover, it contains different approach to the results obtained for non-perforated case which easily carry forward to the perforated case. Though this article announces only the results concerning perforated domains, the results remain valid even for non-perforated domains with necessary modifications.

We end this section by recalling some preliminary notions used in the sequel. We first define the sequential notion of  $K$ -lower limit in its full generality. A detailed study of this and related notions can be found in [5]. Let  $X$  be a topological space and  $E_n$  be a sequence of subsets of  $X$ .

**Definition 1.1.** A point  $x \in X$  is said to be in the *sequential  $K$ -lower limit* of  $E_n$  w.r.t. the topology in  $X$  if and only if there exists  $k \in \mathbb{N}$  and a sequence  $\{x_n\}$  converging to  $x$  in  $X$  such that  $x_n \in E_n$ , for all  $n \geq k$ .

We now recall the notion of two-scale convergence. We refer to [1,16,20,21] for a detailed study of the same and certain applications.

**Definition 1.2.** A sequence of functions  $\{v_\varepsilon\}$  in  $L^2(\Omega)$  is said to *two-scale converge* to a limit  $v \in L^2(\Omega \times Y)$  (denoted as  $v_\varepsilon \xrightarrow{2s} v$ ) if

$$\int_{\Omega} v_\varepsilon \phi \left( x, \frac{x}{\varepsilon} \right) dx \rightarrow \int_{\Omega} \int_Y v(x, y) \phi(x, y) dy dx, \quad \forall \phi \in L^2[\Omega; C_{\text{per}}(Y)].$$

The most interesting property of two-scale convergence is the following compactness result.

**Theorem 1.3.** For any bounded sequence  $v_\varepsilon$  in  $L^2(\Omega)$ , there exist a subsequence and  $v \in L^2(\Omega \times Y)$  such that,  $v_\varepsilon$  two-scale converges to  $v$  along the subsequence. Also, if  $v_\varepsilon$  is bounded in  $H^1(\Omega)$ , then  $v$  is independent of  $y$  and is in  $H^1(\Omega)$ , and there exists  $v_1 \in L^2[\Omega; H^1_{\text{per}}(Y)]$  such that, up to a subsequence,  $\nabla v_\varepsilon$  two-scale converges to  $\nabla v + \nabla_y v_1$ .

The article is organized as follows: In Section 2, we prove  $H$ -convergence for weak data under some assumptions, one of the main results of this article. This is significant by itself and we use this result for our analysis. In this section, we work in the non-perforated set-up to fix ideas. However the results remain valid in the perforated case and are used with necessary modification in the subsequent sections. In Section 3, we study (1.1) where  $F_\varepsilon$  is the Dirichlet-type integral as given in (1.3). In Section 4, we homogenize the system (1.1)–(1.2) under the situation (1.4).

## 2. $H$ -convergence for weak data

We begin by recalling the notion of  $H$ -convergence. Let  $g_\varepsilon$  be a sequence in  $H^{-1}(\Omega)$  which converges to  $g$  strongly in  $H^{-1}(\Omega)$ . If  $v_\varepsilon$  is the solution of

$$\begin{cases} -\text{div} \left( A \left( x, \frac{x}{\varepsilon} \right) \nabla v_\varepsilon \right) = g_\varepsilon & \text{in } \Omega, \\ v_\varepsilon = 0 & \text{on } \partial\Omega, \end{cases} \tag{2.1}$$

then, there exist  $v_0$  and a matrix  $A_0$  such that

$$\left. \begin{aligned} v_\varepsilon \rightharpoonup v_0 & \text{ weakly in } H_0^1(\Omega), \\ A\left(x, \frac{x}{\varepsilon}\right) \nabla v_\varepsilon \rightharpoonup A_0 \nabla v_0 & \text{ weakly in } (L^2(\Omega))^n. \end{aligned} \right\} \tag{2.2}$$

Here  $v_0 \in H_0^1(\Omega)$  is the unique solution of the homogenized problem

$$\begin{cases} -\operatorname{div}(A_0 \nabla v_0) = g & \text{in } \Omega, \\ v_0 = 0 & \text{on } \partial\Omega. \end{cases} \tag{2.3}$$

Further, the  $ij$ th entry of the matrix  $A_0$  is given by

$$(A_0)_{ij} = \int_Y A(x, y) [\nabla_y \mu_i(x, y) + e_i] \cdot [\nabla_y \mu_j(x, y) + e_j] dy. \tag{2.4}$$

The function  $\mu_i$ , for  $1 \leq i \leq n$ , is the solution of the cell problem

$$\begin{cases} -\operatorname{div}_y(A(x, y) [\nabla_y \mu_i(x, y) + e_i]) = 0 & \text{in } Y, \\ y \mapsto \mu_i(x, y) & \text{is } Y\text{-periodic,} \end{cases} \tag{2.5}$$

where  $\{e_1, \dots, e_n\}$  is the standard basis of  $\mathbb{R}^n$ .

The above result is not true, in general, when  $g_\varepsilon$  converge weakly in  $H^{-1}(\Omega)$ . However, it was shown in [6] that if, in addition,  $g_\varepsilon$  are positive distributions of  $H^{-1}(\Omega)$ , the  $H$ -convergence result remains valid. A very general study of the weak converging data was done by L. Tartar and his result can be found in [3]. It involves a homogenized problem defined in a complicated way (involving corrector functions) and which reduces to (2.3) under the strong convergence hypothesis of data in  $H^{-1}(\Omega)$ .

In this section, we give another hypothesis on the data under which the  $H$ -convergence remains valid.

**Theorem 2.1.** *Let  $\alpha < 1$  be a fixed real number. Let  $v_\varepsilon \in H_0^1(\Omega)$  be the weak solution of (2.1), where  $g_\varepsilon \in L^2(\Omega)$  is such that  $g_\varepsilon \rightharpoonup g$  weakly in  $H^{-1}(\Omega)$  and  $\varepsilon^\alpha g_\varepsilon$  is bounded in  $L^2(\Omega)$ . Then, there exist  $v_0$  and a matrix  $A_0$  such that (2.2) is satisfied, where  $v_0 \in H_0^1(\Omega)$  is the unique solution of (2.3) and  $A_0$  is given by (2.4) and (2.5).*

**Proof.** Let  $w_\varepsilon \in H_0^1(\Omega)$  be the solution of

$$\begin{cases} -\Delta w_\varepsilon = g_\varepsilon & \text{in } \Omega, \\ w_\varepsilon = 0 & \text{on } \partial\Omega. \end{cases} \tag{2.6}$$

Then it is easy to see that  $w_\varepsilon$  is bounded in  $H_0^1(\Omega)$ . In fact,

$$\|w_\varepsilon\|_{H_0^1(\Omega)} \leq \|g_\varepsilon\|_{H^{-1}(\Omega)}.$$

Thus, it follows from the two-scale convergence theory, that there exist  $w_0 \in H_0^1(\Omega)$  and  $w_1 \in L^2[\Omega; H_{\text{per}}^1(Y)/\mathbb{R}]$  such that, up to a subsequence,

$$\left. \begin{aligned} w_\varepsilon &\xrightarrow{2s} w_0, \\ \nabla w_\varepsilon &\xrightarrow{2s} \nabla w_0(x) + \nabla_y w_1(x, y). \end{aligned} \right\} \tag{2.7}$$

Further, we see that the homogenized equation of (2.6) is

$$\begin{cases} -\Delta w_0 = g & \text{in } \Omega, \\ w_0 = 0 & \text{on } \partial\Omega. \end{cases} \tag{2.8}$$

By choosing  $\phi_1 \in \mathcal{D}(\Omega)$ ,  $\phi_2 \in C_{\text{per}}^\infty(Y)$  and setting

$$\phi\left(x, \frac{x}{\varepsilon}\right) = \varepsilon \phi_1(x) \phi_2\left(\frac{x}{\varepsilon}\right) \in \mathcal{D}[\Omega; C_{\text{per}}^\infty(Y)]$$

as a two-scale test function for (2.6), we have

$$\varepsilon \int_\Omega \nabla w_\varepsilon \cdot \nabla \phi_1 \phi_2\left(\frac{x}{\varepsilon}\right) dx + \int_\Omega \nabla w_\varepsilon \cdot \nabla_y \phi_2\left(\frac{x}{\varepsilon}\right) \phi_1 dx = \varepsilon^{1-\alpha} \int_\Omega \varepsilon^\alpha g_\varepsilon \phi_1(x) \phi_2\left(\frac{x}{\varepsilon}\right) dx.$$

By passing to the two-scale limit, we get

$$\int_{\Omega} \int_Y [\nabla w_0 + \nabla_y w_1(x, y)] \cdot \nabla_y \phi_2(y) \phi_1(x) \, dx \, dy = 0.$$

Since  $\int_Y \nabla_y \phi_2(y) \, dy = 0$ , we have for every  $\phi_1 \in \mathcal{D}(\Omega)$ ,

$$\int_{\Omega} \phi_1(x) \left[ \int_Y \nabla_y w_1(x, y) \cdot \nabla_y \phi_2(y) \, dy \right] \, dx = 0.$$

Thus, for almost every  $x \in \Omega$ ,

$$\int_Y \nabla_y w_1(x, y) \cdot \nabla_y \phi_2(y) \, dy = 0,$$

which implies  $\Delta_y w_1(x, y) = 0$ , where  $w_1$  is  $Y$ -periodic. Thus,  $w_1$  is a function independent of  $y$  and  $\nabla_y w_1 = 0$ . Therefore in (2.7) we, in fact, have

$$\nabla w_{\varepsilon} \xrightarrow{2s} \nabla w_0(x).$$

Observe that the solution  $v_{\varepsilon}$  of (2.1) is bounded in  $H_0^1(\Omega)$ . Thus, there exist  $v_0 \in H_0^1(\Omega)$  and  $v_1 \in L^2[\Omega; H_{\text{per}}^1(Y)/\mathbb{R}]$  such that, up to a subsequence,

$$\left. \begin{aligned} v_{\varepsilon} &\xrightarrow{2s} v_0, \\ \nabla v_{\varepsilon} &\xrightarrow{2s} \nabla v_0(x) + \nabla_y v_1(x, y). \end{aligned} \right\} \tag{2.9}$$

Now an usual two-scale analysis of (2.1) by replacing  $g_{\varepsilon}$  with Eq. (2.6) and by choosing  $\phi(x, \frac{x}{\varepsilon}) = \phi_1(x) + \varepsilon \phi_2(x, \frac{x}{\varepsilon})$  as the two-scale test function will yield,

$$\int_{\Omega} \int_Y \{A(x, y) [\nabla v_0 + \nabla_y v_1(x, y)] - \nabla w_0\} \cdot \{\nabla \phi_1(x) + \nabla_y \phi_2(x, y)\} \, dx \, dy = 0.$$

Then, the usual arguments of decoupling and Eq. (2.8) together will yield (2.3) and the convergences (2.2).  $\square$

### 3. Dirichlet-type integral of state in cost functional

Let  $U_{\varepsilon}$  be a closed convex subset of  $L^2(\Omega_{\varepsilon})$ . The optimal control problem is to minimise in  $U_{\varepsilon}$ , the cost functional  $J_{\varepsilon}$  given as,

$$J_{\varepsilon}(\theta_{\varepsilon}) = \frac{1}{2} \int_{\Omega_{\varepsilon}} B\left(x, \frac{x}{\varepsilon}\right) \nabla u_{\varepsilon} \cdot \nabla u_{\varepsilon} \, dx + \frac{\varepsilon}{2} \|\theta_{\varepsilon}\|_{2, \Omega_{\varepsilon}}^2, \quad \forall \theta_{\varepsilon} \in U_{\varepsilon}, \tag{3.1}$$

where  $u_{\varepsilon} \in H^1(\Omega_{\varepsilon})$  is the unique solution of

$$\begin{cases} -\text{div}\left(A\left(x, \frac{x}{\varepsilon}\right) \nabla u_{\varepsilon}\right) = f + \theta_{\varepsilon} & \text{in } \Omega_{\varepsilon}, \\ A\left(x, \frac{x}{\varepsilon}\right) \nabla u_{\varepsilon} \cdot \nu = 0 & \text{on } \partial\Omega_{\varepsilon} \setminus \partial\Omega, \\ u_{\varepsilon} = 0 & \text{on } \partial\Omega \cap \partial\Omega_{\varepsilon}, \end{cases} \tag{3.2}$$

and  $\nu$  is the unit outward normal on  $\partial\Omega_{\varepsilon} \setminus \partial\Omega$ .

Let  $\theta_{\varepsilon}^* \in U_{\varepsilon}$  be the unique optimal control and  $u_{\varepsilon}^*$  be the state corresponding to  $\theta_{\varepsilon}^*$  (cf. (3.12)). The solution space of (3.2) is given as,

$$V_{\varepsilon} = \{u \in H^1(\Omega_{\varepsilon}) \mid u = 0 \text{ on } \partial\Omega \cap \partial\Omega_{\varepsilon}\}$$

and  $u_{\varepsilon}^* \in V_{\varepsilon}$ . We now state a Poincaré inequality result of  $\Omega_{\varepsilon}$ , proved in [2].

**Lemma 3.1.** (See [2, Lemma A.4].) *There exists a positive constant  $C$ , independent of  $\varepsilon$ , such that*

$$\|v\|_{2, \Omega_{\varepsilon}} \leq C \|\nabla v\|_{2, \Omega_{\varepsilon}}, \quad \forall v \in V_{\varepsilon}.$$

The above lemma gives an equivalent norm on  $V_\varepsilon$  as,  $\|v\|_{V_\varepsilon} = \|\nabla v\|_{2,\Omega_\varepsilon}$ .

We introduce the adjoint optimal state associated to the optimal control problem. Let  $p_\varepsilon^* \in V_\varepsilon$  be the solution of,

$$\begin{cases} -\operatorname{div}\left({}^t A\left(x, \frac{x}{\varepsilon}\right) \nabla p_\varepsilon^*\right) = -\operatorname{div}\left(B\left(x, \frac{x}{\varepsilon}\right) \nabla u_\varepsilon^*\right) & \text{in } \Omega_\varepsilon, \\ \left({}^t A\left(x, \frac{x}{\varepsilon}\right) \nabla p_\varepsilon^* - B\left(x, \frac{x}{\varepsilon}\right) \nabla u_\varepsilon^*\right) \cdot \nu = 0 & \text{on } \partial\Omega_\varepsilon \setminus \partial\Omega, \\ p_\varepsilon^* = 0 & \text{on } \partial\Omega \cap \partial\Omega_\varepsilon. \end{cases} \quad (3.3)$$

Then the optimality condition, in terms of the adjoint optimal state, is

$$\int_{\Omega_\varepsilon} (p_\varepsilon^* + \varepsilon\theta_\varepsilon^*)(\theta_\varepsilon - \theta_\varepsilon^*) dx \geq 0, \quad \forall \theta_\varepsilon \in U_\varepsilon. \quad (3.4)$$

Note that the symmetry hypothesis on  $B$  comes in hand only to derive the optimality condition (3.4).

In the rest of the section we will consider control sets,  $U_\varepsilon$ , satisfying the following hypothesis:

(H1) There exists  $\eta_\varepsilon \in U_\varepsilon$  such that  $\{\tilde{\eta}_\varepsilon\}$  is bounded in  $L^2(\Omega)$ .

The motivation for this hypothesis will be clear in next lemma. We shall now show that the admissible sets satisfying (H1) are in abundance. Let  $U$  be a closed convex subset of  $L^2(\Omega)$  and  $U_\varepsilon$  be the set of all elements of  $U$  restricted to  $\Omega_\varepsilon$ . Observe that  $U_\varepsilon$  is a closed convex subset of  $L^2(\Omega_\varepsilon)$ . Let  $\theta \in U$ , then by choosing  $\eta_\varepsilon = \theta|_{\Omega_\varepsilon}$  we note that  $U_\varepsilon$  satisfies (H1), since  $\|\eta_\varepsilon\|_{2,\Omega_\varepsilon}^2 \leq \|\theta\|_2^2$ .

**Lemma 3.2.** *Let the admissible set  $U_\varepsilon$  satisfy the hypothesis (H1), then  $\{u_\varepsilon^*\}$  and  $\{p_\varepsilon^*\}$  are bounded in  $V_\varepsilon$ . Also, there exists  $\theta^* \in H^{-1}(\Omega)$  such that, for a subsequence,*

$$\tilde{\theta}_\varepsilon^* \rightharpoonup \theta^* \quad \text{weakly in } H^{-1}(\Omega) \quad (3.5)$$

and

$$\varepsilon^{1/2}\tilde{\theta}_\varepsilon^* \rightharpoonup 0 \quad \text{weakly in } L^2(\Omega). \quad (3.6)$$

**Proof.** Let  $v_\varepsilon$  be the state corresponding to the control  $\theta_\varepsilon = \eta_\varepsilon$  in (3.2). Then, it is clear, from the hypotheses on  $A(x, \frac{x}{\varepsilon})$  that  $v_\varepsilon$  is bounded uniformly with respect to  $\varepsilon$  in  $V_\varepsilon$ , since,

$$\|v_\varepsilon\|_{V_\varepsilon} \leq \frac{C}{a} \left\| \chi\left(\frac{x}{\varepsilon}\right) f + \tilde{\eta}_\varepsilon \right\|_2.$$

Hence, it follows from (1.5) that

$$J_\varepsilon(\theta_\varepsilon^*) \leq J_\varepsilon(\eta_\varepsilon).$$

Therefore, from (3.1) we obtain that  $\{u_\varepsilon^*\}$  and  $\{\varepsilon^{1/2}\theta_\varepsilon^*\}$  are bounded in  $V_\varepsilon$  and  $L^2(\Omega_\varepsilon)$ , respectively. Moreover, from (3.3),  $\{p_\varepsilon^*\}$  is bounded in  $V_\varepsilon$ . For  $v \in H_0^1(\Omega)$ , consider

$$\left| \int_{\Omega} \tilde{\theta}_\varepsilon^* v dx \right| = \left| \int_{\Omega_\varepsilon} \theta_\varepsilon^* v dx \right| = \left| \int_{\Omega} A_\varepsilon \widetilde{\nabla u_\varepsilon^*} \cdot \nabla v dx - \int_{\Omega} \chi\left(\frac{x}{\varepsilon}\right) f v dx \right| \leq \left( b \|u_\varepsilon^*\|_{V_\varepsilon} + C \left\| \chi\left(\frac{x}{\varepsilon}\right) f \right\|_2 \right) \|v\|_{H_0^1(\Omega)}.$$

Therefore  $\{\tilde{\theta}_\varepsilon^*\}$  is bounded in  $H^{-1}(\Omega)$  and thus there exists  $\theta^* \in H^{-1}(\Omega)$  such that, up to a subsequence, (3.5) holds and hence (3.6) holds.  $\square$

Our objective is to know whether  $\theta^*$  is an optimal control of a problem similar to (3.1)–(3.2). We deduce from the *a priori* estimates obtained in Lemma 3.2 that there exist  $u^*, p^* \in H_0^1(\Omega)$  and  $u_1, p_1 \in L^2[\Omega; H_{\text{per}}^1(Y')/\mathbb{R}]$  such that, up to a subsequence,

$$\begin{cases} \widetilde{u_\varepsilon^*} \xrightarrow{2s} u^*(x)\chi(y), \\ \widetilde{\nabla u_\varepsilon^*} \xrightarrow{2s} \chi(y)[\nabla u^*(x) + \nabla_y u_1(x, y)], \\ \widetilde{p_\varepsilon^*} \xrightarrow{2s} p^*(x)\chi(y), \\ \widetilde{\nabla p_\varepsilon^*} \xrightarrow{2s} \chi(y)[\nabla p^*(x) + \nabla_y p_1(x, y)]. \end{cases}$$

The above results can be derived from the two-scale convergence analysis which we skip (cf. [1]).

Observe that the right-hand side of both (3.2) (written for  $\theta_\varepsilon = \theta_\varepsilon^*$ ) and (3.3) involve functions whose extensions to  $\Omega$  are only weakly compact in  $H^{-1}(\Omega)$ . A homogenization result is not available, in general, for such equations.

We shall now define some cell problems which will be used in the sequel to identify the limit problem. For  $1 \leq i \leq n$ , let the function  $\mu_i \in H^1_{\text{per}}(Y')/\mathbb{R}$  be the solution of the cell problem

$$\begin{cases} -\text{div}_y(A(x, y)[\nabla_y \mu_i(x, y) + e_i]) = 0 & \text{in } Y', \\ A(x, y)[\nabla_y \mu_i(x, y) + e_i] \cdot \nu = 0 & \text{on } \partial Y' \setminus \partial Y, \\ y \mapsto \mu_i(x, y) & \text{is } Y\text{-periodic.} \end{cases} \tag{3.7}$$

A note of caution that we are using the same notation  $\mu_i$  as in (2.5). But in the rest of the article  $\mu_i$  will always denote the solution of (3.7). The distinction is clear if one keeps in mind that we are dealing with the perforated situation in this article.

Let  $\omega_i \in H^1_{\text{per}}(Y')/\mathbb{R}$  be the solution of the adjoint cell problem

$$\begin{cases} -\text{div}_y({}^t A(x, y)[\nabla_y \omega_i(x, y) + e_i]) = 0 & \text{in } Y', \\ {}^t A(x, y)[\nabla_y \omega_i(x, y) + e_i] \cdot \nu = 0 & \text{on } \partial Y' \setminus \partial Y, \\ y \mapsto \omega_i(x, y) & \text{is } Y\text{-periodic,} \end{cases} \tag{3.8}$$

and let  $\psi_i \in H^1_{\text{per}}(Y')/\mathbb{R}$  be the solution of the cell problem

$$\begin{cases} -\text{div}_y({}^t A(x, y)\nabla_y \psi_i(x, y) + B(x, y)\{\nabla_y \mu_i + e_i\}) = 0 & \text{in } Y', \\ [{}^t A(x, y)\nabla_y \psi_i(x, y) + B(x, y)\{e_i + \nabla_y \mu_i\}] \cdot \nu = 0 & \text{on } \partial Y' \setminus \partial Y, \\ y \mapsto \psi_i(x, y) & \text{is } Y\text{-periodic.} \end{cases} \tag{3.9}$$

We now provide the homogenization theorem for the state and adjoint-state equations. The non-perforated analogue of the following theorem was proved in [8,19] under the hypothesis that the admissible control set is the positive cone of  $L^2(\Omega)$ . The theorem given below relaxes this hypothesis on the admissible control set, thus proving the result in its full generality both for fixed domains and varying domains with Neumann condition on the holes.

Let the homogenized matrix  $A_0$  be defined as,

$$(A_0)_{ij} = \int_{Y'} A(x, y)[\nabla_y \mu_i(x, y) + e_i] \cdot [\nabla_y \mu_j(x, y) + e_j] dy \tag{3.10}$$

and the transpose of  $A_0$ ,  ${}^t A_0$ , is given by a similar equation written for the function  $\omega_i$  (cf. (4.9)). Let  $B^\sharp$  be given as,

$$(B^\sharp)e_i = \int_{Y'} \{B(x, y)[\nabla_y \mu_i(x, y) + e_i] + {}^t A(x, y)\nabla_y \psi_i(x, y)\} dy. \tag{3.11}$$

**Theorem 3.3.** *Let  $U_\varepsilon$  satisfy the hypothesis (H1) and let  $u_\varepsilon^*$  be the solution of*

$$\begin{cases} -\text{div}\left(A\left(x, \frac{x}{\varepsilon}\right)\nabla u_\varepsilon^*\right) = f + \theta_\varepsilon^* & \text{in } \Omega_\varepsilon, \\ A\left(x, \frac{x}{\varepsilon}\right)\nabla u_\varepsilon^* \cdot \nu = 0 & \text{on } \partial\Omega_\varepsilon \setminus \partial\Omega, \\ u_\varepsilon^* = 0 & \text{on } \partial\Omega \cap \partial\Omega_\varepsilon, \end{cases} \tag{3.12}$$

and  $p_\varepsilon^*$  be the solution of (3.3). Then,  $\chi_0 u^*$  and  $\chi_0 p^*$  are the weak limits of  $\tilde{u}_\varepsilon^*$  and  $\tilde{p}_\varepsilon^*$ , respectively, in  $H^1_0(\Omega)$  where  $(u^*, p^*) \in H^1_0(\Omega) \times H^1_0(\Omega)$  solves

$$\begin{cases} -\text{div}(A_0 \nabla u^*) = \chi_0 f + \theta^* & \text{in } \Omega, \\ -\text{div}({}^t A_0 \nabla p^* - B^\sharp \nabla u^*) = 0 & \text{in } \Omega, \\ u^* = p^* = 0 & \text{on } \partial\Omega. \end{cases} \tag{3.13}$$

The control  $\theta^*$  is as obtained in (3.5).

**Proof.** Since  $U_\varepsilon$  satisfies (H1), the results of Lemma 3.2 remain valid. Let  $\rho_\varepsilon^* \in V_\varepsilon$  be the solution of

$$\begin{cases} -\Delta \rho_\varepsilon^* = \theta_\varepsilon^* & \text{in } \Omega_\varepsilon, \\ \nabla \rho_\varepsilon^* \cdot \nu = 0 & \text{on } \partial\Omega_\varepsilon \setminus \partial\Omega, \\ \rho_\varepsilon^* = 0 & \text{on } \partial\Omega \cap \partial\Omega_\varepsilon. \end{cases} \tag{3.14}$$

Then,

$$\|\rho_\varepsilon^*\|_{V_\varepsilon}^2 \leq \|\theta_\varepsilon^*\|_{H^{-1}(\Omega)} \|P_\varepsilon \rho_\varepsilon^*\|_{H_0^1(\Omega)},$$

where  $P_\varepsilon$  is an extension operator (cf. [4, Lemma 1]) from  $V_\varepsilon$  to  $H_0^1(\Omega)$  such that  $\|\nabla(P_\varepsilon v)\|_2 \leq C\|\nabla v\|_{2,\Omega_\varepsilon}$ , for all  $v \in V_\varepsilon$ , and the constant  $C$  is independent of  $\varepsilon$ . Thus,  $\widetilde{\nabla \rho_\varepsilon^*}$  is bounded in  $(L^2(\Omega))^n$ . Also, by Lemma 3.1,  $\widetilde{\rho_\varepsilon^*}$  is bounded in  $L^2(\Omega)$ . Thus, by the two-scale convergence theory, there exist  $\rho^* \in H_0^1(\Omega)$  and  $\rho_1 \in L^2[\Omega; H_{\text{per}}^1(Y')/\mathbb{R}]$  such that, up to a subsequence,

$$\begin{cases} \widetilde{\rho_\varepsilon^*} \xrightarrow{2S} \rho^*(x)\chi(y), \\ \widetilde{\nabla \rho_\varepsilon^*} \xrightarrow{2S} \chi(y)[\nabla \rho^*(x) + \nabla_y \rho_1(x, y)]. \end{cases} \tag{3.15}$$

It is easy to see that the homogenized equation of (3.14) is

$$\begin{cases} -\Delta \rho^* = \frac{1}{\chi_0} \theta^* & \text{in } \Omega, \\ \rho^* = 0 & \text{on } \partial\Omega. \end{cases} \tag{3.16}$$

Arguing as in the proof of Theorem 2.1 with the obvious modification for perforated domain, we have the first equation of (3.13) as the homogenized state equation. We also have that,

$$u_1(x, y) = \sum_{i=1}^n \mu_i(x, y) \frac{\partial u^*}{\partial x_i}(x).$$

We shall now show that the second equation of (3.13) is the homogenized equation corresponding to adjoint equation (3.3). In the case of (3.3),  $g_\varepsilon = -\text{div}(B(x, \frac{x}{\varepsilon})\nabla u_\varepsilon^*)$  may not satisfy the hypotheses of Theorem 2.1, but here we obtain the limit equation using the divergence form of  $g_\varepsilon$ .

We choose  $\phi(x, \frac{x}{\varepsilon}) = \phi_1(x) + \varepsilon\phi_2(x, \frac{x}{\varepsilon})$  as a two-scale test function in (3.3), and passing to the limit, we have

$$\begin{aligned} & \int_{\Omega} \int_Y A(x, y)\chi(y)\{\nabla p^* + \nabla_y p_1(x, y)\} \cdot [\nabla \phi_1(x) + \nabla_y \phi_2(x, y)] dy dx \\ &= \int_{\Omega} \int_Y B(x, y)\chi(y)\{\nabla u^* + \nabla_y u_1(x, y)\} \cdot [\nabla \phi_1(x) + \nabla_y \phi_2(x, y)] dy dx. \end{aligned}$$

By choosing  $\phi_1 = 0$  and the cell problems (3.8) and (3.9), we deduce that

$$p_1 = \sum_{i=1}^n \left[ \psi_i(x, y) \frac{\partial u^*}{\partial x_i}(x) + \omega_i(x, y) \frac{\partial p^*}{\partial x_i}(x) \right]$$

and by choosing  $\phi_2 = 0$ , we get the second equation of (3.13).  $\square$

**Remark 3.4.** Let  $u_\varepsilon$  be the solution of (3.2) and if  $\widetilde{\theta_\varepsilon} \rightarrow \theta$  strongly in  $H^{-1}(\Omega)$ , then

$$\int_{\Omega_\varepsilon} B_\varepsilon \nabla u_\varepsilon \cdot \nabla u_\varepsilon dx \rightarrow \int_{\Omega} B^\sharp \nabla u_0 \cdot \nabla u_0 dx \tag{3.17}$$

where  $B^\sharp$  is as given in (3.11) and  $u_0 = u_0(\theta) \in H_0^1(\Omega)$  is the unique solution of

$$\begin{cases} -\text{div}(A_0 \nabla u_0) = \chi_0 f + \theta & \text{in } \Omega, \\ u_0 = 0 & \text{on } \partial\Omega. \end{cases}$$

$A_0$  is given as in (3.10). This result is, in a sense, the ‘generalised’ energy convergence. This fact has been observed in [9,11]. We, however, remark that (3.17) is not true, in general, when  $\widetilde{\theta_\varepsilon} \rightarrow \theta$  weakly in  $H^{-1}(\Omega)$ . A one dimensional example establishing this fact for the non-perforated domain can be found in [19].

**Remark 3.5.** If  $B_0$  is the homogenized limit of the sequence  $B(x, \frac{x}{\varepsilon})$ , then, in general,  $B^\sharp \neq B_0$ . For more such interesting properties of  $B^\sharp$  and its applications we refer to [7,9–11,13,22].



Now that we have identified the limit state equation, it now remains for us to find the limit cost functional and to prove that  $\theta^*$  is its optimal control.

Using  $p_\varepsilon^*$  as a test function in (3.12) and  $u_\varepsilon^*$  as a test function in (3.3), one can rewrite the cost functional  $J_\varepsilon(\theta_\varepsilon^*)$  as,

$$J_\varepsilon(\theta_\varepsilon^*) = \frac{1}{2} \int_{\Omega_\varepsilon} (f + \theta_\varepsilon^*) p_\varepsilon^* dx + \frac{\varepsilon}{2} \|\theta_\varepsilon^*\|_{2, \Omega_\varepsilon}^2. \tag{3.18}$$

The above functional (3.18) involves product of two weakly converging sequence, namely,  $\int_{\Omega_\varepsilon} \theta_\varepsilon^* p_\varepsilon^* dx$ , whose limit is unknown in general. This makes the problem difficult to tackle in a general admissible control set  $U_\varepsilon$ .

We could identify the limit cost functional only in the situation when the control set is the set of all positive elements of  $L^2(\Omega_\varepsilon)$ , i.e., the positive cone of  $L^2(\Omega_\varepsilon)$ . More precisely, let  $U_\varepsilon = \{\theta \in L^2(\Omega) \mid \theta \geq 0 \text{ a.e. in } \Omega_\varepsilon\}$  be the closed convex subset of  $L^2(\Omega)$ . Observe that the elements of  $U_\varepsilon$  restricted to  $\Omega_\varepsilon$  is the set of all positive elements of  $L^2(\Omega_\varepsilon)$ . Let  $U'$  be the sequential  $K$ -lower limit of  $U_\varepsilon$  with respect to the strong topology in  $L^2(\Omega)$ . We shall now establish that  $U'$  is non-empty. If  $U$  denotes the positive cone of  $L^2(\Omega)$ , i.e.,  $U = \{\theta \in L^2(\Omega) \mid \theta \geq 0 \text{ a.e. in } \Omega\}$ , then  $U$  is a subset of  $U_\varepsilon$  for all  $\varepsilon$ . Thus,  $U$  is a subset of  $U'$  and hence  $U'$  is non-empty.

Let  $V'$  be the strong closure of  $U'$  in  $H^{-1}(\Omega)$ . Since  $U'$  is convex,  $V'$  is same as the weak closure of  $U'$  in  $H^{-1}(\Omega)$ .

**Remark 3.6.** If  $V$  is the strong closure of  $U$  in  $H^{-1}(\Omega)$  then  $V$  is the set of all positive distributions of  $H^{-1}(\Omega)$ . In other words,  $V$  is the positive cone of  $H^{-1}(\Omega)$ . We refer to [19] for a proof of this fact. Also, observe that  $V$  is a subset of  $V'$ . Since  $U$  is convex,  $V$  is same as the weak closure of  $U$  in  $H^{-1}(\Omega)$ .

**Theorem 3.7.** Let  $U_\varepsilon = \{\theta \in L^2(\Omega) \mid \theta \geq 0 \text{ a.e. in } \Omega_\varepsilon\}$ . Then

$$J_\varepsilon(\theta_\varepsilon^*) \rightarrow \frac{1}{2} \int_{\Omega} B^\sharp \nabla u^* \cdot \nabla u^* dx - \frac{1}{2} \langle \theta^*, p^* \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} \tag{3.19}$$

where  $\theta^*$  is as obtained in Lemma 3.2,  $u^*, p^*$  are as obtained in (3.13) and  $B^\sharp$  is given by (3.11). Also,  $p^* \in U$  and hence in  $V'$ .

Further, if  $\langle \theta^*, p^* \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} = 0$ , then  $\theta^*$  minimizes

$$J(\theta) = \frac{1}{2} \int_{\Omega} B^\sharp \nabla u \cdot \nabla u dx \tag{3.20}$$

over  $V'$ , where  $u = u(\theta)$  solves

$$\begin{cases} -\text{div}(A_0 \nabla u) = \chi_0 f + \theta & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \tag{3.21}$$

and the convergences (3.5) and (3.6) hold for the entire sequence.

**Proof.** Observe that  $0 \in U_\varepsilon$  for all  $\varepsilon$ . Thus, the set of all elements of  $U_\varepsilon$  restricted to  $\Omega_\varepsilon$  (the positive cone of  $L^2(\Omega_\varepsilon)$ ) satisfies the hypothesis (H1). Hence the results of Lemma 3.2 and Theorem 3.3 remain valid.

Since the admissible set is now the positive cone of  $L^2(\Omega_\varepsilon)$ , the optimality condition (3.4) implies that  $\varepsilon \theta_\varepsilon^* = (p_\varepsilon^*)^-$ . This equality will in turn imply that

$$\int_{\Omega_\varepsilon} \theta_\varepsilon^* p_\varepsilon^* dx = -\varepsilon \|\theta_\varepsilon^*\|_{2, \Omega_\varepsilon}^2. \tag{3.22}$$

Thus,  $J_\varepsilon(\theta_\varepsilon^*)$  (cf. (3.18)) reduces to

$$J_\varepsilon(\theta_\varepsilon^*) = \frac{1}{2} \int_{\Omega_\varepsilon} f p_\varepsilon^* dx$$

and passing to the limit will yield

$$J_\varepsilon(\theta_\varepsilon^*) \rightarrow \frac{1}{2} \int_{\Omega} \chi_0 f p^* dx.$$

Using  $p^*$  and  $u^*$  as test functions in the state and adjoint-state equation of (3.13), respectively, we have

$$\int_{\Omega} \chi_0 f p^* dx = \int_{\Omega} B^\sharp \nabla u^* \cdot \nabla u^* dx - \langle \theta^*, p^* \rangle_{H^{-1}(\Omega), H_0^1(\Omega)}.$$

Thus, (3.19) holds true.

Using (3.22), we rewrite the optimality condition (3.4) as,

$$\int_{\Omega_\varepsilon} p_\varepsilon^* \theta \, dx + \int_{\Omega_\varepsilon} (\varepsilon^{\frac{1}{2}} \theta_\varepsilon^*) (\varepsilon^{\frac{1}{2}} \theta) \, dx \geq 0, \quad \forall \theta \in U_\varepsilon.$$

In particular, the above inequality holds for all  $\theta \in U$ . Thus, passing to the limit, we deduce that

$$\int_{\Omega} \chi_0 p^* \theta \, dx \geq 0, \quad \forall \theta \in U.$$

This implies  $p^* \geq 0$ , i.e.,  $p^* \in U$  and hence  $p^* \in V'$ .

For  $\theta \in V'$ , let  $J$  be defined as in (3.20) where  $u$  solves (3.21). Recall that  $\theta_\varepsilon^*$  is the minimiser of  $J_\varepsilon$  over  $U_\varepsilon$ , i.e.,

$$J_\varepsilon(\theta_\varepsilon^*) \leq J_\varepsilon(\theta), \quad \forall \theta \in U_\varepsilon. \quad (3.23)$$

Given  $\theta \in U'$ , there exists  $\delta > 0$  and a sequence  $\theta_\varepsilon \rightarrow \theta$  strongly in  $L^2(\Omega)$  such that  $\theta_\varepsilon \in U_\varepsilon$  for all  $\varepsilon < \delta$ . Observe that

$$\|\theta_\varepsilon\|_{2, \Omega_\varepsilon}^2 \leq \|\theta_\varepsilon\|_2^2, \quad \forall \varepsilon < \delta,$$

and hence

$$\|\tilde{\theta}_\varepsilon\|_2^2 \leq \|\theta_\varepsilon\|_2^2, \quad \forall \varepsilon < \delta.$$

Thus  $\varepsilon \|\theta_\varepsilon\|_{2, \Omega_\varepsilon}^2 \rightarrow 0$ . Thus, passing to the limit in (3.23) (also using Remark 3.4), we deduce

$$\frac{1}{2} \int_{\Omega} B^\sharp \nabla u^* \cdot \nabla u^* \, dx - \frac{1}{2} \langle \theta^*, p^* \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} \leq J(\theta), \quad \forall \theta \in U'.$$

Since  $V'$  is the strong closure of  $U'$  in  $H^{-1}(\Omega)$ , we actually have (using Remark 3.4)

$$J(\theta^*) - \frac{1}{2} \langle \theta^*, p^* \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} \leq J(\theta), \quad \forall \theta \in V'.$$

Observe that  $\tilde{\theta}_\varepsilon^* \in U \subset U_\varepsilon$ . Thus, the weak limit of  $\tilde{\theta}_\varepsilon^*$  in  $H^{-1}(\Omega)$  is in  $V \subset V'$  by Remark 3.6. Therefore, if  $\langle \theta^*, p^* \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} = 0$ , then  $\theta^*$  minimises  $J$  over  $V'$ . The strict convexity of  $J$  would then imply the uniqueness of  $\theta^*$  and thus the convergences (3.5) and (3.6) hold for the entire sequence.  $\square$

**Remark 3.8.** We note that though we have identified the possible limit cost functional as (3.20), the possible limit admissible set as  $V'$  and the possible homogenized state equation as (3.21), we are unable to prove that  $\theta^*$  minimises  $J$  in  $V'$  in general. However, we prove  $\theta^*$  is a minimiser of  $J$  in  $V'$  under the hypothesis that  $\langle \theta^*, p^* \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} = 0$ . Considering the fact that  $\int_{\Omega_\varepsilon} \theta_\varepsilon^* p_\varepsilon^* \, dx \leq 0$  (cf. (3.22)) and  $\langle \theta^*, p^* \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} \geq 0$ , one would expect that  $\langle \theta^*, p^* \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} = 0$ . However, we have no means of proving this hypothesis.

In the next section, we shall study an optimal control problem with a different cost functional (cf. (1.4)) where the difficulties noted in the above remark are overcome for the positive cone situation.

#### 4. $L^2$ -norm of state in the cost functional

Let  $U_\varepsilon$  be a closed convex subset of  $L^2(\Omega_\varepsilon)$ . The optimal control problem is to minimise the cost functional  $J_\varepsilon$  given as,

$$J_\varepsilon(\theta_\varepsilon) = \frac{1}{2} \|u_\varepsilon\|_{2, \Omega_\varepsilon}^2 + \frac{\varepsilon}{2} \|\theta_\varepsilon\|_{2, \Omega_\varepsilon}^2, \quad (4.1)$$

for all  $\theta_\varepsilon \in U_\varepsilon$ , where  $u_\varepsilon \in H^1(\Omega_\varepsilon)$  is the unique solution of (3.2). From classical theory, there exists a unique  $\theta_\varepsilon^* \in U_\varepsilon$  which minimises  $J_\varepsilon$  in  $U_\varepsilon$ . Let  $u_\varepsilon^* \in V_\varepsilon$  be the state corresponding to  $\theta_\varepsilon^*$ .

We now introduce the adjoint optimal state associated to the optimal control problem. Let  $p_\varepsilon^* \in V_\varepsilon$  be the solution of,

$$\begin{cases} -\operatorname{div} \left( {}^t A \left( x, \frac{x}{\varepsilon} \right) \nabla p_\varepsilon^* \right) = u_\varepsilon^* & \text{in } \Omega_\varepsilon, \\ {}^t A \left( x, \frac{x}{\varepsilon} \right) \nabla p_\varepsilon^* \cdot \nu = 0 & \text{on } \partial \Omega_\varepsilon \setminus \partial \Omega, \\ p_\varepsilon^* = 0 & \text{on } \partial \Omega \cap \partial \Omega_\varepsilon. \end{cases} \quad (4.2)$$

Then the optimality condition is as given in (3.4).

**Lemma 4.1.** Let  $U_\varepsilon$  satisfy the hypothesis (H1), then  $\{p_\varepsilon^*\}$  is bounded in  $V_\varepsilon$ . Also,  $\{u_\varepsilon^*\}$  and  $\{\varepsilon^{1/2}\theta_\varepsilon^*\}$  are bounded in  $L^2(\Omega_\varepsilon)$ . Thus, there exist  $u_0(x, y) \in L^2(\Omega \times Y)$ ,  $p^* \in H_0^1(\Omega)$  and  $p_1 \in L^2[\Omega; H_{\text{per}}^1(Y)/\mathbb{R}]$  such that, for a subsequence,

$$\varepsilon^{1/2}\tilde{\theta}_\varepsilon^* \rightharpoonup 0 \quad \text{weakly in } L^2(\Omega), \tag{4.3}$$

$$\tilde{u}_\varepsilon^* \xrightarrow{2s} u_0(x, y), \tag{4.4}$$

$$\tilde{p}_\varepsilon^* \xrightarrow{2s} p^*(x)\chi(y), \tag{4.5}$$

and

$$\widetilde{\nabla p}_\varepsilon^* \xrightarrow{2s} \chi(y)[\nabla p^*(x) + \nabla_y p_1(x, y)]. \tag{4.6}$$

The limit  $p^*$  solves

$$\begin{cases} -\text{div}({}^t A_0 \nabla p^*) = \int_{Y'} u_0(x, y) dy & \text{in } \Omega, \\ p^* = 0 & \text{on } \partial\Omega, \end{cases} \tag{4.7}$$

and

$$p_1 = \sum_{i=1}^n \omega_i(x, y) \frac{\partial p^*}{\partial x_i}(x). \tag{4.8}$$

The entries of  ${}^t A_0$  are given as

$$({}^t A_0)_{ij} = \int_{Y'} {}^t A(x, y)[\nabla_y \omega_i(x, y) + e_i] \cdot [\nabla_y \omega_j(x, y) + e_j] dy \tag{4.9}$$

where  $\omega_i(x, y)$  is the solution to the cell problem (3.8).

**Proof.** Let  $v_\varepsilon$  be the state corresponding to the control  $\eta_\varepsilon$ , given by the hypothesis (H1), in (3.2). Then, as argued in Lemma 3.2,  $v_\varepsilon$  is bounded uniformly with respect to  $\varepsilon$  in  $V_\varepsilon$ . Hence, it follows from (1.5) that  $\{u_\varepsilon^*\}$  and  $\{\varepsilon^{1/2}\theta_\varepsilon^*\}$  are both bounded in  $L^2(\Omega_\varepsilon)$ . Moreover, from (4.2),  $\{p_\varepsilon^*\}$  is bounded in  $V_\varepsilon$ . Thus there exist  $\theta' \in L^2(\Omega)$ ,  $u_0(x, y) \in L^2(\Omega \times Y)$ ,  $p^* \in H_0^1(\Omega)$  and  $p_1 \in L^2[\Omega; H_{\text{per}}^1(Y)/\mathbb{R}]$  such that, for a subsequence,

$$\varepsilon^{1/2}\tilde{\theta}_\varepsilon^* \rightharpoonup \theta' \quad \text{weakly in } L^2(\Omega),$$

and (4.4)–(4.6) are satisfied.

Now an usual two-scale analysis of (4.2) will yield (4.7) and (4.8). It now only remains to prove that  $\theta' = 0$ .

Let  $q_\varepsilon \in V_\varepsilon$  be the solution of

$$\begin{cases} -\text{div}\left(A\left(x, \frac{x}{\varepsilon}\right)\nabla q_\varepsilon\right) = \varepsilon^{\frac{1}{2}}(f + \theta_\varepsilon^*) & \text{in } \Omega_\varepsilon, \\ A\left(x, \frac{x}{\varepsilon}\right)\nabla q_\varepsilon \cdot \nu = 0 & \text{on } \partial\Omega_\varepsilon \setminus \partial\Omega, \\ q_\varepsilon = 0 & \text{on } \partial\Omega \cap \partial\Omega_\varepsilon. \end{cases} \tag{4.10}$$

By linearity,  $q_\varepsilon = \varepsilon^{\frac{1}{2}}u_\varepsilon^*$ . Since,  $\varepsilon^{\frac{1}{2}}\theta_\varepsilon^*$  is bounded in  $L^2(\Omega_\varepsilon)$ ,  $q_\varepsilon$  is bounded in  $V_\varepsilon$  and thus, by Lemma 3.1,  $q_\varepsilon$  is bounded in  $L^2(\Omega_\varepsilon)$ . This together with the fact that  $\tilde{u}_\varepsilon^*$  is bounded in  $L^2(\Omega)$  will imply that, for a subsequence,

$$\tilde{q}_\varepsilon \rightarrow 0 \quad \text{strongly in } L^2(\Omega).$$

Thus, the homogenized equation of (4.10) is the zero equation, which implies that  $\theta' = 0$ .  $\square$

**Remark 4.2.** In the above lemma, we conclude that  $\theta' = 0$  which cannot be done, in general. This argument is also valid in Lemma 3.2. However, there the estimate on the optimal controls was enough to conclude this, which is not available in the present situation.

**Remark 4.3.** The above lemma is the counterpart of Lemma 3.2 for the problem considered in this section. In contrast to the problem in Section 3, here we do not have a *a priori* bound on  $\tilde{\theta}_\varepsilon^*$  even in  $H^{-1}(\Omega)$  (as in Lemma 3.2). Hence, we also do not have an estimate of  $u_\varepsilon^*$  in  $V_\varepsilon$ . Hence the homogenization of the state equation is not achieved as in Theorem 3.3. In other words, one is unable to address this problem for an arbitrary admissible set  $U_\varepsilon$ , in general. However, in the next theorem we shall study the positive cone case which will offer some estimates on the optimal controls. Also, in the above lemma, we have homogenized the adjoint-state equation (as in Theorem 3.3). In the next theorem we homogenize the state equation, as well, for the positive cone case.

Let  $U_\varepsilon = \{\theta \in L^2(\Omega) \mid \theta \geq 0 \text{ a.e. in } \Omega_\varepsilon\}$  be the closed convex subset of  $L^2(\Omega)$ . The elements of  $U_\varepsilon$  restricted to  $\Omega_\varepsilon$  is the set of all positive elements of  $L^2(\Omega_\varepsilon)$ . Let  $U'$  be the sequential  $K$ -lower limit of  $U_\varepsilon$  with respect to the strong topology in  $L^2(\Omega)$ .  $U'$  is non-empty. Let  $V'$  be the strong closure of  $U'$  in  $H^{-1}(\Omega)$ . Let  $U$  denote the positive cone of  $L^2(\Omega)$ . Let  $V$  be the strong closure of  $U$  in  $H^{-1}(\Omega)$ .

**Theorem 4.4.** Let  $U_\varepsilon = \{\theta \in L^2(\Omega) \mid \theta \geq 0 \text{ a.e. in } \Omega_\varepsilon\}$ . Then  $\{u_\varepsilon^*\}$  and  $\{\tilde{\theta}_\varepsilon^*\}$  are bounded in  $V_\varepsilon$  and  $H^{-1}(\Omega)$ , respectively, and there exists  $\theta^* \in H^{-1}(\Omega)$  such that

$$\tilde{\theta}_\varepsilon^* \rightharpoonup \theta^* \text{ weakly in } H^{-1}(\Omega). \tag{4.11}$$

Also,  $u_0(x, y) = \chi(y)u^*(x)$  where  $u^* \in H_0^1(\Omega)$  solves

$$\begin{cases} -\operatorname{div}(A_0 \nabla u^*) = \chi_0 f + \theta^* & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases} \tag{4.12}$$

Moreover,  $p^* \in U$  and  $\langle \theta^*, p^* \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} = 0$ . Further,  $\theta^*$  minimises

$$J(\theta) = \frac{1}{2} \int_{\Omega} \chi_0 u^2 dx$$

over  $V'$ , where  $u = u(\theta)$  solves (3.21) and

$$J_\varepsilon(\theta_\varepsilon^*) \rightarrow J(\theta^*). \tag{4.13}$$

**Proof.** Observe that  $U_\varepsilon$  satisfies the hypothesis (H1), since  $0 \in U_\varepsilon$  for all  $\varepsilon$ . Therefore the results of Lemma 4.1 remain valid. Also, the optimality condition (3.4) implies that  $\varepsilon \theta_\varepsilon^* = (p_\varepsilon^*)^-$  and thus (3.22) holds. An argument similar to the one in Theorem 3.7 will show that  $p^* \in U$ .

Using  $u_\varepsilon^*$  as a test function in the weak form of the state equation satisfied by  $u_\varepsilon^*$  (cf. (3.12)), we have

$$\int_{\Omega_\varepsilon} A\left(x, \frac{x}{\varepsilon}\right) \nabla u_\varepsilon^* \cdot \nabla u_\varepsilon^* dx = \int_{\Omega_\varepsilon} (f + \theta_\varepsilon^*) u_\varepsilon^* dx = \int_{\Omega_\varepsilon} f u_\varepsilon^* dx + \varepsilon^{-1} \int_{\Omega_\varepsilon} (p_\varepsilon^*)^- u_\varepsilon^* dx.$$

Now, using  $(p_\varepsilon^*)^-$  as a test function in the weak form of the adjoint equation (4.2), we have

$$\int_{\Omega_\varepsilon} (p_\varepsilon^*)^- u_\varepsilon^* dx = \int_{\Omega_\varepsilon} {}^t A\left(x, \frac{x}{\varepsilon}\right) \nabla (p_\varepsilon^*)^- \cdot \nabla p_\varepsilon^* dx = - \int_{\Omega_\varepsilon} {}^t A\left(x, \frac{x}{\varepsilon}\right) \nabla (p_\varepsilon^*)^- \cdot \nabla (p_\varepsilon^*)^- dx$$

and hence we derive the equality,

$$\int_{\Omega_\varepsilon} A\left(x, \frac{x}{\varepsilon}\right) \nabla u_\varepsilon^* \cdot \nabla u_\varepsilon^* dx + \varepsilon^{-1} \int_{\Omega_\varepsilon} A\left(x, \frac{x}{\varepsilon}\right) \nabla (p_\varepsilon^*)^- \cdot \nabla (p_\varepsilon^*)^- dx = \int_{\Omega_\varepsilon} f u_\varepsilon^* dx. \tag{4.14}$$

Since  $\{u_\varepsilon^*\}$  is bounded in  $L^2(\Omega_\varepsilon)$ , we deduce from (4.14) that  $\{u_\varepsilon^*\}$  and  $\{\varepsilon^{-1/2}(p_\varepsilon^*)^-\}$  are bounded in  $V_\varepsilon$ . Therefore, there exist  $u^* \in H_0^1(\Omega)$  and  $u_1 \in L^2[\Omega; H_{\text{per}}^1(Y)/\mathbb{R}]$  such that  $u_0(x, y) = \chi(y)u^*(x)$  and, up to a subsequence,

$$\widetilde{\nabla u_\varepsilon^*} \xrightarrow{2s} \chi(y)[\nabla u^*(x) + \nabla_y u_1(x, y)].$$

Now, arguing as in Lemma 3.2, we show that  $\{\tilde{\theta}_\varepsilon^*\}$  is bounded in  $H^{-1}(\Omega)$  and thus there exists  $\theta^* \in H^{-1}(\Omega)$  such that (4.11) holds, for a subsequence. Now, arguing as in the proof of Theorem 3.3, we show that  $u^*$  solves (4.12).

Moreover, we have (cf. [2, Theorem A.1])

$$\left. \begin{aligned} \|u_\varepsilon^*\|_{2, \Omega_\varepsilon}^2 &\rightarrow \chi_0 \|u^*\|_2^2, \\ \|\varepsilon^{-1/2}(p_\varepsilon^*)^-\|_{2, \Omega_\varepsilon} &= \|\varepsilon^{1/2}\theta_\varepsilon^*\|_{2, \Omega_\varepsilon} \rightarrow 0. \end{aligned} \right\}$$

Thus, passing to the limit in  $J_\varepsilon(\theta_\varepsilon^*)$  we have (4.13).

On the other hand, using  $p_\varepsilon^*$  as a test function in (3.12) and  $u_\varepsilon^*$  as a test function in (3.3), one can rewrite the cost functional  $J_\varepsilon(\theta_\varepsilon^*)$  as in (3.18). Now, using (3.22),  $J_\varepsilon(\theta_\varepsilon^*)$  reduces to

$$J_\varepsilon(\theta_\varepsilon^*) = \frac{1}{2} \int_{\Omega_\varepsilon} f p_\varepsilon^* dx$$

and passing to the limit will yield

$$J_\varepsilon(\theta_\varepsilon^*) \rightarrow \frac{1}{2} \int_{\Omega} \chi_0 f p^* dx.$$

Using  $p^*$  and  $u^*$  as test functions in (4.12) and (4.7), respectively, we have

$$\int_{\Omega} \chi_0 f p^* dx = J(\theta^*) - \frac{1}{2} \langle \theta^*, p^* \rangle_{H^{-1}(\Omega), H_0^1(\Omega)}.$$

Therefore, (4.13) implies that  $\langle \theta^*, p^* \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} = 0$ . Now an argument analogous to the one in the proof of Theorem 3.7 will show that  $\theta^*$  minimises  $J$  in  $V'$ . The strict convexity of  $J$  would then imply the uniqueness of  $\theta^*$  and thus the convergences (4.11) and (4.3) hold for the entire sequence.  $\square$

**Remark 4.5.** The above theorem differs from Theorem 3.7 in the sense that we now have the fact that

$$\int_{\Omega_\varepsilon} \theta_\varepsilon^* p_\varepsilon^* dx = -\varepsilon \|\theta_\varepsilon^*\|_{2, \Omega_\varepsilon}^2 \rightarrow 0.$$

In other words, we have identified the limit of the product of the two weakly converging sequence. This was possible due to (4.14) and similar equality is not available for the problem in Section 3.

## 5. Conclusion

We have studied the asymptotic behaviour of some low-cost control problems left open in [12,19]. The results are proved for perforated domains but they remain valid for non-perforated domains as well, with necessary modification. We prove an  $H$ -convergence result for weak data (cf. Theorem 2.1) which forms the backbone of the article. It would be interesting to see if the difficulties discussed in Remark 3.8 could be overcome.

## Acknowledgments

The authors wish to thank the University Grants Commission (UGC), India for their support. The first author would like to thank the National Board of Higher Mathematics (NBHM), India for their financial support. This work was also partially supported by the CEFIPRA project No. 3701-1.

## References

- [1] G. Allaire, Homogenization and two-scale convergence, *SIAM J. Math. Anal.* 23 (1992) 1482–1518.
- [2] G. Allaire, F. Murat, Homogenization of the Neumann problem with non-isolated holes, *Asymptot. Anal.* 7 (1993) 81–95, with an appendix jointly written with A.K. Nandakumaran.
- [3] D. Cioranescu, P. Donato, An Introduction to Homogenization, Oxford Lecture Ser. Math. Appl., vol. 17, Oxford University Press, 1999.
- [4] D. Cioranescu, J. Saint Jean Paulin, Homogenization in open sets with holes, *J. Math. Anal. Appl.* 71 (1979) 590–607.
- [5] G. Dal Maso, An Introduction to  $\Gamma$ -Convergence, *Progr. Nonlinear Differential Equations Appl.*, vol. 8, Birkhäuser, Boston, 1993.
- [6] G. Dal Maso, F. Murat, Asymptotic behaviour and correctors for linear Dirichlet problems with simultaneously varying operators and domains, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 21 (2004) 445–486.
- [7] S. Kesavan, T. Muthukumar, Homogenization of an optimal control problem with state-constraints, *Differential Equations Dynam. Systems*, in press.
- [8] S. Kesavan, T. Muthukumar, Low-cost control problems on perforated and non-perforated domains, *Proc. Indian Acad. Sci. Math. Sci.* 118 (2008) 133–157.
- [9] S. Kesavan, M. Rajesh, On the limit matrix obtained in the homogenization of an optimal control problem, *Proc. Indian Acad. Sci. Math. Sci.* 112 (2002) 337–346.
- [10] S. Kesavan, J. Saint Jean Paulin, Homogenization of an optimal control problem, *SIAM J. Control Optim.* 35 (1997) 1557–1573.
- [11] S. Kesavan, J. Saint Jean Paulin, Optimal control on perforated domains, *J. Math. Anal. Appl.* 229 (1999) 563–586.
- [12] S. Kesavan, J. Saint Jean Paulin, Low cost control problems, *Trends Indust. Appl. Math.* (2002) 251–274.
- [13] S. Kesavan, M. Vanninathan, L'homogénéisation d'un problème de contrôle optimal, *C. R. Acad. Sci. Paris Ser. A* 285 (1977) 441–444.
- [14] J.L. Lions, Remarks on "cheap" control, in: J.J.H. Miller (Ed.), *Topics in Numerical Analysis*, Academic Press Inc., ISBN 0-12-496950-X, 1973, pp. 203–209.
- [15] J.L. Lions, Some Methods in the Mathematical Analysis of Systems and Their Control, Gordon and Breach, Science Publishers, Inc., ISBN 0-677-60200-6, 1981.
- [16] D. Lukkassen, G.Nguetseng, P. Wall, Two-scale convergence, *Int. J. Pure Appl. Math.* 2 (2002) 35–86.
- [17] F. Murat,  $H$ -convergence, in: *Séminaire d'Analyse Fonctionnelle et Numérique de l'Université d'Alger*, March 1978, mimeographed notes. English translation in [18].

- [18] F. Murat, L. Tartar, *H*-convergence, in: A. Cherkaev, R. Kohn (Eds.), *Topics in the Mathematical Modelling of Composite Materials*, in: *Progr. Nonlinear Differential Equations Appl.*, vol. 31, Birkhäuser, Boston, 1997, pp. 21–43. English translation of [17,23].
- [19] T. Muthukumar, *Asymptotic behaviour of some optimal control problems*, PhD thesis, University of Madras, Chennai, Institute of Mathematical Sciences, Chennai, India, May 2006.
- [20] A.K. Nandakumar, *Steady and evolution Stokes equations in a porous media with nonhomogeneous boundary data: A homogenization process*, *Differential Integral Equations* 5 (1992) 79–93.
- [21] G. Nguetseng, *A general convergence result for a functional related to the theory of homogenization*, *SIAM J. Math. Anal.* 20 (1989) 608–623.
- [22] J. Saint Jean Paulin, H. Zoubairi, *Optimal control and “strange term” for a Stokes problem in perforated domains*, *Port. Math.* 59 (2002) 161–178.
- [23] L. Tartar, *Cours Peccot au Collège de France*, partially written in [17]. English translation in [18], March 1977.