



Candidate Gene Expression in *Bos indicus* Ovarian Tissues: Prepubertal and Postpubertal Heifers in Diestrus

Mayara Morena Del Cambre Amaral Weller^{1,2}, Marina Rufino S. Fortes^{3*}, Laercio R. Porto-Neto⁴, Matthew Kelly^{2†}, Bronwyn Venus², Lisa Kidd⁵, João Paulo Arcelino do Rego⁶, Sophia Edwards^{5†}, Gry B. Boe-Hansen⁵, Emily Piper^{5†}, Sigrid A. Lehnert⁴, Simone Eliza Facioni Guimarães¹ and Stephen Stewart Moore²

OPEN ACCESS

Edited by:

Ramanathan K. Kasimanickam,
Washington State University, USA

Reviewed by:

Adriana Santana Carmo,
Deoxi Biotecnologia, Brazil
Sherrie Gayle Clark-Deener,
Virginia Tech, USA

*Correspondence:

Marina Rufino S. Fortes
m.fortes@uq.edu.au

†Present address:

Matthew Kelly,
The Australian Agricultural Company,
Brisbane, QLD, Australia
Sophia Edwards,
Vetoquinol Australia Pty Ltd,
Brisbane, QLD, Australia
Emily Piper,
Zoetis Inc. Pharmaceuticals,
Kalamazoo, MI, USA

Specialty section:

This article was submitted to Animal
Reproduction - Theriogenology,
a section of the journal
Frontiers in Veterinary Science

Received: 25 July 2016

Accepted: 04 October 2016

Published: 18 October 2016

Citation:

Weller MMDCA, Fortes MRS,
Porto-Neto LR, Kelly M, Venus B,
Kidd L, do Rego JPA, Edwards S,
Boe-Hansen GB, Piper E,
Lehnert SA, Guimarães SEF and
Moore SS (2016) Candidate Gene
Expression in *Bos indicus* Ovarian
Tissues: Prepubertal and
Postpubertal Heifers in Diestrus.
Front. Vet. Sci. 3:94.
doi: 10.3389/fvets.2016.00094

¹Animal Science Department, Universidade Federal de Viçosa, Viçosa, Minas Gerais, Brazil, ²Queensland Alliance for Agriculture and Food Innovation, Centre for Animal Science, The University of Queensland, Brisbane, QLD, Australia, ³School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, QLD, Australia, ⁴CSIRO Agriculture Flagship, Queensland Bioscience Precinct, Brisbane, QLD, Australia, ⁵School of Veterinary Sciences, The University of Queensland, Gatton, QLD, Australia, ⁶Instituto Federal de Educação, Ciência e Tecnologia do Ceará, Fortaleza, Ceará, Brazil

Growth factors such as bone morphogenetic proteins 6, 7, 15, and two isoforms of transforming growth factor-beta (BMP6, BMP7, BMP15, TGFB1, and TGFB2), and insulin-like growth factor system act as local regulators of ovarian follicular development. To elucidate if these factors as well as others candidate genes, such as estrogen receptor 1 (*ESR1*), growth differentiation factor 9 (*GDF9*), follicle-stimulating hormone receptor (*FSHR*), luteinizing hormone receptor (*LHR*), bone morphogenetic protein receptor, type 2 (*BMPR2*), type 1 insulin-like growth factor receptor (*IGFR1*), and key steroidogenic enzymes cytochrome P450 aromatase and 3- β -hydroxysteroid dehydrogenase (*CYP19A1* and *HSD3B1*) could modulate or influence diestrus on the onset of puberty in Brahman heifers, their ovarian mRNA expression was measured before and after puberty (luteal phase). Six postpubertal (POST) heifers were euthanized on the luteal phase of their second cycle, confirmed by corpus luteum observation, and six prepubertal (PRE) heifers were euthanized in the same day. Quantitative real-time PCR analysis showed that the expression of *FSHR*, *BMP7*, *CYP19A1*, *IGF1*, and *IGFR1* mRNA was greater in PRE heifers, when contrasted to POST heifers. The expression of *LHR* and *HSD3B1* was lower in PRE heifers. Differential expression of ovarian genes could be associated with changes in follicular dynamics and different cell populations that have emerged as consequence of puberty and the luteal phase. The emerging hypothesis is that *BMP7* and *IGF1* are co-expressed and may modulate the expression of *FSHR*, *LHR* and *IGFR1*, and *CYP19A1*. *BMP7* could influence the downregulation of *LHR* and upregulation of *FSHR* and *CYP19A1*, which mediates the follicular dynamics in heifer ovaries. Upregulation of *IGF1* expression prepuberty, compared to postpuberty diestrus, correlates with increased levels *FSHR* and *CYP19A1*. Thus, *BMP7* and *IGF1* may play synergic roles and were predicted to interact, from the expression data ($P = 0.07$, $r = 0.84$). The role of these co-expressed genes in puberty and heifers luteal phase merits further research.

Keywords: bovine species, intraovarian factors, quantitative real-time PCR, corpus luteum, folliculogenesis

INTRODUCTION

Ovarian activity and hormones are paramount for pubertal development and normal reproductive performance (1). Improving the reproductive performance of *Bos indicus* cattle is an industry priority in tropical and subtropical regions of the world, because of its impact on farm productivity (2). *B. indicus* breeds have a later onset of puberty (16–40 months of age), which has a negative impact on their overall reproductive performance (3–6). Although the heritability of age at puberty measured by first detected corpus luteum (CL) has been reported to be moderate (0.52–0.57) (7), the phenotypic identification of animals that undergo puberty at an early age is expensive. Enhancing our comprehension of ovarian genes and their interactions involved in bovine puberty could have practical implications in the animal breeding context.

The regulation of ovarian activity is an integrated process that involves FSH and LH, their receptors, ovarian steroids, and intraovarian factors (8). Some of the most important intraovarian factors are members of the transforming growth factor-beta (*TGFB*) superfamily, such as bone morphogenetic proteins (BMPs) 6, 7, and 15 (*BMP6*, *BMP7*, and *BMP15*) and two isoforms of *TGFB* (*TGFB1* and *TGFB2*), which are expressed by ovarian somatic cells and oocytes in a stage-specific manner throughout folliculogenesis (9, 10). These genes function as local regulators of ovarian follicular development and subsequently affect fertility (11–13). Experiments in mice indicate that the different isoforms of *TGFB* are responsible for diverse physiological functions (14, 15). An *in vivo* study showed that *BMP7* promotes the “recruitment” of primordial follicles into the growing follicle pool while inhibiting progesterone production and ovulation (16). Similar to *BMP7*, *BMP15*, and *BMP6* are part of a group of luteinization inhibitors (17). *In vitro*, *BMP7* induced expression of follicle-stimulating hormone receptor (*FSHR*) mRNA in human granulosa cells (18). In contrast, *BMP15* suppressed *FSHR* and luteinizing hormone receptor (*LHR*) expression (19). Given the discussed roles affecting ovulation, these intraovarian factors could modulate or influence the onset of puberty that leads to the first ovulation event.

Growth differentiation factor 9 (*GDF9*) gene expression is relevant for oocyte competence, and *GDF9* follows a similar expression pattern to *BMP15* in cows stimulated with FSH treatment (20). The relevance of *GDF9* expression in ovarian tissues of peripubertal *B. indicus* heifers is unclear, but its link to *BMP15* and FSH pathways merits investigation.

The insulin-like growth factors (IGFs) system plays a key role in follicular development and female fertility (21–23). Insulin-like growth factor 1 (*IGF1*) and insulin-like growth factor 2 (*IGF2*) have been reported to act in synergy with gonadotropins *LH* and *FSH* to stimulate growth and differentiation of ovarian follicles and subsequent synthesis and secretion of estradiol and progesterone production (24–27). Moreover, experiments in cattle (28, 29) showed that *IGF2* was the main intrafollicular IGF ligand regulating follicular growth and highly expressed in theca cells. All of these previous reports suggest that the IGFs and *TGFB* superfamily genes may have different roles in ovarian activity, related to the cycle phase and possible to endocrine regulation of puberty.

Estrogen produced by ovarian tissue is relevant for GnRH release from the hypothalamus and the expression of its receptor, *ESR1*, has been associated with puberty in mice (30). In cattle, *ESR1* expression is required for normal follicular development and follicular dominance (31). Given the relevance of steroid hormones for pubertal development and ovarian function, key steroidogenic enzymes such as cytochrome P450 aromatase (*CYP19A1*) and 3- β -hydroxysteroid dehydrogenase (*HSD3B1*) were also targeted in this study.

So far, studies have focused on the endocrine regulation of the hypothalamic–pituitary–ovarian system for onset of puberty (32–34). Information about specific changes in ovarian gene expression-associated specific cycle phases and puberty is sparse. Thus, the aim of the present study was to elucidate the expression pattern of candidate genes, such as *ESR1*, *GDF9*, *FSHR*, *LHR*, *BMPR2*, *TGFB1*, *TGFB2*, *BMP15*, *BMP6*, *BMP7*, *IGF1*, *IGF2*, *CYP19A1*, and *HSD3B1* in pre- (PRE) and postpubertal (POST) heifers in the diestrus phase. Evidence of differential expression will help to support or disprove the hypothesis that these genes modulate or are influenced by the presence of circulating progesterone and the onset of puberty.

MATERIALS AND METHODS

Animal Management and Puberty Observation

Management, handling, and euthanasia of animals were approved by the Animal Ethics Committee of The University of Queensland, Production and Companion Animal group (certificate number QAAFI/279/12). Twenty Brahman heifers, which were not pedigree animals, but had a characteristic *B. indicus* phenotype and were typical beef industry animals, were sourced as young weaners born at the same season (<250 kg) and kept at grazing conditions, from two commercial herds in Queensland, Australia. After being sourced, heifers were kept at the Gatton Campus facilities of the University of Queensland; they were all under the same conditions and pasture based diet until project end. Precise day of birth information was not available for these heifers as they were sourced from industry; the effect of age differences in pubertal development is possible but could not be tested in this experiment.

Heifers were examined every 2 weeks for observation of pubertal development, from October 2012 to May 2013. Ovarian activity was observed using ultrasonography [HS-2000(VET), Honda Electronics Inc.]. Pubertal status was defined by presence of a CL observed with the ultrasound (7). Euthanasia plans were based on date of the first CL observation. Six heifers were chosen to be prepubertal (PRE) and six heifer POST. The POST heifer when identified was then paired with PRE heifer which was randomly drawn from the remaining animals and processed on the same day. The animals were weighted, and the body condition (BCS) was measured. POST heifers were euthanized 23 days, on average, after observation of the first CL. POST heifers were euthanized on the luteal phase of their second estrous cycle, confirmed by CL presence on ovarian tissue post euthanasia. Euthanasia was carried out by stunning with a non-penetrating

captive bolt followed by exsanguination. Concentrations of progesterone were measured with a radioimmunoassay (RIA) from blood samples collected at exsanguination, to confirm that POST heifers had a functional CL, which was observed at euthanasia. RIA was performed by the Laboratory for Animal Endocrinology of the University of Queensland (Dr. Stephen Anderson). Plasma progesterone concentrations were measured by RIA as described by J. D. Curlew, M. Axelson, and G. M. Stone (35) with the difference that progesterone antiserum C-9817 (Bioquest, North Ryde, NSW, Australia) was used. Extraction efficiency was 75%, and the values reported herein were not corrected for these losses. The sensitivity of the assay was 0.1 ng/ml, and the intra- and inter-assay coefficient of variation (CV) was 5.0%.

Post-euthanasia, left and right ovaries were harvested and preserved by snap freezing in liquid nitrogen, then kept at -80°C until RNA extractions were carried out. In total, 24 ovaries (2×6 PRE and 2×6 POST heifers) were processed separately for RNA extraction and quantitative real-time PCR (qRT-PCR) measurements.

RNA Extraction and Quantitative Real-Time Transcription PCR Analysis

Prior to RNA extraction, the whole ovary tissue was ground under liquid nitrogen to form a powder of which 25 mg was used for RNA extractions; left and right ovaries were kept separately. Total RNA was isolated separately from 25 mg of the homogenized sample of left and right ovaries from PRE and POST heifers, using Trizol method (Life Technologies, Inc.). The total RNA was resuspended in RNase-free ultrapure water and stored at -80°C until further use. RNA concentrations were measured by Nanodrop ND-1000 spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA) with an optimal 260/280 ratio between 1.8 and 2.1. Intact 28S and 18S rRNA subunit integrity was assessed by Bioanalyser (RIN 6.9 or above for all samples).

Reverse transcription was performed using GoScript Reverse Transcription System (Promega) and oligo (dT) primers (Invitrogen). The reactions were performed with 6 μg of total RNA and 2 μl of 50 μM oligo (dT)23VN primer, following the manufacturer's recommended protocol. The cDNA concentrations from the samples were estimated on a Nanodrop ND-1000 spectrophotometer (Thermo Fisher Scientific). Finally, the single-stranded cDNA samples were stored at -20°C until analysis by qRT-PCR.

Quantitative real-time PCR reactions were performed in triplicate using SYBR[®] Select Master Mix (Applied Biosystem) following the manufacturer's instructions in a ViiA[™] 7 Real-Time PCR System (Applied Biosystem). The CV of Ct values from replicates within each sample was low <3% indicating acceptable accuracy and reproducibility (not shown). The oligonucleotide primers used for the reactions were designed using PrimerQuest software provided by Integrated DNA Technologies, Inc. from *Bos taurus* sequences available in GeneBank database. In the present study, glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*) was used as an internal control gene because it was stably expressed in our study and previously in dairy cattle (36). The list of primer sequences used for each target gene is listed in **Table 1**.

Prior to performing qRT-PCR, the amplification efficiency and optimal primer concentration were determined for each gene using serial dilution of cDNA. For this purpose, four concentrations of cDNA (0.625, 2.5, 10, and 40 ng/reaction) and two primer dilutions (100, 200 nM) were tested. The PCR efficiencies for all primers pairs were obtained using the formula $E = 10^{(-1/\text{slope})} \times 100$, where E is efficiency and slope is the gradient of dilution series in the linear phase. These results are summarized in **Table 1**, where E is represented as a percentage [$\%E = (E - 1) \times 100$]. Samples were amplified separately in triplicate using a ViiA[™] 7 Real-Time PCR System (Applied Biosystem, Foster City, CA, USA) with the following amplification program: an initial step at 95°C for 10 min, a second step of 40 cycles of 95°C for 20 s, and a final extension step at 60°C for 30 s. At the end of each reaction, a denaturation curve was plotted to assure that each reaction produced a single fragment, that is, the curve contained only one dissociation peak.

Statistical Analysis

Progesterone, body weight, and condition score data were analyzed in a completed randomized design using SAS (version 9.1.3), and means were compared by the t -test to the level of 5%. Statistical analysis of Ct data was performed using %QPCR_MIXED macro developed in SAS (version 9.1.3) (https://msu.edu/~steibelj/JP_files/QPCR.html) developed to generate codes in SAS PROC MIXED suitable to analyze data from qRT-PCR, assuming independent random effects for reference and target genes in each biological replicate (37). The following model was used:

$$y_{gikr} = \text{TG}_{gi} + C_{gik} + D_{ik} + e_{gikr}$$

where, y_{gikr} corresponds to the Ct value obtained from the thermocycler software for the g th gene (reference or targets) from the r th well, which corresponds to the k th animal in i th physiological state (PRE or POST); TG_{gi} is the effect of the i th physiological state on the expression of gene g ; $C_{gik} \sim N(0, \sigma_c^2)$ is the gene-specific random effect of the k th animal; $D_{ik} \sim N(0, \sigma_d^2)$ is the sample-specific random effect (common to reference and target genes); and $e_{gikr} \sim N(0, \sigma_e^2)$ is the residual term.

The relative expression was estimated using ΔCt method (target gene Ct – GAPDH gene Ct) as previously reported (38), where Ct is the PCR cycle number at which the fluorescence generated within a reaction crosses an arbitrary threshold. For each target gene, the comparison of gene expression between physiological state (PRE or POST) was performed by CONTRAST statement of the GLM procedure (SAS software) using Student's t -test to the level of 5%. The "estimates" generated by CONTRAST analyses were used to estimate fold-change (relative expression) for pair-wise contrast of interest and were obtained by using $2^{-(\text{Estimate})}$. The contrast between PRE versus POST heifers (both ovaries) was analyzed considering the average of Ct values from right and left ovary of each heifer for each gene. The other two contrasts were: (1) both ovaries from PRE versus only ovaries with CL from POST heifers and (2) ovaries from POST heifers without CL versus ovaries from POST heifers with CL. These two other contrasts should help to elucidate if differences in gene expression observe are related to the local presence of the

TABLE 1 | Gene names, primer pair sequences, annealing temperatures, and amplification efficiencies of each target.

Gene ^a	GenBank accession no.	Primer sequence (5'–3')	Amplicon (bp)	Efficiency (%)
<i>ESR1</i>	NM_001001443.1	F-CTCTCTGCCTTTGCTACCTTAC R-CCC GGTCACAAATAGCTAAA	114	98.87
<i>FSHR</i>	NM_174061.1	F-CATGCTCATCTTCACCGACTT R-GACCAGGAGGATCTTTGACTTG	112	99.64
<i>LHR</i>	NM_174381.1	F-GAGTGGCTGGGATTATGACTATG R-CAGCCAAATCAGGACTCTAAGG	127	99.87
<i>BMPR2</i>	NM_001304285.1	F-GTACCGGCATGACCACTATATC R-AGTCTTCTTCGGTCAGTTGTAAG	120	99.74
<i>TGFB1</i>	NM_001166068.1	F-TGCTTCAGCTCCACAGAAA R-GTATCCAGGCTCCAGATGTAAG	149	99.74
<i>TGFB2</i>	NM_001113252.1	F-CACGAATGGCTCCACCATAA R-AGCGTGCTTCTAGTTCTTCCAC	127	98.67
<i>GDF9</i>	NM_174681.2	F-GCATTCCCTCCACCCTAAA R-GGTGACGGGACAATCTTACA	113	99.89
<i>BMP15</i>	NM_001031752.1	F-GTAGTGAGGTTTCGTGAGTTCTG R-TAGGGAGAGGTTTGGTCTTCT	111	98.85
<i>BMP6</i>	XM_005223892.2	F-TTCTCAACGACGCCGATATG R-ACCCTCAGGAATCTGGGATAG	122	98.56
<i>BMP7</i>	NM_001206015.1	F-ATGACAGCTCCAACGTCATC R-AGAGACCCAGGATCCAGAAA	123	98.99
<i>CYP19A1</i>	NM_174305.1	F-GTGTCGGAAGTTGTGCCTAT R-GACCTGGTATTGAGGATGTGTC	101	98.95
<i>HSD3B1</i>	NM_174343.3	F-ACACCAGCACCATAGAAGTG R-CTTGCTGTATGGGTATGGAGAG	113	98.88
<i>IGF1</i>	NM_001077828.1	F-TCCCATCTCCCTGGATTTCT R-GGGTTGGAAGACTGCTGATT	105	98.86
<i>IGFR1</i>	NM_001244612.1	F-GTATGGAGGAGCCAAGCTAAA R-GTCTTGGCCTGAACGTAGAA	123	98.98
<i>IGF2</i>	NM_174087.3	F-CATTGTGGGAAGGTGTGTC R-TTACATTCTGTGGGCTGTGG	113	99.02
<i>B2M</i>	NM_173893.3	F-ACCTGAAGTCTATGTGTATGG R-GTGGGACAGCAGGTAGAAAAG	134	99.95
<i>GAPDH</i>	NM_001034034.2	F-GATGCTGGTGCTGAGTATGT R-GCAGAAGGTGCAGAGATGAT	113	99.96

^aGene symbol: *ESR1*, estrogen receptor α ; *FSHR*, follicle-stimulating hormone receptor; *LHR*, luteinizing hormone receptor; *BMPR2*, bone morphogenetic protein receptor, type 2; *TGF β 1*, transforming growth factor beta 1; *TGF β 2*, transforming growth factor beta 2; *GDF9*, growth differentiation factor 9; *BMP15*, bone morphogenetic protein 15; *BMP6*, bone morphogenetic protein 6; *BMP7*, bone morphogenetic protein 7; *CYP19A1*, cytochrome P450 aromatase; *HSD3B1*, 3- β -hydroxysteroid dehydrogenase; *IGF1*, insulin-like growth factor 1; *IGFR1*, type 1 insulin-like growth factor receptor; *IGF2*, insulin-like growth factor 2; *B2M*, beta-2-microglobulin; *GAPDH*, glyceraldehyde-3-phosphate dehydrogenase.

CL itself or not (are more generally related to PRE versus POST processes). Once the efficiency (E) of the qRT-PCR reaction was close to 100%, one PCR cycle of difference between two samples means twice as much expression in the first sample in comparison with the second.

Pair-wise correlations between gene expression values were calculated, and its significance was tested using R software for analyses. Significant correlations were interpreted as predicted gene interactions.

RESULTS

Serum Progesterone Concentrations, Body Weight, and Condition


Average serum concentration of progesterone for the PRE and POST *B. indicus* Brahman heifers were 0.4 ± 0.2 and 2.0 ± 0.7 ng/ml, respectively ($n = 6$ per group). The difference in progesterone concentration corresponds to the observation of CL in POST heifers. Associated data collected prior to euthanasia of PRE and POST heifers indicated no significant difference in body weight

or body condition score between PRE and POST heifers. Body weight averages were 338 ± 54.17 and 362.6 ± 38.62 kg ($P = 0.38$), and condition score averages were 3.5 ± 0.44 and 3.75 ± 0.41 ($P = 0.18$), respectively.

Ovarian Gene Expression

Ovarian gene expression was measured with real-time quantitative PCR in three contrasts: (1) average gene expression from both ovaries PRE versus POST heifers in diestrus; (2) average expression of PRE ovaries versus only ovaries with a CL in POST heifers in diestrus; and (3) ovaries with CL versus ovaries without using only POST heifers in diestrus.

In the contrast between PRE versus POST heifers (both ovaries), *FSHR* ($P < 0.05$), *BMP7* ($P < 0.05$), *IGF1* ($P < 0.05$), *TGFB1* ($P < 0.01$), and *IGFR1* ($P < 0.01$) expression were greater in PRE heifers, whereas *LHR* ($P < 0.05$) expression was lower in PRE heifers (**Figure 1**). The abundance of key steroidogenic enzyme genes, such as *CYP19A1* ($P < 0.05$) was greater in PRE heifers, whereas *HSD3B1* mRNA expression was greater ($P < 0.01$) in POST heifers.



In the contrast between both ovaries from PRE versus only ovaries with CL from POST heifers, *FSHR* ($P < 0.05$), *CYP19A1* ($P < 0.05$), and *TGFB1* ($P < 0.01$) expression was greater in PRE heifers, whereas expression of *LHR* ($P < 0.05$), and *HSD3B1* ($P < 0.01$) were greater in POST heifers' ovaries with CL (Figure 2).

In the third and last contrast, we did not observe any differential gene expression between ovaries with a CL versus the contralateral ovary (without CL) from POST heifers (not shown).

Pair-wise correlations predicted four gene-gene interactions from co-expression data. The predicted significant correlations were between: *LHR* and *HSD3B1* ($P < 0.01$, $r = -0.97$); and *HSD3B1* and *CYP19A1* ($P = 0.03$, $r = -0.90$). Correlations between *IGF1* and *BMP7* ($P = 0.07$, $r = 0.84$); *IGF1* and *IGFR1* ($P = 0.10$, $r = 0.79$) were high ($r > 0.79$) but not statistically significant ($P > 0.05$ and < 0.10). Other pair-wise correlations between gene expression values were low and not significant ($P > 0.1$).


DISCUSSION

This study has demonstrated that the diestrus phase of pubertal heifers is associated with changes in ovarian gene expression of gonadotropin receptors (*FSHR* and *LHR*), key steroidogenic enzymes (*HSD3B1* and *CYP19A1*) and important intraovarian factors and their receptor, such as *BMP7*, *TGFB1*, and *IGFR1*. The regulation of ovarian activity is determined by coordinated action between FSH and LH with their receptors, ovarian steroids, and intraovarian factors (8). Genes of the TGF β superfamily, such as *GDF9*, *BMP6*, *BMP7*, *BMP15*, *TGFB1*, and *TGFB2*, and their cell receptors are known paracrine and

autocrine modulators of ovarian function and fertility (9–13). The IGF system plays a key role in follicular development and is an additional important intraovarian factor involved in pubertal development (24–27, 39). Our results fit with current theory and point to specific intraovarian factors relevant to the diestrus phase of pubertal heifers in *B. indicus*.

We did not detect differential gene expression between ovaries from POST heifers with a CL versus the contralateral ovary (without a CL), which could confirm the systemic effect of circulating progesterone on ovarian dynamics. Therefore, we focused on the differences between PRE and POST heifers. In the present study, the significant differences in ovarian gene expression between two physiological states, PRE and POST (luteal phase), are likely related to onset of puberty and influenced by progesterone signaling. It is important to note that the comparison is between PRE heifers, which had never experienced a luteal phase and the related progesterone signaling, while the POST heifers were in their second diestrus with the corresponding levels of progesterone being produced by CL. Future research examining the other phases of the cycle in POST heifers would complement our findings.

We identified greater expression of *FSHR* in the ovaries of PRE than POST heifers in diestrus, which could be related to follicular waves characterized by synchronous development of groups of growing follicles, one of which would become dominant and achieved the greatest diameter, suppressing the growth of the other smaller subordinate follicles. Transrectal ultrasonography has detected the sequential growth and regression of large follicles in heifers, near the time of first ovulation (40). FSH and its receptor *FSHR* play an important role in follicle progression from



the primary to the advanced stage of follicular development (41). Studies reported the expression of *FSHR* on granulosa cells in primary and secondary follicles, which is evidence for the role of FSH in follicular development at early stages (41, 42). Our findings are further supported by previous studies that demonstrated an increase in FSH binding sites in the rat ovary during the PRE period (43, 44). The lower expressions of *FSHR* in POST heifers in diestrus is in accord with the higher progesterone concentrations during luteal phase because progesterone exerts a negative feedback inhibition on GnRH and suppress further follicular growth and maturation (45).

The POST heifers in diestrus exhibited greater *LHR* mRNA expression than PRE heifers. These animals are in luteal phase of their estrous cycle, and it is well known that LH has essential role for maintenance and normal function of the CL (46). Therefore, LHR signaling is likely to play an essential role in suppressing ovarian activity in the remainder of the diestrus ovary.

The BMPs are important for the regulation of follicular development, ovulation, and CL morphogenesis (12, 16, 19). Interestingly, in the present study, no significant differences in mRNA abundance of *GDF9*, *BMP15*, *BMP6*, *TGFβ2*, and their receptor *BMPR2* were observed between the PRE versus POST contrasts. Although these genes are known as modulators of mammalian folliculogenesis and were therefore selected for investigation, it seems they are not involved with the changes in follicular dynamics observed in this contrast of PRE versus POST (luteal phase). The specific BMP associated with diestrus and pubertal development seems to be *BMP7*.

In the PRE versus POST (luteal phase) contrast, *BMP7* was differentially expressed. Our findings demonstrated higher

abundance of ovarian *BMP7* in PRE than in POST heifers, which may be associated with the different physiological phases represented in this contrast. The ovaries of PRE heifers are likely to be constantly initiating in follicular recruitment. The recruitment stage is characterized by the presence of a cohort of growing follicles at the beginning of the follicular wave. Our finding is consistent with previous studies in mice and human showing that *BMP7* promoted the recruitment of primordial follicles into the growing follicles and enhanced estradiol secretion when secretion of progesterone was concomitantly suppressed (9, 47, 48). Because progesterone is important in the process of ovulation (49), inhibition of progesterone production by *BMP7* suggests a functional role as luteinization inhibitor delaying ovulation (50, 51). Furthermore, in an *in vitro* study, *BMP7* increased *FSHR* mRNA expression in rat and human granulosa cells (18, 52), while it decreased *LHR* expression in human granulosa cells (18). In this context, the results of the present study suggest an important mechanism of *FSHR* and *LHR* regulation by *BMP7* in PRE heifers. Greater ovarian *BMP7* contributes to increased FSH sensitivity of granulosa cells *via* upregulation of *FSHR* and down-regulation of *LHR*, thus promoting folliculogenesis (recruitment, selection, and atresia) and suppressing ovulation (18).

We observed differential expression of key steroidogenic genes *CYP19A1* and *HSD3B1* between PRE and POST heifers. *CYP19A1* codes for the key regulator enzyme in the steroid biosynthesis pathways: P450aromatase, which catalyzes the conversion of androgens to estrogens (53). The upregulation of *BMP7* may explain the increased expression of *CYP19A1* in PRE heifers when compared to the diestrus POST heifers. An *in vivo* study demonstrated that rat ovaries treated with *BMP7*

had enhanced expression of *CYP19A1*, which in turn increased estradiol production, as well as increased the number of preantral and antral follicles (47, 54).

Greater expression of *HSD3B1* mRNA in POST heifers in their luteal phase could be the result of the presence of an active CL and consequent progesterone signaling. Theca- and granulosa-derived luteal cells express the enzyme HSD3B1 that converts pregnenolone to progesterone (55). In short, the greater expression of *HSD3B1* is necessary for production of progesterone, which is generally not occurring in PRE heifers.

Transforming growth factor-beta isoforms (TGFB1 and TGFB2) are multifunctional regulatory molecules because it can either stimulate or inhibit proliferation, differentiation, and other critical cell functions according to species, stage of differentiation of ovarian cells, and presence of others growth factors (48, 56, 57). In rodents, TGFB1 increased proliferation of FSH-stimulated granulosa cells (58). However, in cattle (TGFB2) and sheep (TGFB1 and TGFB2), these growth factors have inhibitory effects on granulosa cell proliferation (59, 60). The effect of TGFB isoforms are species specific and might be different in *B. indicus* and *B. taurus* animals. *TGFB1* mRNA and protein expression in bovine granulosa cells decrease during progress of folliculogenesis (61). Furthermore, in cattle, TGFB1 was present in granulosa cells of earliest stage of development (early preantral and early antral follicles) but absent in larger more advanced follicles (62). We observed greater expression of *TGFB1* mRNA in PRE heifers compared to POST heifers (luteal phase), which highlights the important role of TGFB1 in modulating both granulosa cell growth and differentiation. We can speculate that the proliferation inhibition induced by TGFB1 may promote differentiation of granulosa cell in order to acquire *FSHR* and express steroidogenic enzymes, which allow the cell to be more responsive to FSH and enables secretion of ovarian steroids. An improved understanding of the mechanisms of TGFB1 in follicular growth and pubertal development in bovine ovary requires further investigation. Especially, it will be relevant to understand the interactions between BMP7 and TGFB1 in regulation of bovine folliculogenesis. To achieve a better understanding of this interaction, it will be relevant to study the expression of BMP7 and TGFB1 in all phases of the estrus cycle for POST heifers.

Intrafollicular IGF1 and IGF2 have been reported to act in synergy with the gonadotropins LH and FSH to stimulate growth and differentiation of ovarian follicles and subsequent synthesis and secretion of estradiol and progesterone production (24–27). The type 1 IGF receptor (IGFR1) mediates most of the actions of both IGF1 and IGF2 (63), and its expression increases during follicular development (64). In this current study, we observed greater expression of *IGF1* and *IGFR1* in PRE compared to POST heifers (luteal phase). Our findings suggest that IGF1, as well as BMP7, could be underpinning the increased expression of *FSHR* and *CYP19A1* mRNA found in PRE heifers ovaries. This is in agreement with a previous *in vivo* study that showed IGF1 can regulate *FSHR* gene expression and indirectly aromatase expression (65). *FSHR* expression has been shown to be severely reduced in preantral follicles of *IGF1* knockout mice, as well as aromatase expression, and was restored to wild-type expression levels after

2 weeks of exogenous IGF1 supplementation (66). Reinforcing this hypothesis an *in vivo* study showed that FSH enhanced *IGFR1* expression (65). Thus, local IGF1 creates a positive feedback loop where IGF1 enhances FSH action and FSH enhances IGF1 action through mutual receptor upregulation.

Taken together, our findings suggest that BMP7 and IGF1 may interact and regulate intrafollicular steroidogenesis and follicular response to gonadotropins during pubertal development. The emerging hypothesis is that two regulators of ovarian activity prepuberty are co-expressed: *BMP7* and *IGF1*. These two genes may play synergic roles modulating the expression of *FSHR*, *LHR*, and *IGFR1*, and steroidogenesis (*CYP19A1*). *BMP7* could be associated with downregulation of *LHR* and upregulation of *FSHR* and *CYP19A1*, which mediates the follicular dynamics in PRE heifer ovaries as compared to POST (luteal phase). Upregulation of *IGF1* expression in PRE compared to POST (luteal phase) correlates with increased levels *FSHR* and *CYP19A1*. Furthermore, TGFB1 likely plays important role in follicular dynamics in prepuberty. The role of these genes and predicted interactions merits further research to elucidate the molecular pathways in cattle ovarian tissue, pre- and postpuberty, especially to elucidate these genes expression in all cycle phases.

In summary, our results suggest that differential expression of ovarian genes could be related to changes in follicular dynamics and differences of gene expression levels within the cell population that formed the ovarian tissue, which had emerged as consequence of the luteal phase and puberty. The comparative expression of these genes in granulosa versus luteal cells in cattle PRE versus POST (including all cycle phases) should be the focus of further research to better understand their role in puberty. The data presenting herein are a starting point that reinforces the differentially expressed genes as relevant genes for ovarian dynamics in *B. indicus* heifers.

AUTHOR CONTRIBUTIONS

MW, MF, LP-N, SL, and SM participated in the design of the study. MF, LP-N, LK, JR, SE, GB-H, and EP carried the field trial and euthanasia of heifers. BV contributed to handling of tissue samples and RNA extraction. MW performed qRT-PCR experiment. MW and MK performed statistical analyses. MW, MF, SG, GB-H, SL, and SM drafted and revised the manuscript. All authors read and approved the final manuscript.

ACKNOWLEDGMENTS

The authors gratefully acknowledge The University of Queensland (UQ) in permitting the study to be performed and the Animal Genetics Laboratory (AGL at UQ) for their technical support.

FUNDING

Financial support for this Ph.D. project was provided by The University of Queensland and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES/project number A125/2013). MF is supported by the UQ Postdoctoral Fellowship.

REFERENCES

- Bliss SP, Navratil AM, Xie J, Roberson MS. GnRH signaling, the gonadotrope and endocrine control of fertility. *Front Neuroendocrinol* (2010) 31(3):322–40. doi:10.1016/j.yfrne.2010.04.002
- Burrow HM. Importance of adaptation and genotype x environment interactions in tropical beef breeding systems. *Animal* (2012) 6(5):729–40. doi:10.1017/S175173111200002X
- Abeygunawardena H, Dematawewa CM. Pre-pubertal and postpartum anestrus in tropical Zebu cattle. *Anim Reprod Sci* (2004) 8(2–83):373–87. doi:10.1016/j.anireprosci.2004.05.006
- Burns BM, Fordyce G, Holroyd RG. A review of factors that impact on the capacity of beef cattle females to conceive, maintain a pregnancy and wean a calf—implications for reproductive efficiency in northern Australia. *Anim Reprod Sci* (2010) 122(1–2):1–22. doi:10.1016/j.anireprosci.2010.04.010
- Nogueira GP. Puberty in South American *Bos indicus* (Zebu) cattle. *Anim Reprod Sci* (2004) 8(2–83):361–72. doi:10.1016/j.anireprosci.2004.04.007
- Brito LF, Silva AE, Unanian MM, Dode MA, Barbosa RT, Kastelic JP. Sexual development in early- and late-maturing *Bos indicus* and *Bos indicus* x *Bos taurus* crossbred bulls in Brazil. *Theriogenology* (2004) 62(7):1198–217. doi:10.1016/j.theriogenology.2004.01.006
- Johnston DJ, Barwick SA, Corbet NJ, Fordyce G, Holroyd RG, Williams PJ, et al. Genetics of heifer puberty in two tropical beef genotypes in Northern Australia and associations with heifer- and steer-production traits. *Anim Prod Sci* (2009) 49(6):399–412. doi:10.1071/EA08276
- Van den Hurk R, Zhao J. Formation of ovarian follicles and their growth, differentiation and maturation within ovarian follicles. *Theriogenology* (2005) 63(6):1717–51. doi:10.1016/j.theriogenology.2004.08.005
- Knight PG, Glistler C. TGF-beta superfamily members and ovarian follicle development. *Reproduction* (2006) 132(2):191–206. doi:10.1530/rep.1.01074
- Otsuka F. Multiple endocrine regulation by bone morphogenetic protein system. *Endocr J* (2010) 57(1):3–14. doi:10.1507/endocrj.K09E-310
- Corduk N, Abbas G, Yildirim B, Sarioglu-Buke A. The effect of vitamin D on expression of TGF beta1 in ovary. *Exp Clin Endocrinol Diabetes* (2012) 120(8):490–3. doi:10.1055/s-0032-1314858
- Nagashima T, Kim J, Li Q, Lydon JP, DeMayo FJ, Lyons KM, et al. Connective tissue growth factor is required for normal follicle development and ovulation. *Mol Endocrinol* (2011) 25(10):1740–59. doi:10.1210/me.2011-1045
- Paradis F, Novak S, Murdoch GK, Dyck MK, Dixon WT, Foxcroft GR. Temporal regulation of BMP2, BMP6, BMP15, GDF9, BMP1A, BMP1B, BMP2 and TGFBR1 mRNA expression in the oocyte, granulosa and theca cells of developing preovulatory follicles in the pig. *Reproduction* (2009) 138(1):115–29. doi:10.1530/REP-08-0538
- Proetzel G, Pawlowski SA, Wiles MV, Yin M, Boivin GP, Howles PN, et al. Transforming growth factor-beta 3 is required for secondary palate fusion. *Nat Genet* (1995) 11:409–14. doi:10.1038/ng1295-409
- Sanford LP, Ormsby I, Gittenberger-de Groot AC, Sariola H, Friedman R, Boivin GP, et al. TGFbeta 2 knockout mice have multiple developmental defects that are nonoverlapping with other TGFbeta knockout phenotypes. *Development* (1997) 124(13):2659–70.
- Otsuka F, Moore RK, Shimasaki S. Biological function and cellular mechanism of bone morphogenetic protein-6 in the ovary. *J Biol Chem* (2001) 276(35):32889–95. doi:10.1074/jbc.M103212200
- Otsuka F, Yamamoto S, Erickson GF, Shimasaki S. Bone morphogenetic protein-15 inhibits follicle-stimulating hormone (FSH) action by suppressing FSH receptor expression. *J Biol Chem* (2001) 276(14):11387–92. doi:10.1074/jbc.M010043200
- Shi J, Yoshino O, Osuga Y, Nishii O, Yano T, Taketani Y. Bone morphogenetic protein 7 (BMP-7) increases the expression of follicle-stimulating hormone (FSH) receptor in human granulosa cells. *Fertil Steril* (2010) 93(4):1273–9. doi:10.1016/j.fertnstert.2008.11.014
- Shimasaki S, Moore RK, Otsuka F, Erickson GF. The bone morphogenetic protein system in mammalian reproduction. *Endocr Rev* (2004) 25(1):72–101. doi:10.1210/er.2003-0007
- Barros CM, Satrapa RA, Castilho AC, Fontes PK, Razza EM, Ereno RL, et al. Effect of superstimulatory treatments on the expression of genes related to ovulatory capacity, oocyte competence and embryo development in cattle. *Reprod Fertil Dev* (2012) 25:17–25. doi:10.1071/RD12271
- Adashi EY, Roban RM. Intraovarian regulation: peptidergic signalling systems. *Trends Endocrinol Metab* (1992) 3(7):243–8. doi:10.1016/1043-2760(92)90125-K
- Armstrong DG, Webb R. Ovarian follicular dominance: the role of intraovarian growth factors and novel proteins. *Rev Reprod* (1997) 2(3):139–46. doi:10.1530/ror.0.0020139
- Giudice LC. Insulin-like growth factors and ovarian development. *Endocr Rev* (1992) 13(4):641–69. doi:10.1210/edrv-13-4-641
- Bergh C, Olsson JH, Hillensjo T. Effect of insulin-like growth factor I on steroidogenesis in cultured human granulosa cells. *Acta Endocrinol* (1991) 125(2):177–85. doi:10.1530/acta.0.1250177
- Adashi EY, Resnick CE, Brodie AM, Svoboda ME, Van Wyk JJ. Somatomedin-C-mediated potentiation of follicle-stimulating hormone-induced aromatase activity of cultured rat granulosa cells. *Endocrinology* (1985) 117(6):2313–20. doi:10.1210/endo-117-6-2313
- Spicer LJ, Aad PY. Insulin-like growth factor (IGF) 2 stimulates steroidogenesis and mitosis of bovine granulosa cells through the IGF1 receptor: role of follicle-stimulating hormone and IGF2 receptor. *Biol Reprod* (2007) 77(1):18–27. doi:10.1095/biolreprod.106.058230
- Veldhuis JD, Rodgers RJ. Mechanisms subserving the steroidogenic synergism between follicle-stimulating hormone and insulin-like growth factor I (somatomedin C). Alterations in cellular sterol metabolism in swine granulosa cells. *J Biol Chem* (1987) 262(16):7658–64.
- Perks CM, Peters AR, Wathes DC. Follicular and luteal expression of insulin-like growth factors I and II and the type 1 IGF receptor in the bovine ovary. *J Reprod Fertil* (1999) 116(1):157–65. doi:10.1530/jrf.0.1160157
- Armstrong DG, Gutierrez CG, Baxter G, Glazyrin AL, Mann GE, Woad KJ, et al. Expression of mRNA encoding IGF-I, IGF-II and type 1 IGF receptor in bovine ovarian follicles. *J Endocrinol* (2000) 165(1):101–13. doi:10.1677/joe.0.1650101
- Cheong RY, Czielesky K, Porteous R, Herbison AE. Expression of ESR1 in glutamatergic and GABAergic neurons is essential for normal puberty onset, estrogen feedback, and fertility in female mice. *J Neurosci* (2015) 35:14533–43. doi:10.1523/JNEUROSCI.1776-15.2015
- Rovani MT, Gasperin BG, Ilha GF, Ferreira R, Bohrer RC, Duggavathi R, et al. Expression and molecular consequences of inhibition of estrogen receptors in granulosa cells of bovine follicles. *J Ovarian Res* (2014) 7:96. doi:10.1186/s13048-014-0096-0
- Evans AC, Adams GP, Rawlings NC. Endocrine and ovarian follicular changes leading up to the first ovulation in prepubertal heifers. *J Reprod Fertil* (1994) 100(1):187–94. doi:10.1530/jrf.0.1000187
- Nakada K, Moriyoshi M, Nakao T, Watanabe G, Taya K. Changes in concentrations of plasma immunoreactive follicle-stimulating hormone, luteinizing hormone, estradiol-17beta, testosterone, progesterone, and inhibin in heifers from birth to puberty. *Domest Anim Endocrinol* (2000) 18(1):57–69. doi:10.1016/S0739-7240(99)00063-6
- Ojeda SR, Dubay C, Lomniczi A, Kaidar G, Matagne V, Sandau US, et al. Gene networks and the neuroendocrine regulation of puberty. *Mol Cell Endocrinol* (2010) 324(1–2):3–11. doi:10.1016/j.mce.2009.12.003
- Curlew JD, Axelson M, Stone GM. Identification of the major steroids in ovarian and adrenal venous plasma of the brush-tail possum (*Trichosurus vulpecula*) and changes in the peripheral plasma-levels of estradiol and progesterone during the reproductive-cycle. *J Endocrinol* (1985) 105:53–62. doi:10.1677/joe.0.1050053
- Weller MM, Fortes MRS, Marcondes MI, Rotta PP, Gionbeli TR, Valadares Filho SC, et al. Effect of maternal nutrition and days of gestation on pituitary gland and gonadal gene expression in cattle. *J Dairy Sci* (2016) 99:3056–71. doi:10.3168/jds.2015-9673
- Steibel JP, Poletto R, Coussens PM, Rosa GJM. A powerful and flexible linear mixed model framework for the analysis of relative quantification RT-PCR data. *Genomics* (2009) 94(2):146–52. doi:10.1016/j.ygeno.2009.04.008
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2 DDCT method. *Methods* (2001) 25(4):402–8. doi:10.1006/meth.2001.1262
- Fortes MR, Li Y, Collis E, Zhang Y, Hawker RJ. The IGF1 pathway genes and their association with age of puberty in cattle. *Anim Genet* (2013) 44(1):91–5. doi:10.1111/j.1365-2052.2012.02367.x

40. Roche JF, Boland MP. Turnover of dominant follicles in cattle of different reproductive states. *Theriogenology* (1991) 35(1):81–90. doi:10.1016/0093-691X(91)90149-8
41. Richards JS. Hormonal control of gene expression in the ovary. *Endocr Rev* (1994) 15(6):725–51. doi:10.1210/edrv-15-6-725
42. Bao B, Garverick HA, Smith GW, Smith MF, Salfen BE, Youngquist RS. Changes in messenger RNA encoding LH receptor, cytochrome P450 side chain cleavage, and aromatase are associated with recruitment and selection of bovine ovarian follicles. *Biol Reprod* (1997) 56(5):1158–68. doi:10.1095/biolreprod56.5.1158
43. Tanaka Y, Nakada K, Moriyoshi M, Sawamukai Y. Appearance and number of follicles and change in the concentration of serum FSH in female bovine fetuses. *Reproduction* (2001) 121(5):777–82. doi:10.1530/rep.0.1210777
44. Peluso B, Steger RW, Hafez ESE. Development of gonadotrophin-binding sites in the immature rat ovary. *J Reprod Fertil* (1976) 47:55–8. doi:10.1530/jrf.0.0470055
45. Sokka T, Huhtaniemi I. Ontogeny of gonadotrophin receptors and gonadotrophin stimulated cyclic AMP production in the neonatal rat ovary. *J Endocrinol* (1990) 127:297–303. doi:10.1677/joe.0.1270297
46. Webb R, Gong JG, Law AS, Rusbridge SM. Control of ovarian function in cattle. *J Reprod Fertil Suppl* (1992) 5:141–56.
47. Niswender GD, Juengel JL, Silva PJ, Rollyson MK, McIntush EW. Mechanisms controlling the function and life span of the corpus luteum. *Physiol Rev* (2000) 80(1):1–29.
48. Lee WS, Otsuka F, Moore RK, Shimasaki S. Effect of bone morphogenetic protein-7 on folliculogenesis and ovulation in the rat. *Biol Reprod* (2001) 65(4):994–9. doi:10.1095/biolreprod65.4.994
49. Zhang H, Klausen C, Zhu H, Chang HM, Leung PC. BMP4 and BMP7 suppress StAR and progesterone production via ALK3 and SMAD1/5/8-SMAD4 in human granulosa-lutein cells. *Endocrinology* (2015) 156(11):4269–80. doi:10.1210/en.2015-1494
50. Yoshimura Y, Wallach EE. Studies of the mechanism(s) of mammalian ovulation. *Fertil Steril* (1987) 47(1):22–34. doi:10.1016/S0015-0282(16)49931-0
51. Shimasaki S, Zachow RJ, Li D, Kim H, Iemura S, Ueno N, et al. A functional bone morphogenetic protein system in the ovary. *Proc Natl Acad Sci U S A* (1999) 96(13):7282–7. doi:10.1073/pnas.96.13.7282
52. Findlay JK, Drummond AE, Dyson ML, Baillie AJ, Robertson DM, Ethier JF. Recruitment and development of the follicle; the roles of the transforming growth factor- β superfamily. *Mol Cell Endocrinol* (2002) 191(1):35–43. doi:10.1016/S0303-7207(02)00053-9
53. Lee WS, Yoon SJ, Yoon TK, Cha KY, Lee SH, Shimasaki S, et al. Effects of bone morphogenetic protein-7 (BMP-7) on primordial follicular growth in the mouse ovary. *Mol Reprod Dev* (2004) 69(2):159–63. doi:10.1002/mrd.20163
54. Meinhardt U, Mullis PE. The essential role of the aromatase/p450arom. *Semin Reprod Med* (2002) 20(3):277–84. doi:10.1055/s-2002-35374
55. Gougeon A. Human ovarian follicular development: from activation of resting follicles to preovulatory maturation. *Ann Endocrinol* (2010) 71(3):132–43. doi:10.1016/j.ando.2010.02.021
56. Hanukoglu I. Steroidogenic enzymes: structure, function, and role in regulation of steroid hormone biosynthesis. *J Steroid Biochem* (1992) 43(8):779–804. doi:10.1016/0960-0760(92)90307-5
57. Lerner AA, Salamone DF, Chiappe ME, Baranao JL. Comparative studies between freshly isolated and spontaneously immortalized bovine granulosa cells: protein secretion, steroid metabolism, and responsiveness to growth factors. *J Cell Physiol* (1995) 164(2):395–403. doi:10.1002/jcp.1041640220
58. Juengel JL, McNatty KP. The role of proteins of the transforming growth factor- β superfamily in the intraovarian regulation of follicular development. *Hum Reprod Update* (2005) 11(2):143–60. doi:10.1093/humupd/dmh061
59. Saragueta PE, Lanuza GM, Baranao JL. Autocrine role of transforming growth factor β 1 on rat granulosa cell proliferation. *Biol Reprod* (2002) 66(6):1862–8. doi:10.1095/biolreprod66.6.1862
60. Gilchrist RB, Morrissey MP, Ritter LJ, Armstrong DT. Comparison of oocyte factors and transforming growth factor- β in the regulation of DNA synthesis in bovine granulosa cells. *Mol Cell Endocrinol* (2003) 201(1–2):87–95. doi:10.1016/S0303-7207(02)00429-X
61. Juengel JL, Bibby AH, Reader KL, Lun S, Quirke LD, Haydon LJ, et al. The role of transforming growth factor- β (TGF- β) during ovarian follicular development in sheep. *Reprod Biol Endocrinol* (2004) 2:78–88. doi:10.1186/1477-7827-2-78
62. Nilsson EE, Doraiswamy V, Skinner MK. Transforming growth factor- β isoform expression during bovine ovarian antral follicle development. *Mol Reprod Dev* (2003) 66(3):237–46. doi:10.1002/mrd.10350
63. Oullette Y, Price CA, Carrière PD. Follicular fluid concentration of transforming growth factor- β 1 is negatively correlated with estradiol and follicle size at the early stage of development of the first-wave cohort of bovine ovarian follicles. *Domest Anim Endocrinol* (2005) 29(4):623–33. doi:10.1016/j.domaniend.2005.04.008
64. Lucy MC. Regulation of ovarian follicular growth by somatotropin and insulin-like growth factors in cattle. *J Dairy Sci* (2000) 83(7):1635–47. doi:10.3168/jds.S0022-0302(00)75032-6
65. Wandji SA, Pelletier G, Sirard MA. Ontogeny and cellular localization of 125I-labeled insulin-like growth factor-I, 125I-labeled follicle-stimulating hormone, and 125I-labeled human chorionic gonadotropin binding sites in ovaries from bovine fetuses and neonatal calves. *Biol Reprod* (1992) 47(5):814–22. doi:10.1095/biolreprod47.5.814
66. Zhou J, Chin E, Bondy CA. Cellular pattern of IGF-I and IGF-I receptor gene expression in the developing and mature follicle. *Endocrinology* (1991) 129(6):3281–8. doi:10.1210/endo-129-6-3281

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Weller, Fortes, Porto-Neto, Kelly, Venus, Kidd, do Rego, Edwards, Boe-Hansen, Piper, Lehnert, Guimarães and Moore. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.