

Multilocus sequence typing methods for the emerging *Campylobacter* species *C. hyointestinalis*, *C. lanienae*, *C. sputorum*, *C. concisus*, and *C. curvus*

William G. Miller^{1*}, Mary H. Chapman¹, Emma Yee¹, Stephen L. W. On², Desmond K. McNulty^{3†}, Albert J. Lastovica⁴, Anne M. Carroll³, Eleanor B. McNamara³, Geraldine Duffy⁵ and Robert E. Mandrell¹

¹ USDA, ARS, WRRRC, Produce Safety and Microbiology, Albany, CA, USA

² Christchurch Science Centre, Institute of Environmental Science and Research Limited, Christchurch, New Zealand

³ Public Health Laboratory, Health Service Executive, Cherry Orchard Hospital, Ballyfermot, Dublin, Ireland

⁴ Department of Biotechnology, University of the Western Cape, Bellville, South Africa

⁵ Food Safety Department, Teagasc Food Research Centre, Ashtown, Dublin, Ireland

Edited by:

Alain Stintzi, Ottawa Institute of Systems Biology, Canada

Reviewed by:

Qijing Zhang, Iowa State University, USA

Eduardo Taboada, Public Health Agency of Canada, Canada

*Correspondence:

William G. Miller, USDA, ARS, WRRRC, Produce Safety and Microbiology Research Unit, 800 Buchanan Street, Albany, CA 94710, USA.
e-mail: william.miller@ars.usda.gov

†Present address:

Desmond K. McNulty, ICON Central Laboratories, South County Business Park, Leopardstown, Dublin, Ireland.

Multilocus sequence typing (MLST) systems have been reported previously for multiple food- and food animal-associated *Campylobacter* species (e.g., *C. jejuni*, *C. coli*, *C. lari*, and *C. fetus*) to both differentiate strains and identify clonal lineages. These MLST methods focused primarily on campylobacters of human clinical (e.g., *C. jejuni*) or veterinary (e.g., *C. fetus*) relevance. However, other, emerging, *Campylobacter* species have been isolated increasingly from environmental, food animal, or human clinical samples. We describe herein four MLST methods for five emerging *Campylobacter* species: *C. hyointestinalis*, *C. lanienae*, *C. sputorum*, *C. concisus*, and *C. curvus*. The *concisus/curvus* method uses the loci *aspA*, *atpA*, *glnA*, *gltA*, *glyA*, *ilvD*, and *pgm*, whereas the other methods use the seven loci defined for *C. jejuni* (i.e., *aspA*, *atpA*, *glnA*, *gltA*, *glyA*, *pgm*, and *tkf*). Multiple food animal and human clinical *C. hyointestinalis* ($n=48$), *C. lanienae* ($n=34$), and *C. sputorum* ($n=24$) isolates were typed, along with 86 human clinical *C. concisus* and *C. curvus* isolates. A large number of sequence types were identified using all four MLST methods. Additionally, these methods speciated unequivocally isolates that had been typed ambiguously using other molecular-based speciation methods, such as 16S rDNA sequencing. Finally, the design of degenerate primer pairs for some methods permitted the typing of related species; for example, the *C. hyointestinalis* primer pairs could be used to type *C. fetus* strains. Therefore, these novel *Campylobacter* MLST methods will prove useful in differentiating strains of multiple, emerging *Campylobacter* species.

Keywords: MLST, emerging, *Campylobacter hyointestinalis*, *Campylobacter lanienae*, *Campylobacter concisus*, *Campylobacter curvus*, *Campylobacter sputorum*

INTRODUCTION

Campylobacters are a major cause of human bacterial gastrointestinal illness in the industrialized world (Mølbak and Havelaar, 2008; Olson et al., 2008); campylobacterioses (12.68 cases per 100,000) were second only to *Salmonella* infections (16.2/100,000) in the United States in 2008 (Anonymous, 2009). The majority of *Campylobacter* strains isolated from human clinical samples have been identified as *C. jejuni* subsp. *jejuni* or, to a lesser extent, *C. coli* (Lastovica and Allos, 2008). Recently, pathogenic campylobacters outside of the *C. jejuni/C. coli* group, termed here as emerging *Campylobacter* species, have been isolated more frequently from food and/or food animals. Recovery of these more fastidious, emerging *Campylobacter* species from food has not been reported often; isolation of such strains is likely limited by the culture conditions employed, conditions that favor *Campylobacter* species such as *C. jejuni* and *C. coli*. However, Lynch et al. (2011) using novel culture conditions, reported the isolation of multiple emerging *Campylobacter* spp., e.g., *C. concisus*, *C. curvus*, and *C. sputorum*,

from chicken, beef, and pork samples. Emerging campylobacters isolated from food animals are often strains of species associated typically with livestock, such as *C. hyointestinalis* in sheep, cattle, and swine (Hakkinen et al., 2007; Salihi et al., 2009; Oporto and Hurtado, 2011), *C. lanienae* in cattle and swine (Sasaki et al., 2003; Inglis et al., 2004; Oporto and Hurtado, 2011), and *C. sputorum* in cattle and sheep (Terzolo, 1988; On et al., 1998).

The clinical relevance of the emerging *Campylobacter* spp. is as yet undetermined. Many of the emerging campylobacters are isolated infrequently from human clinical samples, although, as with isolation from food, recovery of these strains from clinical samples is probably limited by the isolation methods and media used. Nevertheless, emerging *Campylobacter* species are isolated from human clinical samples (Edmonds et al., 1987; Gorkiewicz et al., 2002; Lastovica and Allos, 2008; Bullman et al., 2011). Although the frequency of human illness associated with emerging *Campylobacter* spp. might be quite low, especially when compared to *C. jejuni*-associated gastroenteritis, it is possible that some

emerging species could be associated with more severe illness. One such example is *C. concisus*, for which a strong association with Crohn's disease and ulcerative colitis, has been reported recently (Man et al., 2010; Mahendran et al., 2011; Mukhopadhyaya et al., 2011).

Although molecular detection methods exist for many of the emerging campylobacters, population analyses, epidemiology, and source tracking of these organisms are limited by the strain typing methods available for these taxa. Molecular typing methods such as amplified fragment length polymorphism (AFLP) analysis and pulsed field gel electrophoresis (PFGE) methods have been employed on emerging *Campylobacter* strains (reviewed in On et al., 2008); however, sequence-based typing methods are not available for many species. One such sequence-based typing method is multilocus sequence typing (MLST). MLST methods amplify and sequence defined regions of moderately conserved housekeeping loci. At each locus, regions with distinct sequences receive arbitrary but unique allele numbers; similarly, each different allelic profile is assigned a unique sequence type (ST). The first *Campylobacter* MLST method was developed for *C. jejuni* (Dingle et al., 2001). This method sequences portions of seven genes: *aspA*, *atpA* (*uncA*), *glnA*, *gltA*, *glyA*, *pgm* (*glmM*), and *tkl*. The *C. jejuni* MLST method has been used in multiple typing studies and has been used successfully for strain typing and characterization, identification of clonal complexes and lineages, epidemiology, and investigation of host/source-associations (reviewed in Maiden and Dingle, 2008). Since the description of the *C. jejuni* MLST method, other *Campylobacter* MLST methods have been constructed that type *C. coli* (Dingle et al., 2005; Miller et al., 2005), *C. lari* (Miller et al., 2005), *C. upsaliensis* (Miller et al., 2005), *C. helveticus* (Miller et al., 2005), *C. fetus* (van Bergen et al., 2005), and *C. insulaenigra* (Stoddard et al., 2007). Besides the primary use of *Campylobacter* MLST data for strain typing, MLST data for multiple taxa within *Campylobacter* are a valuable resource for studies on lateral gene transfer and evolution. MLST data can be used also to identify putative and perhaps clinically relevant taxonomic subdivisions within a species (Miller et al., 2005); additionally, MLST can provide genotypic information for novel species that are diverse phenotypically (Stoddard et al., 2007), especially those for which the only molecular speciation method is 16S rDNA sequencing.

Development of several *Campylobacter* MLST methods was assisted by the availability of draft genome sequences (Miller et al., 2005). Development of the novel *Campylobacter* MLST methods described in this study utilized recent draft genomes of *C. hyointestinalis*, *C. lanienae*, and *C. sputorum* (Miller et al., unpublished data), in addition to the closed *C. concisus* and *C. curvius* genomes available in the NCBI Microbial Genomes database. We anticipated that the draft genomes would contain some sequencing errors; however, enough reliable sequencing data was available to design MLST primers that could be used to type these five *Campylobacter* species. Therefore, in this study we describe four novel MLST methods that can be used to type: (1) *C. concisus* and *C. curvius*; (2) both subspecies of *C. hyointestinalis* (subsp. *hyointestinalis* and *lawsonii*) and *C. fetus* (subsp. *fetus* and *venerealis*); (3) *C. lanienae*; and (4) all three biovars of *C. sputorum* (bvs. *fecalis*, *paraureolyticus*, and *sputorum*). All four MLST gene sets

are identical to the *C. jejuni* gene set [i.e., *aspA*, *atpA* (*uncA*), *glnA*, *gltA*, *glyA*, *pgm*, and *tkl*], with the exception of the *C. concisus*/*C. curvius* MLST method in which *ilvD* replaces *tkl*. A sample set of 213 isolates of diverse geographic origin and source was typed in this study. For all four methods, a total of 163 STs and 729 alleles were identified, indicating that these new MLST methods provide resolution similar to the previous MLST methods described.

MATERIALS AND METHODS

GROWTH CONDITIONS AND CHEMICALS

All *Campylobacter* strains were cultured routinely under microaerobic conditions (1.5% O₂, 10% H₂, 10% CO₂, and 78.5% N₂) at 37°C on Brain Heart Infusion agar (Becton Dickinson, Sparks, MD, USA) or Anaerobe Basal Agar (ABA; Oxoid, Lenexa, KS, USA) supplemented with 5% (v/v) laked horse blood (Hema Resource and Supply, Aurora, OR, USA). PCR enzymes and reagents were purchased from New England Biolabs (Beverly, MA, USA) or Epicentre (Madison, WI, USA). All chemicals were purchased from Sigma-Aldrich Chemicals (St. Louis, MO, USA) or Fisher Scientific (Pittsburgh, PA, USA). DNA sequencing chemicals and capillaries were purchased from Applied Biosystems (Foster City, CA, USA). PCR and sequencing oligonucleotides were purchased from Eurofins MWG Operon (Huntsville, AL, USA).

ISOLATION OF *CAMPYLOBACTER* FROM FERAL SWINE AND CATTLE

Cattle feces were inoculated into wells of a six-well microtiter plate containing 6 ml 1 × Anaerobe Basal Broth (Oxoid) amended with Preston supplement (Oxoid), using a sterile cotton swab. Plates were placed inside plastic ZipLoc bags and incubated under microaerobic conditions (as above) for 24 h at 37°C, while shaking at 40 rpm. After incubation, a 10-μl loop of each enrichment culture was plated onto ABA amended with 5% laked horse blood and CAT supplement (Oxoid). Feral swine feces were plated directly, using a sterile cotton swab, onto ABA amended with 5% laked horse blood and CAT supplement. All plates were then incubated under microaerobic conditions at 37°C for 24 h. Bacterial cultures were then filtered through 0.2 μm mixed cellulose ester filters onto ABA plates and incubated at 37°C under microaerobic conditions. After 24 h, single colonies were streaked onto new ABA plates and incubated 24–48 h at 37°C for purification.

CAMPYLOBACTER SPECIATION

Campylobacter strains isolated from the feces of California feral swine or cattle were speciated initially by 16S rDNA sequencing, using the primer pairs 27F (5' AGA GTT TGA TCC TGG CTC AG 3') and 1392R (5' GAC GGG CGG TGT GTA C 3'; Lane, 1991). However, the 27F/1392R primers were not able to type *C. hyointestinalis* strains past the species level and several strains could not be typed unequivocally. To improve speciation, the *atpA*/*atpA*R primer pairs from Miller et al. (2005) were used. These primers can amplify all *Campylobacter* taxa described currently, with the exception of *C. avium* (data not shown). Using these *atpA* primers, campylobacters of uncertain type were amplified and sequenced; the sequences were then compared by alignments and phylogenetic analyses to strains of known species/subspecies identification, especially those whose genomes had been sequenced. Unlike the 16S primers, the *atpA* primers could speciate unambiguously all

of the strains isolated in this study and could identify clearly both subspecies of *C. hyointestinalis*. The *atpA* primer pairs, however, could not differentiate the three known biovars of *C. sputorum* (i.e., *sputorum*, *fecalis*, and *paraureolyticus*; On et al., 1998). The MLST results provided further confirmatory speciation data; *atpA* speciation agreed completely with subsequent MLST speciation.

DETECTION OF UREASE AND CATALASE ACTIVITY

The biovar *paraureolyticus* can be distinguished from the other two biovars of *C. sputorum* by the production of urease (On et al., 1998). Therefore, to identify putative bv. *paraureolyticus* strains, *C. sputorum* isolates were assayed for urease activity, as follows: a 10- μ l loop of an overnight *C. sputorum* culture was resuspended in 2 ml urease reagent (3 mM NaH₂PO₄, 110 mM urea, 7 μ g/ml phenol red, pH 6.8) and incubated for 1 h at RT. *C. sputorum* cultures were typed as bv. *paraureolyticus* based on a positive reaction (solution turning from yellow/orange to magenta). Genome-sequenced strains of biovars *sputorum* (strain RM3237) and *paraureolyticus* [strain RM4120 (LMG 11764)] were used as negative and positive controls, respectively. All tests were performed independently at least twice.

The *C. sputorum* biovars are distinguished also by the production of catalase: bv. *fecalis* is catalase-positive while the other two biovars are catalase negative (On et al., 1998). To test for catalase activity, a 10- μ l loop of an overnight *C. sputorum* culture was resuspended in 200 μ l 3% H₂O₂ on a glass slide. Presence of bubbles indicated a positive reaction. Genome-sequenced strains of biovars *sputorum* (strain RM3237) and *fecalis* [strain RM4121 (CCUG 20703)] were used as negative and positive controls, respectively. All tests were performed independently at least twice.

MULTILOCUS SEQUENCE TYPING

Each MLST amplification mixture contained: 1 \times MasterAmp PCR buffer (Epicentre, Madison, WI, USA), 1 \times MasterAmp PCR enhancer (Epicentre), 2.5 mM MgCl₂, 250 μ M (each) dNTPs, 50 pmol each primer, and 1 U *Taq* polymerase (New England Biolabs). For strains where genomic DNA was extracted using kits or standard isolation protocols, 50 ng purified genomic DNA was added to each reaction tube. Otherwise, 2 μ l of a boilate was added. Boilates were prepared by resuspending a 1- μ l loop of a pure culture or a single Microbank bacterial storage bead (Pro-Lab, Austin, TX, USA) in 100 μ l TE and heating at 80°C for 5 min, then 100°C for 20 min, and cooling to 4°C. MLST amplifications were performed on a Tetrad thermocycler (Bio-Rad, Hercules, CA, USA) with the following settings: 94°C for 30 s, 53°C for 30 s, and 72°C for 2 min (30 cycles). Amplicons were purified on a BioRobot 8000 workstation (Qiagen, Valencia, CA, USA). Cycle sequencing reactions were performed on a Tetrad thermocycler, using the ABI PRISM BigDye terminator cycle sequencing kit (version 3.1; Applied Biosystems) and standard protocols. Cycle sequencing extension products were purified using BigDye XTerminator (Applied Biosystems). DNA sequencing was performed on an ABI PRISM 3730 DNA Analyzer (Applied Biosystems), using POP-7 polymer and ABI PRISM Genetic Analyzer Data Collection and ABI PRISM Genetic Analyzer Sequencing Analysis software. Sequences were trimmed, assembled, and analyzed in SeqMan (v 9.1; DNASTAR, Madison, WI, USA).

ALLELE NUMBER/SEQUENCE TYPE ASSIGNMENT

The Perl program MLSTparser3 (Miller et al., 2009) was modified to include the novel MLST methods for *C. concisus*, *C. curvus*, *C. hyointestinalis*, *C. lanienae*, and *C. sputorum*. The expanded MLSTparser3 was used to identify the MLST alleles and ST of each *Campylobacter* strain typed in this study. New *Campylobacter* MLST databases were created¹; allele and ST data generated in this study were deposited in this database and are available online. The allelic profiles for all 213 strains are listed in Table S1 in Supplementary Material.

PHYLOGENETIC ANALYSES

A dendrogram of unique *Campylobacter* STs was constructed by concatenating the allele sequences comprising each ST. Allele sequences for each strain were concatenated in the order *aspA-atpA-glnA-gltA-glyA-pgm-tkt* with the exception of *C. concisus* and *C. curvus* allele sequences, that were concatenated in the order *aspA-atpA-glnA-gltA-glyA-ivlD-pgm*. Composite concatenate lengths were 3345 bp (*C. concisus/C. curvus*), 3312 bp (*C. fetus*, *C. hyointestinalis*, and *C. lanienae*), or 3321 bp (*C. sputorum*). Sequence alignments were performed using CLUSTALX (ver. 2.1)², and dendrograms were constructed using the neighbor-joining method with the Kimura two-parameter distance estimation method (Kimura, 1980). Phylogenetic analyses were performed using MEGA version 5.1 (Tamura et al., 2011). Polymorphic sites and d_n/d_s ratios were calculated using START2 (Jolley et al., 2001).

RESULTS AND DISCUSSION

DESIGN OF THE NOVEL *CAMPYLOBACTER* MLST METHODS

Construction of the novel *Campylobacter* MLST methods was facilitated by the availability of genome sequences for all of the taxa typed in this study. The genome-sequenced strains were: the completed genomes of *C. concisus* strain 13826 (NC_009802.1), *C. curvus* strain 525.92 (NC_009715.1), and *C. fetus* subsp. *fetus* strain 82–40 (NC_008599.1), and the draft genomes of *C. hyointestinalis* subsp. *hyointestinalis* (*Chh*) strain RM4092 (LMG 9260), *C. hyointestinalis* subsp. *lawsonii* (*Chl*) strain RM4096 (CCUG 27631), *C. lanienae* strain RM3663 (NCTC 13004), *C. sputorum* bv. *sputorum* strain RM3237, *C. sputorum* bv. *fecalis* strain RM4121 (CCUG 20703), *C. sputorum* bv. *paraureolyticus* strain RM4120 (LMG 11764), and strain RM6914, exemplar of a novel *C. concisus*-like clade (Mandrell et al., manuscript in preparation).

Primer design based on a sequence from a single strain might not lead to a successful MLST method if the sequence variation within that taxon prevents the design of primer pairs that efficiently amplify all strains. Therefore, MLST gene sequences from related species would be aligned. Based on this alignment, primers would be designed to bind to regions, 100–200 bp upstream and downstream of the allelic endpoints, that demonstrate a high degree of conservation among the aligned taxa. One to four degenerate bases would be incorporated into the MLST primers, if necessary, to optimize primer binding. This approach was used previously to construct successfully other *Campylobacter* MLST

¹<http://pubmlst.org/campylobacter/>

²<http://www.clustal.org/>

methods (Miller et al., 2005). Therefore, the full *aspA*, *atpA*, *glnA*, *gltA*, *glyA*, *pgm*, and *tkt* gene sequences were extracted from the completed and draft genomes and various alignments were performed. Based on sequence similarity between the various *Campylobacter* taxa, we developed four novel MLST methods to type the strains in this study: Method 1 for typing both *C. concisus* and *C. curvus*; Method 2 for typing *C. fetus* and both subspecies of *C. hyointestinalis*; Method 3 for typing *C. lanienae* strains; and Method 4 for typing all three biovars of *C. sputorum*.

The final MLST primer sets are listed in **Table 1**. Methods 2, 3, and 4 use the same seven loci and allelic endpoints of the *C. jejuni* MLST method, i.e., *aspA*, *atpA*, *glnA*, *gltA*, *glyA*, *pgm*, and *tkt*. However, for the *C. concisus/C. curvus* MLST method, the sequence diversity at the *tkt* locus was too great for the construction of suitable primers. Therefore, *tkt* was replaced by *ilvD* in Method 1; *ilvD* was used in a *C. jejuni* MLST method described previously (Manning et al., 2003). The Method 1 *aspA*, *atpA*, *glnA*, *gltA*, *glyA*, and *pgm* alleles also use the same endpoints of their *C. jejuni* counterparts; the *ilvD* allelic endpoints are unique to this study.

CHARACTERIZATION OF THE FOUR NOVEL *CAMPYLOBACTER* MLST METHODS

A total of 213 strains were typed in this study. Complete descriptions of each strain, including isolation source, date, and location (if known), allelic profiles and STs are listed in Table S1 in Supplementary Material. Strains typed in this study were isolated over a 30-year period (1981–2010) and were also geographically diverse: strains from each species were isolated on two to three continents (Table S1 in Supplementary Material). All of the *C. concisus* and *C. curvus* strains typed were isolated from human clinical samples, whereas nearly all of the *Chl* and *C. lanienae* strains typed were isolated from pigs and feral swine; strains from the other three taxa were a mixture of isolates from humans, cattle, and swine (**Table 2**).

With a few exceptions, the primary MLST primers listed in **Table 1** were able to amplify successfully all seven loci and provide high quality sequence data for all 213 strains to generate unambiguous ST. However, despite our best efforts to overcome sequence variation in the initial primer design, in a few instances (~1–2%), the main primer pairs did not provide sequence quality high enough for an unambiguous ST. Thus, alternate primer pairs (annotated as “A” in **Table 1**) were used to amplify and sequence these alleles. No strain was excluded from the final strain list because a ST could not be obtained.

GENETIC DIVERSITY

Phylogenetic analysis of the MLST STs validated the taxonomic relationships observed previously (Debruyne et al., 2008) for the six *Campylobacter* species typed in this study. Although the use of different gene sets comprising Method 1 and Methods 2, 3, and 4 prevented the simultaneous analysis of all six species, a clear delineation between *C. concisus* and *C. curvus* strains was observed (**Figure 1**), as well as unambiguous segregation of the other four species (**Figure 2**), that included the related *C. hyointestinalis*, *C. fetus*, and *C. lanienae* taxa. Moreover, the two *C. hyointestinalis* subspecies, *Chh* and *Chl*, formed distinct clusters (**Figure 2**).

However, the two *C. fetus* subspecies could not be discriminated by phylogenetic analysis, consistent with previous observations (van Bergen et al., 2005). Divergent STs (*C. lanienae* STs 1 and 4; *C. sputorum* STs 8, 13, 14, and 15) were identified within some strains (see below and **Figure 2**). In *C. sputorum*, these divergent STs formed a cluster (termed Csp₂) distinct from a cluster (termed Csp₁) containing the other *C. sputorum* strains.

Many MLST STs and alleles were identified in this study (**Table 3**). With the exception of *C. fetus*, for which a previous study identified also a relatively small number of highly clonal STs (van Bergen et al., 2005), the majority of strains within each of the remaining five species possess unique STs. Phylogenetic analysis of the STs of these five species indicated that the least amount of variation exists within *Chh* and each of the two *C. sputorum* clades, whereas the greatest amount of variation detected here resides clearly within the *C. concisus* and *C. curvus* strain sets (**Figures 1–3**). Even with the inclusion of the divergent *C. lanienae* and *C. sputorum* STs, the average number of base substitutions per site was less in each case (0.0330, *C. lanienae*; 0.0265, *C. sputorum*; **Figure 3**) than that calculated for the *C. concisus* STs (0.0641; **Figure 3**).

The high degree of variation across the *C. concisus* STs is reflected by the large number of alleles and polymorphic sites identified within this strain set: for the 70 *C. concisus* strains, the number of alleles detected at any locus ranged from 55 (*atpA*) to 64 (*ilvD*; **Table 3**). This high density of alleles translated into the large number of polymorphic sites identified at each of the seven *C. concisus* loci (**Table 4A**). Over 100 polymorphic sites were present within the alleles of each of the *C. concisus* MLST loci (**Table 4A**), a large number when compared to the relatively few polymorphic sites detected within the *Chh* alleles, even when the relative sizes of the strain sets were factored into the comparison. For some species, the numbers of polymorphic sites were inflated by the presence of divergent alleles or strains within the strain set. When these alleles and strains were removed from the appropriate strain sets, the number of polymorphic sites decreased substantially. For example, removal of the divergent *C. sputorum* strains present within *C. sputorum* clade Csp₂ (**Figure 3**) eliminated 23 of 24 polymorphic sites at the *glnA* locus (**Table 4A**).

While MLST is often used as a strain typing tool, it can be used also to investigate the population structure of an organism, to identify lineages, for example, that demonstrate a higher association with disease or a particular host. Genes used for MLST methods, typically core housekeeping genes, are usually under purifying or neutral selection. Positively selected genes would be influenced by external/environmental pressures and would evolve more through recombination rather than through the accumulation of point mutations; thus, such genes are not generally used in MLST methods, since they may not provide an accurate representation of the clonal structure of a bacterial population (Maiden, 2006; Perez-Losada et al., 2011). One method of determining the level of selective pressure on a gene is by calculating the ratio between non-synonymous (d_n) and synonymous (d_s) base substitutions. The rate of synonymous base substitution in genes should equal the neutral substitution rate, in the absence of codon usage bias. Non-synonymous base substitutions (that result in an amino acid change) would be the result of positive

Table 1 | *Campylobacter* MLST primers.

Locus	Allele size (bp)	Oligonucleotide primers		Method					
		Forward (5'–3')	Reverse (5'–3')	1	2	3	4		
<i>asp</i>	477	aspCCCFN1	GGHAAAAGCACAATGAYRCTTATCC	aspCCCRN1	GCCWAGDACTGATTTTARGCAAGC	P	P		
		aspCCF1	CAAAAGCACAAATGACACTTATCCAA	aspCCR1	GRACTGATTTTAGGCAAGCCCTCAGG	A	A		
		HFaspF	CITTTGAWAAAAAGCAGARGAGTTTAA	HFaspR	GCTGTAACGATACCGATTGAGTTATAA		P	P	
		HYOaspF	AAGAGTKGCTATTATGAAAGACTTTGTG	HYOaspR	AATGCTGTAACGATACCGATTGAGT			A	
		LANaspF	TTTAGCCACAGCTATGGAGTATCTCAA	LANaspR	ATATGGTTRAAWVGCTGTAACRATACC				P
		HFLaspxF	AAYATGAAYGCAAACGAAAGTTATAGC	LANaspR	ATATGGTTRAAWVGCTGTAACRATACC				A
		SPUTaspF	GAACATTTGCAAGACGAACTAGAT	SPUTaspR	AAATGCTGTACTATACCACACTGATCC				P
		atpCCCFN1	GATACYATCATYAAACAAAAGGTCAGA	atpCCCRN2	GGTATTHGCCTCKATATAWGGATAWAGC	P	P		
		atpCCF1	ACTATCATCAAYCAAAAAGGTC AAG	atpCCR2	GGATAWAGCTCMGCTTCAAATTTTGT	A	A		
		HFatpF	GTATYAAAAGCTATWGACGGTTTGGTTC	HFatpR	GAGYGGGCTATAAGGAGGTTG		P	P	
		HYOatpF	ATGTVGCTATMGGTCAAAARCAATC	HYOatpR	TTTCTACWGGRAGYGGGCTATAAAGG			A	
		LANatpF	AACCAAAAAGGTC AAGATGTTATATGT	LANatpR	ATTTTCTACTGGAAGTGGGCTATAAAGG				P
		HFLatpXF	CMAAAGGHATYATGGCTAGAAAAT	HFLatpXR	TTCRATATCTTTATCHAGWVGCTTTTTT				A
SPUTatpF	ACTATCATAAATCAAAAAGGCGGAG	SPUTatpR	TTCTCAATACCAAGRGTGAATAAG				P		
glnA	477	glnCCCFN1	GSTTGGCAYAGCATAAGYTACAAC	glnCCCRN2	GTTYGTGCTWGGGTTTTGTRAAAGC	P	P		
		glnCCF1	GSYTGCCACAGCATAAGCTACAAC	glnCCR2	ACRRTCTTTCCARACTGATTGATGC	A	A		
		HFglnF	GGC ATCAGTATCKTATAATAATAAAGC	HFglnR	ATGTACGTGCATACCGCTTCC		P	P	
		HYOglnF	TC TTATAATAATAAAGCCGTGAGCGAA	HYOglnR	CCRCTTTCCATATACTTTGATGTAGC			A	
		LANglnF	TGGCAYCAYGTATCWTATAATAATAAAGC	LANglnR	ATGGACRTGCATACCRCTWCCATTATC				P
		HFglnXF	TTYGAATWTTGTRAWGAAAATGAAGT	HFglnXR	AGAGTAWGTWAGAATGCTTGKGGCTTC				A
		SPUTglnF	AGGAAC TTGGC ATCATGTTCTTAT	SPUTglnR	CCATCTTTCCAAAATAGATTGATGAA				P
<i>gltA</i>	402	gltCCCFN1	GGGMATACCTCRACKGCGATGTG	gltCCCRN2	CBCCRTGWGCCCAGCCCC	P	P		
		gltCCF1	TACATCAGCGGKCTAAARGGCGAG	gltCCR2	GATCTCTCWAGCTGGCGGATGAC	A	A		
		HFgltF	CTATAACRTTATMGATGGWGA AAAAGG	HFgltR	ATCAACYCTATCTGSAGTTCCATCAT		P	P	
		HYOgltF	TATCAGTTTTATAGATGGTGAAAAGG	HYOgltR	YCTATCTGGAGTCCATCATCATYCAAG			A	
		LANgltF	ATGCATAGMGGMTATGATAGCGTGG	LANgltR	CATCAACTCTATCTGGAGTWCKKATCA				P
		HFgltXF	TACTGGTATGTTTACRTTTGATAGGGG	HFgltXR	GAATCMACATTTGGATATAPATTTCT				A
		SPUTgltF	AAAAAAGCATATAAAAACATCGTGG	SPUTgltR	TTATCCACACTTCCATCAITTTCTAGTT				P

(Continued)

Table 1 | Continued

Locus	Allele size (bp)	Oligonucleotide primers				Method			
		Forward (5'-3')	Reverse (5'-3')	Co	Cv	Cf	Ch	Cl	Cs
glyA	507	glyCCCFN1	ACAAACAATACGCMGAAGGCTA	glyCCCRN1	GATATCWGCRCTTTWCCGCTAAA	P	P		
		glyCCF1	AARSAGCTTTTGGMTGCGAA	glyCCR1	GATRTCWGCCTTTTCCGCTAA	A	A		
		HFglyF	GCAAYGTTTCAGCCAAATAGC	HFglyR	TTTATTACTGTATTCCWGCRTTACC		P	P	
		HYOglyF	TTGCWAATGTTTCAGCCAAATAGC	HYOglyR	ATTCRRGCATTRCCAAGAGCGATAT				A
		LANGlyF	TGCWAATGTTTCAGCCAAATAGCG	LANGlyR	CAAGAGCGGATATCRGCRCTTTTACC				P
		HF _L glyXF	GATWGTAGTGARAAYTTCACATATCC	HF _L glyXR	GYCCTCTTTCATACCYCTTGC				A
ilvD	492	SPU _T glyF	TGCAAAATGTTCAACCAACTCAG	SPU _T glyR	GTAATTCAGCAATTCCTAAAGCAT	P	P		P
		ilvCCCFN1	CGACTGGGCTAAAAGACGAGGA	ilvCCCRN1	TATGTGAGCGATRTTYGGCTGAT	A	A		
pgm	501	ilvCCF1	CGACYGGGCTAAAAGACGAGGAC	ilvCCR1	CKATGTGAGCGATGTTTTGGCTGAT	P	P		
		pgmCCCFN1	CARATMAAAAATTCHTCCCAAAAAGAG	pgmCCCRN1	CTTTTABGGCATTARGGCTTTTAYRAA	P	P		
		pgmCCF1	ATCAAAAATTCMTCCCAAAAAGAGC	pgmCCR1	ATCAMRTTTCRGRCCAGAGATCTAAA	A	A		
		HFpgmF	AAAAGGTTTRMGAGTTGTTTTGGACGT	HFpgmR	TAACGTTTTCWGTVCCWGAATATCTAAA		P	P	
		HYOpgmF	GCTTACCTTAAAAGGTTTRCGAGTTGT	HYOpgmR	TCATCAGTCTTCAAGCAAAAAG				A
		LANpgmF	GCTTACYTTAAAAGGCCCTRMGAGTTGT	LANpgmR	AAGAAGCAGYCTAATCAAAATYCTGT				P
		HF _L pgmXF	CAATMGCRTTTTTAACCCGAAGATATG	HF _L pgmXR	AAATTTTCYACYCTTCCATYTTTTTA				A
		SPU _T pgmF	TTCCAAAAGAGCTTACAATGTATGG	SPU _T pgmR	TGTTCCCTGAATATCTAAATAGTGAGCG				P
		HFtktF	TTTTTRTGTGCVGATATGTTCAAAA	HFtktR	TATGATWCTTCRCCCAAGMGAGC		P	P	
		HYOtktF	CAATGGGACTTGCTGATTTAATGAG	HYOtktR	TCTTTGCMTCITTTATGATATCTCCG				A
tkt	459, 468 (Cs)	HF _L tktXF	AATAAGATTTTTRTGTGCVGATATGGT	HF _L tktXR	AAGAGTGAATTTARMAGCTCTTTTTTA				P
		LANtktF	CATCTAAAACAYAAATCCMAAAAATCC	LANtktR	ATCTCWKCGCCAAGMGAGC				A
		SPU _T tktF	TTGGGATATATGGTTGTTTTAGC	SPU _T tktR	GATTAATCTCGCGAGTTTTTTTTGC				P

All tkt alleles are 459 bp except for *C. sputorum* (468 bp). Forward and reverse primers were used in both PCR amplification and subsequent amplicon sequencing.

Co, *C. concisus*; Cv, *C. curvus*; Cf, *C. fetus* (both subspecies); Ch, *C. hyointestinalis* (both subspecies); Cl, *C. lariensis*; Cs, *C. sputorum*; P, primary MLST primer; A, alternate MLST primer.

Table 2 | Source of the *Campylobacter* strains typed in this study.

Species	Subspecies	Strains	Human	Cow/cattle	Pig/feral swine	Other/unknown
<i>concisus</i>		70	70	0	0	0
<i>curvus</i>		16	16	0	0	0
<i>fetus</i>		21	6	4	8	3
<i>hyointestinalis</i>	<i>hyointestinalis</i>	39	14	16	9	0
<i>hyointestinalis</i>	<i>lawsonii</i>	9	0	1	8	0
<i>lanienae</i>		34	1	0	32	1
<i>sputorum</i>		24	2	9	8	5

selection. Thus, the ratio of non-synonymous to synonymous base substitutions (d_n/d_s) would be an indicator of potential positive selection: ratios > 1 would be evidence of possible positive selection, whereas ratios < 1 would be more indicative of purifying or stabilizing selection. The d_n/d_s values for previous *Campylobacter* methods were quite low: the highest d_n/d_s values for *C. jejuni* (0.093), *C. coli* (0.173), *C. lari* (0.047), *C. upsaliensis* (0.097), and *C. insulaenigrae* (0.110) were substantially < 1 (Colles et al., 2003; Miller et al., 2005; Stoddard et al., 2007). Similar ratios (highest d_n/d_s values) calculated in this study for *C. concisus* (0.0295), *C. curvus* (0.0468), *Chh* (0.0516), *Chl* (0.0655), *C. lanienae* (0.0562), and *C. sputorum* (0.0426; **Table 4B**) are consistent with the previous methods, indicating that these MLST loci are also not subject to positive selection.

IDENTIFICATION OF PUTATIVE LATERAL GENE TRANSFER EVENTS AND NOVEL TAXA

Characterization of the *Campylobacter* MLST methods also identified putative lateral gene transfer events. An allele that was nearly identical to, and clustered phylogenetically with, alleles from another taxon was determined to represent a putative lateral gene transfer event. For example, strains RM14410 and RM14403 in the *Chh* strain set contain alleles *pgm*-14 and *pgm*-16, respectively (Table S1 in Supplementary Material), that are 97.8–99.6% similar at the nt level to *pgm* alleles from *Chl* but only 95.2–96.2% similar at the nt level to *pgm* alleles from *Chh* (data not shown); these two alleles are also clearly related to other *Chl* *pgm* alleles, based on phylogenetic analysis (**Figure 4A**). Strains RM14410 and RM14403 also contain the *atpA* alleles *atpA*-2 and *atpA*-13, respectively, that cluster with other *Chh* *atpA* alleles (**Figure 4B**). Alleles at the other five loci for these two strains cluster also with other *Chh* alleles (data not shown), indicating that RM14403 and RM14410 are *Chh* strains in which a putative lateral transfer event has occurred at the *pgm* locus. The extent of gene transfer in these two strains could not be determined by MLST and will require further genome sequence analyses. It is not surprising that *Chl* alleles were discovered in *Chh* strains: these two taxa are highly related, being subspecies of the same species, and are isolated often from the same food animals (i.e., cattle and swine). No putative lateral transfer events were observed within *C. concisus* or *C. curvus* and no alleles were identified that originated tentatively in another species.

In some instances, phylogenetically divergent alleles within a strain set were indicative of either highly divergent strains or perhaps novel taxa. Here, as in other MLST studies (Miller et al.,

2005, 2009), putative lateral gene transfer events were identified at only one of the seven MLST loci. For example, *Chh* STs containing *pgm*-14 or *pgm*-16 (ST-26_{hh} and ST-28_{hh}, respectively; **Figure 2**) were divergent at only the *pgm* locus. The alleles for each of the remaining six loci within each ST were of likely *Chh* origin, and phylogenetic analysis of the concatenated allele sequences clearly placed these two STs within the *Chh* clade. However, some STs (ST-1_{lan}, ST-4_{lan}, ST-8_{sp}, ST-13_{sp}, ST-14_{sp}, and ST-15_{sp}) contain three to seven variant alleles (Table S1 in Supplementary Material). The *C. lanienae* STs ST-1_{lan} and ST-4_{lan} differ substantially from the other typed *C. lanienae* strains: concatenated nucleotide sequences across all seven loci for ST-1_{lan} and ST-4_{lan} are on average only approx. 92 and 88% similar, respectively, to the concatenated sequences of the other *C. lanienae* STs (**Figure 2**), which display an average 98% cross-similarity (data not shown). Therefore, these two *C. lanienae* STs may be exemplars of novel *C. lanienae*-related taxa (for comparison, the concatenated *Chh* nucleotide sequences are approx. 94% similar to those of the other *C. hyointestinalis* subspecies and 87% similar to those of the related species *C. fetus*; **Figure 2**). Additionally, six phylogenetically related, urease-negative strains of *C. sputorum*, all isolated from cattle over a 19-month time period, may be members of a novel taxon. Within these six strains, four divergent ST were identified (ST-8_{sp}, ST-13_{sp}, ST-14_{sp}, and ST-15_{sp}). The concatenated allele sequences of these four STs are 95% similar to STs from the three established *C. sputorum* biovars, that display only 1% sequence divergence across the 3321-bp (**Figure 2**). Thus, it is possible that these six strains are members of a *C. sputorum*-like taxon, perhaps a novel *C. sputorum* subspecies or biovar. Nevertheless, for both the divergent *C. lanienae* and *C. sputorum* strains, additional biochemical and molecular tests will need to be performed to definitively establish their taxonomic position within *Campylobacter*.

SUBTYPING OF *C. CONCISUS* AND *C. SPUTORUM* STRAINS

Previous studies investigating the diversity of *C. concisus* organized strains from this species into two major genetically diverse clusters or genomospecies (GS), based on strain typing using 23S rRNA PCR (Engberg et al., 2005; Kalischuk and Inglis, 2011) or AFLP (Aabenhus et al., 2005; Kalischuk and Inglis, 2011). Included in the *C. concisus* strain set here were several strains characterized previously by AFLP (Aabenhus et al., 2005). In agreement with these previous studies, phylogenetic analysis of the *C. concisus* STs identified two clusters: each cluster contained almost exclusively GS1 or GS2 strains (**Figure 1**). Within *C. concisus*, two to eight






FIGURE 2 | Dendrogram of *C. hyointestinalis*, *C. fetus*, *C. lanienae*, and *C. sputorum* STs. Allele sequences for each strain were concatenated in the order *aspA-atpA-glnA-gltA-glyA-pgm-tkt* and aligned using CLUSTALX. The dendrogram was constructed using the neighbor-joining algorithm and the Kimura two-parameter distance estimation method (Kimura, 1980). Bootstrap values >75%, generated from 1000 replicates, are shown at the nodes. Scale bar represents substitutions per site. ST labels indicate taxon: hh, *C. hyointestinalis* subsp. *hyointestinalis*; hl, *C. hyointestinalis* subsp. *lawsonii*; ff, *C. fetus* subsp. *fetus*; fv, *C. fetus* subsp. *venerealis*; lan, *C. lanienae*; sp, *C. sputorum*; sp (sp), *C. sputorum* bv. *sputorum*; sp (f), *C. sputorum* bv. *fecalis*; sp (p), *C. sputorum* bv. *paraureolyticus*. Arrows indicate the STs of the genome-sequenced strains for each taxon. *Urease-positive strains. Values in parentheses at the nodes represent the average %nt similarity of the STs split at each node, following pairwise comparisons of the concatenated allele sequences.

alleles at each of the seven MLST loci were identified in more than one ST (Table S1 in Supplementary Material). It is noteworthy perhaps that of these 33 “common” MLST alleles, only one

Table 3 | Sequence types and alleles identified by the novel MLST methods.

Species	Subspecies	Strains	STs	Alleles								
				<i>aspA</i>	<i>atpA</i>	<i>glnA</i>	<i>gltA</i>	<i>glyA</i>	<i>ilvD</i>	<i>pgm</i>	<i>tkt</i>	
<i>concisus</i>		70	66	60	55	62	59	61	64	59	N/A	
<i>curvus</i>		16	11	8	7	6	6	7	9	7	N/A	
<i>fetus</i>		21	5	1	4	2	2	2	N/A	1	2	
<i>hyointestinalis</i>	<i>hyointestinalis</i>	39	31	8	6	7	5	12	N/A	12	13	
<i>hyointestinalis</i>	<i>lawsonii</i>	9	8	7	8	7	6	7	N/A	7	7	
<i>laniena</i>		34	26	16	9	13	11	13	N/A	16	12	
<i>sputorum</i>		24	16	6	6	3	6	9	N/A	6	7	



allele (*glyA*-31) was identified in both GS1 and GS2 strains, suggesting that minimal genetic exchange occurs between the two genomospecies.

Division of *C. concisus* strains into two primary genogroups is not merely an academic exercise. In a study analyzing *C. concisus* strains isolated from diarrheic and non-diarrheic individuals, Kalischuk and Inglis reported that GS1 strains were isolated predominantly from healthy individuals while the GS2 cluster contained isolates primarily from diarrheic individuals (Kalischuk and Inglis, 2011); this correlation between diarrheal disease and GS2 *C. concisus* species was also observed by Aabenhus et al. (2005). Moreover, GS2 strains were reported to exhibit higher levels of epithelial invasion (Kalischuk and Inglis, 2011). Therefore, GS1 or GS2 strains would be predicted to lead possibly to different clinical outcomes, and the proper placement of *C. concisus* isolates into these two genomospecies would be critical not only clinically

but also for epidemiological studies. The MLST method described here provides another accurate tool for *C. concisus* genotyping.

Campylobacter concisus GS4 strains were isolated from severely immunodeficient patients and identified initially by limited phenotyping; comparative AFLP analysis and other DNA-based testing indicated their taxonomic position required clarification (Aabenhus et al., 2005). Given these data, and our MLST results, it is likely that these strains are *C. curvus* and not *C. concisus*. These species share many phenotypic traits (On et al., 1996) and are difficult to distinguish with limited phenotypic testing.

At present, *C. sputorum* biovars are identified by their catalase or urease phenotypes (On et al., 1998). No PCR/sequencing methods exist that differentiate the three described biovars, although whole-genome macro-restriction profiling has been used to suggest biovar clonality (On et al., 1999). However, putative biovar-associated alleles were observed at three of the *C. sputorum* MLST loci (Table S1 in Supplementary Material). For example, at the *atpA* locus, bv. *sputorum* strains are *atpA*-1, bv. *fecalis* strains are *atpA*-3, and bv. *paraureolyticus* strains are either *atpA*-2 or *atpA*-4; strains of the cow-associated *C. sputorum* clade described above are either *atpA*-5 or *atpA*-6. Similar associations exist at the *glyA* and *pgm* loci. Indeed, the three biovars segregate also when all seven loci are examined phylogenetically (Figure 2), although the differences are quite small. Obviously, the size of the *C. sputorum* strain set typed here is too small to reach any definitive conclusions, although the associations are intriguing.


CONCLUSION

The four MLST methods described in this study typed successfully all 213 *Campylobacter* strains, representing at least ten *Campylobacter* taxa that included both subspecies of *C. fetus* and *C. hyointestinalis* and all three biovars of *C. sputorum*. Each method identified multiple novel STs; the small number of STs present in some taxa were more likely due to the limited size of the sample sets for those taxa than a limitation of the method itself. These methods were successful despite the high degree of variation in some species, e.g., *C. concisus*. Also, the concatenated *C. concisus* and *C. curvus* ST sequences were only 81% similar at the nt level (Figure 1); likewise, the *C. hyointestinalis* and *C. fetus* ST sequences were 87% similar (Figure 2), yet MLST Methods 1 and 2 could type either set of strains unequivocally. Indeed, the ability to sequence such variable strains provided an unexpected bonus to these MLST methods. The methods described here identified putative novel *C.*

Table 4 | Diversity within the *Campylobacter* MLST loci.

Species	Subspecies	Strains	<i>aspA</i>	<i>atpA</i>	<i>glnA</i>	<i>gltA</i>	<i>glyA</i>	<i>ilvD</i>	<i>pgm</i>	<i>tkt</i>
A. POLYMORPHIC SITES										
<i>concisus</i>		70	131	105	120	102	129	140	128	N/A
<i>curvus</i>		16	90	47	47	51	40	63	41	N/A
<i>hyointestinalis</i>	<i>hyointestinalis</i>	39	9	9	10	27	15	N/A	30 (17)	24
<i>hyointestinalis</i>	<i>lawsonii</i>	9	23	36	10	11	64	N/A	19	21
<i>lanienae</i>		34	67 (29)	86 (51)	58 (19)	57 (26)	71 (26)	N/A	142 (31)	96 (38)
<i>sputorum</i>		24	58 (5)	33 (17)	24 (1)	28 (9)	39 (14)	N/A	18 (10)	20 (8)
B. RATIOS OF NON-SYNONYMOUS (d_n) TO SYNONYMOUS (d_s) BASE SUBSTITUTIONS										
<i>concisus</i>		70	0.0257	0.0028	0.0052	0.0036	0.017	0.0295	0.0095	N/A
<i>curvus</i>		16	0.0417	0.0057	0.008	0.0468	0.0168	0.0149	0.0091	N/A
<i>hyointestinalis</i>	<i>hyointestinalis</i>	39	0	0	0	0.0262	0.0516	N/A	0.0249 (0.0110)	0.0381
<i>hyointestinalis</i>	<i>lawsonii</i>	9	0.0169	0.0276	0.0216	0	0.0113	N/A	0.0655	0.0251
<i>lanienae</i>		34	0.015 (0.0149)	0.0204 (0.0235)	0.0026 (0)	0.0502 (0.0298)	0.0112 (0.0257)	N/A	0.0562 (0.0748)	0.0421 (0.0419)
<i>sputorum</i>		24	0.041 (0.0715)	0 (0)	0 (0)	0 (0)	0.0102 (0.0511)	N/A	0.0293 (0.0264)	0.0426 (0.1672)

Numbers in parentheses represent polymorphic sites and d_n/d_s ratios recalculated following removal of the divergent *pgm*-14 and *pgm*-16 alleles within the *Chh* profiles and removal of the divergent *ST*-1_{lan} and *ST*-4_{lan} (*lanienae*) and *ST*-8_{sp}, *ST*-13_{sp}, *ST*-14_{sp}, and *ST*-15_{sp} (*sputorum*) sequence types.



lanienae- and *C. sputorum*-related taxa and it is likely that these methods could further characterize and type as-yet-undescribed *Campylobacter* species or subspecies. For example, MLST Method 2 has been used to type reptile-associated *C. fetus*-like organisms (data not shown).

For many campylobacters, sequence data is restricted currently to ribosomal rRNA loci. While these rDNA sequences can provide crucial speciation data for many taxa, some groups of campylobacters cannot be differentiated readily by 16S rDNA sequencing. One such example includes *C. hyointestinalis* and *C. lanienae* strains.

Some of the strains in this study from these species could not be typed unequivocally by 16S rDNA sequencing; however, MLST could readily place all strains in their proper taxonomic positions. MLST has been shown also to be of value in identifying strains of species with multiple phenogroups, such as *C. insulaenigrae* (Stoddard et al., 2007). In this study, a *C. sputorum* clade was typed that, based on established phenotypic characterization, would likely have been classified as *bv. fecalis*. While additional tests need to be performed, MLST cast some doubt that these strains were *C. sputorum* *bv. fecalis*.

Eighteen of the thirty validly described *Campylobacter* taxa can now be typed by MLST. This number is likely an underestimate, as some of the *C. lari*-like species (e.g., *C. peloridis*) described recently can be typed also using the *C. lari* MLST method (data not shown). The ability of MLST to type and speciate campylobacters, as well as identify putative horizontal gene transfer, indicates that the multiple *Campylobacter* MLST methods now available provide a valuable tool in the epidemiology, typing, and evolution of emerging campylobacters.

ACKNOWLEDGMENTS

We thank Nate Simon for technical assistance in the *C. hyointestinalis* strain typing. We thank Dr. Birgitta Duim for critical reading of this manuscript. We thank also S. Chandler, D. Orthmeyer, G. Baird-Kavanaugh, and S. Hee for providing some of the wildlife and environmental samples; and E. R. Atwill, M.

Jay-Russell for their support in projects that supported partially this research and funded by National Research Initiative Competitive Grant nos. 2006-55212-16927 and 2007-35212-18239 from the USDA National Institute of Food and Agriculture. This work was supported mostly by USDA Agricultural Research Service CRIS projects 5325-42000-044 and -045. This publication made use of the *Campylobacter* Multilocus Sequence Typing website (<http://pubmlst.org/campylobacter/>) developed by Keith Jolley at the University of Oxford (Jolley and Maiden, 2010).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at http://www.frontiersin.org/Cellular_and_Infection_Microbiology/10.3389/fcimb.2012.00045/abstract

Table S1 | *Campylobacter* strains typed in this study.

REFERENCES

- Aabenhus, R., On, S. L., Siemer, B. L., Permin, H., and Andersen, L. P. (2005). Delineation of *Campylobacter concisus* genomospecies by amplified fragment length polymorphism analysis and correlation of results with clinical data. *J. Clin. Microbiol.* 43, 5091–5096.
- Anonymous. (2009). Preliminary Food-Net Data on the incidence of infection with pathogens transmitted commonly through food – 10 States, 2008. *MMWR Morb. Mortal. Wkly. Rep.* 58, 333–337.
- Bullman, S., O’Leary, J., Corcoran, D., Sleator, R. D., and Lucey, B. (2011). Molecular-based detection of non-culturable and emerging campylobacteria in patients presenting with gastroenteritis. *Epidemiol. Infect.* 140, 684–688.
- Colles, F. M., Jones, K., Harding, R. M., and Maiden, M. C. (2003). Genetic diversity of *Campylobacter jejuni* isolates from farm animals and the farm environment. *Appl. Environ. Microbiol.* 69, 7409–7413.
- Debruyne, L., Gevers, D., and Vandamme, P. (2008). “Taxonomy of the family *Campylobacteraceae*,” in *Campylobacter*, 3rd Edn, eds I. Nachamkin, C. M. Szymanski, and M. J. Blaser (Washington, DC: ASM Press), 3–25.
- Dingle, K. E., Colles, F. M., Falush, D., and Maiden, M. C. (2005). Sequence typing and comparison of population biology of *Campylobacter coli* and *Campylobacter jejuni*. *J. Clin. Microbiol.* 43, 340–347.
- Dingle, K. E., Colles, F. M., Wareing, D. R. A., Ure, R., Fox, A. J., Bolton, F. E., Bootsma, H. J., Willems, R. J. L., Urwin, R., and Maiden, M. C. J. (2001). Multilocus sequence typing system for *Campylobacter jejuni*. *J. Clin. Microbiol.* 39, 14–23.
- Edmonds, P., Patton, C. M., Griffin, P. M., Barrett, T. J., Schmid, G. P., Baker, C. N., Lambert, M. A., and Brenner, D. J. (1987). *Campylobacter hyointestinalis* associated with human gastrointestinal disease in the United States. *J. Clin. Microbiol.* 25, 685–691.
- Engberg, J., Bang, D. D., Aabenhus, R., Aarestrup, F. M., Fussing, V., and Gerner-Smidt, P. (2005). *Campylobacter concisus*: an evaluation of certain phenotypic and genotypic characteristics. *Clin. Microbiol. Infect.* 11, 288–295.
- Gorkiewicz, G., Feierl, G., Zechner, R., and Zechner, E. L. (2002). Transmission of *Campylobacter hyointestinalis* from a pig to a human. *J. Clin. Microbiol.* 40, 2601–2605.
- Hakkinen, M., Heiska, H., and Hanninen, M. L. (2007). Prevalence of *Campylobacter* spp. in cattle in Finland and antimicrobial susceptibilities of bovine *Campylobacter jejuni* strains. *Appl. Environ. Microbiol.* 73, 3232–3238.
- Inglis, G. D., Kalischuk, L. D., and Busz, H. W. (2004). Chronic shedding of *Campylobacter* species in beef cattle. *J. Appl. Microbiol.* 97, 410–420.
- Jolley, K. A., Feil, E. J., Chan, M. S., and Maiden, M. C. (2001). Sequence type analysis and recombinational tests (START). *Bioinformatics* 17, 1230–1231.
- Jolley, K. A., and Maiden, M. C. (2010). BIGSdb: scalable analysis of bacterial genome variation at the population level. *BMC Bioinformatics* 11, 595. doi:10.1186/1471-2105-11-595
- Kalischuk, L. D., and Inglis, G. D. (2011). Comparative genotypic and pathogenic examination of *Campylobacter concisus* isolates from diarrheic and non-diarrheic humans. *BMC Microbiol.* 11, 53. doi:10.1186/1471-2180-11-53
- Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.* 16, 111–120.
- Lane, D. J. (1991). “16S/23S rRNA sequencing,” in *Nucleic Acid Sequencing Techniques in Bacterial Systematics*, eds E. Stakebrandt and M. Goodfellow (New York, NY: Wiley and Sons), 115–175.
- Lastovica, A. J., and Allos, B. M. (2008). “Clinical significance of *Campylobacter* and related species other than *Campylobacter jejuni* and *Campylobacter coli*,” in *Campylobacter*, 3rd Edn, eds I. Nachamkin, C. M. Szymanski, and M. J. Blaser (Washington, DC: ASM Press), 123–149.
- Lynch, O. A., Cagney, C., McDowell, D. A., and Duffy, G. (2011). Occurrence of fastidious *Campylobacter* spp. in fresh meat and poultry using an adapted cultural protocol. *Int. J. Food Microbiol.* 150, 171–177.
- Mahendran, V., Riordan, S. M., Grimm, M. C., Tran, T. A., Major, J., Kaakoush, N. O., Mitchell, H., and Zhang, L. (2011). Prevalence of *Campylobacter* species in adult Crohn’s disease and the preferential colonization sites of *Campylobacter* species in the human intestine. *PLoS ONE* 6, e25417. doi:10.1371/journal.pone.0025417
- Maiden, M. C. (2006). Multilocus sequence typing of bacteria. *Annu. Rev. Microbiol.* 60, 561–588.
- Maiden, M. C., and Dingle, K. E. (2008). “Population biology of *Campylobacter jejuni* and related organisms,” in *Campylobacter*, 3rd Edn, eds I. Nachamkin, C. Szymanski, and M. J. Blaser (Washington, DC: ASM Press), 27–40.
- Man, S. M., Zhang, L., Day, A. S., Leach, S. T., Lemberg, D. A., and Mitchell, H. (2010). *Campylobacter concisus* and other *Campylobacter* species in children with newly diagnosed Crohn’s disease. *Inflamm. Bowel Dis.* 16, 1008–1016.
- Manning, G., Dowson, C. G., Bagnall, M. C., Ahmed, I. H., West, M., and Newell, D. G. (2003). Multilocus sequence typing for comparison of veterinary and human isolates of *Campylobacter jejuni*. *Appl. Environ. Microbiol.* 69, 6370–6379.
- Miller, W. G., On, S. L., Wang, G., Fontanoz, S., Lastovica, A. J., and Mandrell, R. E. (2005). Extended multilocus sequence typing system for *Campylobacter coli*, *C. lari*, *C. upsaliensis*, and *C. helveticus*. *J. Clin. Microbiol.* 43, 2315–2329.
- Miller, W. G., Wesley, I. V., On, S. L., Houf, K., Megraud, F., Wang, G., Yee, E., Srijan, A., and Mason, C. J. (2009). First multi-locus sequence typing scheme for *Arcobacter* spp. *BMC Microbiol.* 9, 196. doi:10.1186/1471-2180-9-196
- Mølbaek, K., and Havelaar, A. (2008). “Burden of illness of campylobacteriosis and sequelae,” in *Campylobacter*, 3rd Edn, eds I. Nachamkin, C. M. Szymanski, and M. J. Blaser (Washington, DC: ASM Press), 151–162.
- Mukhopadhyaya, I., Thomson, J. M., Hansen, R., Berry, S. H., El-Omar, E. M., and Hold, G. L. (2011). Detection of *Campylobacter concisus* and other *Campylobacter* species in colonic biopsies from adults with ulcerative colitis. *PLoS ONE* 6, e21490. doi:10.1371/journal.pone.0021490

- Olson, C. K., Ethelberg, S., Van Pelt, W., and Tauxe, R. V. (2008). "Epidemiology of *Campylobacter jejuni* infections in industrialized nations," in *Campylobacter*, 3rd Edn, eds I. Nachamkin, C. M. Szymanski, and M. J. Blaser (Washington, DC: ASM Press), 163–189.
- On, S. L., Atabay, H. I., and Corry, J. E. (1999). Clonality of *Campylobacter sputorum* bv. paraureolyticus determined by macrorestriction profiling and biotyping, and evidence for long-term persistent infection in cattle. *Epidemiol. Infect.* 122, 175–182.
- On, S. L., Atabay, H. I., Corry, J. E., Harrington, C. S., and Vandamme, P. (1998). Emended description of *Campylobacter sputorum* and revision of its infrasubspecific (biovar) divisions, including *C. sputorum* biovar paraureolyticus, a urease-producing variant from cattle and humans. *Int. J. Syst. Bacteriol.* 48(Pt 1), 195–206.
- On, S. L., Holmes, B., and Sackin, M. J. (1996). A probability matrix for the identification of campylobacters, helicobacters and allied taxa. *J. Appl. Bacteriol.* 81, 425–432.
- On, S. L. W., McCarthy, N., Miller, W. G., and Gilpin, B. J. (2008). "Molecular epidemiology of *Campylobacter* species," in *Campylobacter*, 3rd Edn, eds I. Nachamkin, C. Szymanski, and M. J. Blaser (Washington, DC: ASM Press), 191–212.
- Oporto, B., and Hurtado, A. (2011). Emerging thermotolerant *Campylobacter* species in healthy ruminants and swine. *Foodborne Pathog. Dis.* 8, 807–813.
- Perez-Losada, M., Porter, M. L., Viscidi, R. P., and Crandall, K. A. (2011). "Multilocus sequence typing of pathogens," in *Genetics and Evolution of Infectious Diseases*, ed. M. Tibayrenc (Burlington, MA: Elsevier), 503–522.
- Salihu, M. D., Abdulkadir, J. U., Oboegbulem, S. I., Egwu, G. O., Magaji, A. A., Lawal, M., and Hassan, Y. (2009). Isolation and prevalence of *Campylobacter* species in cattle from Sokoto state, Nigeria. *Vet. Ital.* 45, 501–505.
- Sasaki, Y., Fujisawa, T., Ogikubo, K., Ohzono, T., Ishihara, K., and Takahashi, T. (2003). Characterization of *Campylobacter lanienae* from pig feces. *J. Vet. Med. Sci.* 65, 129–131.
- Stoddard, R. A., Miller, W. G., Foley, J. E., Lawrence, J., Gulland, F. M., Conrad, P. A., and Byrne, B. A. (2007). *Campylobacter insulaenigrae* isolates from northern elephant seals (*Mirounga angustirostris*) in California. *Appl. Environ. Microbiol.* 73, 1729–1735.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., and Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–2739.
- Terzolo, H. R. (1988). Identification of campylobacters from bovine and ovine faeces. *Rev. Argent. Microbiol.* 20, 53–68.
- van Bergen, M. A., Dingle, K. E., Maiden, M. C., Newell, D. G., Van Der Graaf-Van Bloois, L., Van Putten, J. P., and Wagenaar, J. A. (2005). Clonal nature of *Campylobacter fetus* as defined by multilocus sequence typing. *J. Clin. Microbiol.* 43, 5888–5898.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 05 December 2011; accepted: 16 March 2012; published online: 02 April 2012.

Citation: Miller WG, Chapman MH, Yee E, On SLW, McNulty DK, Lastovica AJ, Carroll AM, McNamara EB, Duffy G and Mandrell RE (2012) Multilocus sequence typing methods for the emerging *Campylobacter* species *C. hyointestinalis*, *C. lanienae*, *C. sputorum*, *C. concisus*, and *C. curvus*. *Front. Cell. Inf. Microbio.* 2:45. doi: 10.3389/fcimb.2012.00045
Copyright © 2012 Miller, Chapman, Yee, On, McNulty, Lastovica, Carroll, McNamara, Duffy and Mandrell. This is an open-access article distributed under the terms of the Creative Commons Attribution Non Commercial License, which permits non-commercial use, distribution, and reproduction in other forums, provided the original authors and source are credited.