



Identification and Roles of Photosystem II Assembly, Stability, and Repair Factors in Arabidopsis

Yan Lu*

Department of Biological Sciences, Western Michigan University, Kalamazoo, MI, USA

OPEN ACCESS

Edited by:

Julian Eaton-Rye,
University of Otago, New Zealand

Reviewed by:

Biswapriya Biswas Misra,
University of Florida, USA
Peter Julian Nixon,
Imperial College London, UK

*Correspondence:

Yan Lu
yan.1.lu@wmich.edu

Specialty section:

This article was submitted to
Plant Cell Biology,
a section of the journal
Frontiers in Plant Science

Received: 18 November 2015

Accepted: 31 January 2016

Published: 16 February 2016

Citation:

Lu Y (2016) Identification and Roles of Photosystem II Assembly, Stability, and Repair Factors in Arabidopsis. *Front. Plant Sci.* 7:168. doi: 10.3389/fpls.2016.00168

Photosystem II (PSII) is a multi-component pigment-protein complex that is responsible for water splitting, oxygen evolution, and plastoquinone reduction. Components of PSII can be classified into core proteins, low-molecular-mass proteins, extrinsic oxygen-evolving complex (OEC) proteins, and light-harvesting complex II proteins. In addition to these PSII subunits, more than 60 auxiliary proteins, enzymes, or components of thylakoid protein trafficking/targeting systems have been discovered to be directly or indirectly involved in *de novo* assembly and/or the repair and reassembly cycle of PSII. For example, components of thylakoid-protein-targeting complexes and the chloroplast-vesicle-transport system were found to deliver PSII subunits to thylakoid membranes. Various auxiliary proteins, such as PsbP-like (Psb stands for PSII) and light-harvesting complex-like proteins, atypical short-chain dehydrogenase/reductase family proteins, and tetratricopeptide repeat proteins, were discovered to assist the *de novo* assembly and stability of PSII and the repair and reassembly cycle of PSII. Furthermore, a series of enzymes were discovered to catalyze important enzymatic steps, such as C-terminal processing of the D1 protein, thiol/disulfide-modulation, peptidylprolyl isomerization, phosphorylation and dephosphorylation of PSII core and antenna proteins, and degradation of photodamaged PSII proteins. This review focuses on the current knowledge of the identities and molecular functions of different types of proteins that influence the assembly, stability, and repair of PSII in the higher plant *Arabidopsis thaliana*.

Keywords: Photosystem II assembly, Photosystem II stability, Photosystem II repair, *Arabidopsis thaliana*, identification and roles

INTRODUCTION

Photosystem II (PSII) is a multi-subunit pigment-protein complex found in thylakoid membranes of oxygenic photosynthetic organisms, including cyanobacteria, algae, and plants (Nickelsen and Rengstl, 2013; Järvi et al., 2015). Driven by light, PSII catalyzes electron transfer from water to plastoquinone. Therefore, PSII is also known as a water-plastoquinone oxidoreductase. Proteomics, X-ray crystallography, and single-particle electron cryo-microscopy studies revealed that PSII components include core proteins, low-molecular-mass (LMM, i.e., <10 kDa) proteins, extrinsic oxygen-evolving complex (OEC) proteins, and light-harvesting complex (LHC) proteins (da Fonseca et al., 2002; Kashino et al., 2002; Liu et al., 2004; Aro et al., 2005; Nield and Barber, 2006; Umena et al., 2011; Suga et al., 2015). Except for some minor differences in the composition of LMM proteins, the core of PSII is conserved from cyanobacteria to land plants


(Umena et al., 2011; Nickelsen and Rengstl, 2013). Proteins that form the PSII core complex in land plants include PSII reaction center core proteins D1 and D2 (i.e., PsbA and PsbD; Psb stands for PSII), core antenna proteins CP43 and CP47 (i.e., PSII chlorophyll proteins of 43 and 47 kDa, also known as PsbC and PsbB, respectively), cytochrome *b*₅₅₉ subunits alpha and beta (i.e., PsbE and PsbF), and LMM proteins PsbH, PsbI, PsbJ, PsbK,

Abbreviations: ALB3, Albino3; BN-PAGE, blue native-polyacrylamide gel electrophoresis; CP43, Photosystem II chlorophyll protein of 43 kDa, also known as PsbC; CP47, Photosystem II chlorophyll protein of 47 kDa, also known as PsbB; CPRabA5e, chloroplast Rab GTPase A5e; cpFtsY, filamentation temperature sensitive protein Y; cpSec, chloroplast secretory; cpSecA, cpSecE, cpSecY, chloroplast secretory translocase A, E, and Y; cpSRP, chloroplast signal recognition particle; cpSRP43 and cpSRP54, chloroplast Signal Recognition Particle protein of 43 and 54 kDa; cpTat, chloroplast twin-arginine translocation; cpTatC, chloroplast twin-arginine translocation protein C; CtpA1 and CtpA2, C-terminal processing peptidases A 1 and 2; CYO1, Shiyou 1 (Shiyou means cotyledon in Japanese), also known as SCO2; CYP20-3, 20-kDa cyclophilin 3; CYP38, cyclophilin of 38 kDa; Cyt *b*₅₅₉, cytochrome *b*₅₅₉; Deg1; Deg2, Deg5, Deg7, and Deg8, Degradation-of-periplasmic-proteins proteases 1, 2, 5, 7, and 8; FKBP20-2, 20-kDa FK506 (tacrolimus)-binding protein; ELIP1 and ELIP2, early light-induced proteins 1 and 2; FtsH1, FtsH2, FtsH5, FtsH6, FtsH8, FtsH9, FtsH11, and FtsH12, filamentation temperature sensitive proteins H 1, 2, 5, 6, 8, 9, 11, and 12; FtsH2 and FtsH5 are also known as VAR2 and VAR1, respectively; fgl1, fu-gaer1; GTPase, GTP hydrolase; HCF106, HCF136, HCF173, HCF243, and HCF244, High Chlorophyll Fluorescence 106, 136, 173, 243, and 244; HHL1, Hypersensitive to High Light 1; HLIP, high-light-induced protein; LHCI, light-harvesting complex II; LHCB1, LHCB2, LHCB3, LHCB4, LHCB5, and LHCB6, Photosystem II light-harvesting chlorophyll *a/b*-binding proteins 1, 2, 3, 4, 5, and 6; LHCP, light-harvesting chlorophyll *a/b*-binding protein; LIL3:1 and LIL3:2, light-harvesting-like proteins 3:1 and 3:2; LMM, low molecular mass; LPA1, LPA2, LPA3, and LPA19, Low Photosystem II Accumulation 1, 2, 3, and 19; LPA1 is also known as Prata; LQY1, Low Quantum Yield of Photosystem II 1; LTO1, Lumen Thiol Oxidoreductase 1; MET1, Mesophyll-Enriched Thylakoid protein 1; MPH1, Maintenance of Photosystem II under High light 1; OEC, oxygen-evolving complex; OHP1 and OHP2, one-helix proteins 1 and 2; PAM68, Photosynthesis Affected Mutant 68; PBCP, Photosystem II core phosphatase; PBF1, Photosystem Biogenesis Factor 1; pD1, precursor D1; PDI6, Protein Disulfide Isomerase 6, also known as PDIL1-2; PDIase, protein disulfide isomerase; PDIL1-2, Protein Disulfide Isomerase-Like 1-2, also known as PDI6; PPH1, Protein Phosphatase 1; PPIase, Peptidylprolyl Isomerase; PPL1 and PPL2, PsbP-Like proteins 1 and 2; Prata, Processing-associated tetratricopeptide repeat protein A, also known as LPA1; PSI, Photosystem I; PSII, Photosystem II; Psa, Photosystem II; PsaH and PsaK, Photosystem I proteins H and K; Psb, Photosystem II; PSB28, PSB29, and PSB33, Photosystem II proteins 28, 29 and 33; PSB29 is also known as THF1; PsbA, PsbB, PsbC, PsbD, PsbE, PsbF, PsbH, PsbI, PsbJ, PsbK, PsbL, PsbM, PsbN, PsbO, PsbP, PsbQ, PsbR, PsbT, PsbU, PsbV, PsbW, PsbX, PsbY, PsbZ, Photosystem II protein A, B, C, D, E, F, H, I, J, K, L, M, N, O, P, Q, R, T, U, V, W, X, Y and Z; PsbA, PsbB, PsbC, and PsbD are also known as D1, CP47, CP43, and D2; PsbE and PsbF are also known as cytochrome *b*₅₅₉ subunit alpha and beta, respectively; PsbN is also known as PBF1; PsbO, PsbP, and PsbQ are also known as the 33-, 23-, and 17-kDa protein of the oxygen-evolving complex, respectively; PsbTc, chloroplast-encoded Photosystem II protein T; PsbTn, nuclear-encoded PSII protein T; RC, minimal reaction-center complex which lacks CP47 and CP43; RC47a and RC 47b, reaction-center complexes that contain CP47 but lack CP43; RBD1, rubredoxin 1; RNAi, RNA interference; SCO2, Snowy Cotyledon 2, also known as CYO1; SCP, small chlorophyll-binding-like protein; SDR, short-chain dehydrogenase/reductase; SRP, signal recognition particle; STN7 and STN8, state transition 7 and 8; ROC4, rotamase cyclophilin 4; SEP3.1 and SEP3.2, stress-enhanced protein 3.1 and 3.2; TerC, Tellurite-resistance protein C; T-DNA, transfer DNA; TAP38, Thylakoid-Associated Phosphatase of 38 kDa; THA4, Thylakoid Assembly 4; THF1, Thylakoid Formation 1, also known as PSB29; TLP18.3 and TLP40, Thylakoid Lumen Proteins of 18.3 and 40 kDa; TPR, tetratricopeptide repeat; TRX, thioredoxin; VAR1 and VAR2, Yellow Variegated 1 and 2, also known as FtsH5 and FtsH2, respectively; VIPP1, Vesicle-Inducing Protein in Plastids 1; YCF48, hypothetical chloroplast reading frame number 48.

PsbL, PsbM, PsbR, PsbTc (chloroplast-encoded PSII protein T), PsbTn (nuclear-encoded PSII protein T), PsbW, PsbX, PsbY, and PsbZ (Nickelsen and Rengstl, 2013). Due to the loss of PsbU and PsbV during green plant evolution, cyanobacterial OEC has PsbO, PsbP, PsbQ, PsbU, and PsbV subunits but land plant OEC only contains PsbO, PsbP, and PsbQ subunits (Thornton et al., 2004; Bricker et al., 2012). The PSII-light-harvesting antenna in cyanobacteria is made of phycobilisomes, which are attached to the cytoplasmic side of PSII (Liu et al., 2005). The PSII-light-harvesting antenna (i.e., light-harvesting complex II, abbreviated as LHCI) in land plants is an integral membrane complex. LHCI contains three major trimeric PSII light-harvesting chlorophyll *a/b*-binding (LHCB) proteins LHCB1, LHCB2, and LHCB3 and three minor monomeric LHCB proteins LHCB4, LHCB5, and LHCB6 (Jansson, 1999; Liu et al., 2004). In addition to PSII subunits, more than 60 auxiliary proteins or enzymes have been found to be involved in the assembly, stability, and repair of PSII complexes (Nixon et al., 2010; Nickelsen and Rengstl, 2013; Järvi et al., 2015). This article focuses on the identification and roles of different types of proteins that influence the assembly, stability, and repair of PSII in the higher plant *Arabidopsis thaliana*.

DE NOVO ASSEMBLY OF PSII

De novo (Latin for “anew” or “from the beginning”) PSII assembly is a sequential and highly coordinated process. The principal steps were revealed by the use of radioactive pulse-chase experiments, two-dimensional blue native/sodium dodecyl sulfate-polyacrylamide gel electrophoresis, and subsequent proteomics and mass spectrometry analysis (Aro et al., 2005; Rokka et al., 2005; Boehm et al., 2012a). *De novo* PSII assembly in higher plants include: (1) assembly of the precursor D1-PsbI (pD1-PsbI) and D2-cytochrome *b*₅₅₉ (D2-Cyt *b*₅₅₉) precomplexes, (2) assembly of the minimal reaction-center complex (RC), which lacks CP47 and CP43, (3) assembly of the reaction-center complex (RC47a) that contains CP47 but lacks CP43, (4) incorporation of LMM subunits, such as PsbH, PsbM, PsbT, and PsbR, to form RC47b, (5) incorporation of CP43, along with LMM subunit PsbK, to form the OEC-less PSII monomer, (6) assembly of the OEC and additional LMM subunits, such as PsbW and PsbZ, to form the PSII core monomer, and (7) dimerization and formation of the PSII-LHCI supercomplex (Figure 1; Rokka et al., 2005; Nixon et al., 2010; Komenda et al., 2012a; Nickelsen and Rengstl, 2013). A similar pathway exists in cyanobacteria, algae, and lower plants, suggesting that the core components of PSII and the assembly process of PSII complexes are conserved (Nixon et al., 2010; Komenda et al., 2012a; Nickelsen and Rengstl, 2013). From cyanobacteria to green algae to land plants, the initial assembly steps of photosynthetic complexes appear to be spatially separated from sites of active photosynthesis (Nickelsen and Rengstl, 2013). For instance, in *Chlamydomonas reinhardtii*, initial steps of *de novo* PSII assembly occur in discrete regions near the pyrenoid, called translation zones (Uniacke and Zerges, 2007).




Assembly of LHCII is recently thought to initiate on the chloroplast envelope in developing chloroplasts and on thylakoid membranes in developed chloroplasts (Tanz et al., 2012; Khan et al., 2013). LHCII assembly on the chloroplast envelope consists of four major steps: (1) partial insertion of LHCP (light-harvesting chlorophyll *a/b*-binding protein) apoproteins into the inner chloroplast envelope, (2) binding of chlorophyll to reach a stable conformation in the membrane, (3) insertion of the rest of the protein domains, and (4) further pigment binding and protein assembly into a fully assembled pigment-protein complex (Hooper et al., 2007; Dall’Osto et al., 2015). The pigment-protein complexes on the inner chloroplast envelope can be transferred to thylakoid membranes via the chloroplast-vesicle-transport system, the primary source of lipids and proteins for developing thylakoids in young chloroplasts (Tanz et al., 2012; Khan et al., 2013; Karim et al., 2014). In developed chloroplasts, LHCP proteins are primarily transported and integrated into thylakoid membranes via the chloroplast signal recognition particle (cpSRP) pathway (Cline

and Dabney-Smith, 2008; Albinia et al., 2012; Dall’Osto et al., 2015).

DAMAGE, REPAIR, AND REASSEMBLY OF PSII

The PSII repair cycle is a sequential process as well. The major steps in higher plants include: (1) high-light-induced phosphorylation, damage, and disassembly of the PSII-LHCII supercomplex and the PSII core dimer in grana stacks, (2) lateral migration of the PSII core monomer to stroma-exposed thylakoid membranes, (3–5) dephosphorylation, partial disassembly of the PSII core monomer, and degradation of photodamaged D1, (6) synthesis and reassembly of new D1, (7) re-incorporation of CP43, (8) re-attachment of OEC, (9) migration of the PSII core monomer back to grana stacks, and (10) dimerization into PSII core dimers and reformation of PSII-LHCII



supercomplexes (Figure 2; Mulo et al., 2008; Järvi et al., 2015).

PROTEINS THAT INFLUENCE THE ASSEMBLY, STABILITY, AND REPAIR OF PSII

Assistance from a series of protein factors is required for the assembly, stability, and repair of PSII (Figures 1, 2; Mulo et al., 2008; Nickelsen and Rengstl, 2013; Järvi et al., 2015). The types of protein factors include: (1) components of thylakoid-protein-targeting complexes; (2) components of the chloroplast-vesicle-transport system, (3) PSII subunit-like proteins, e.g., PsbP-like and LHCP-like proteins, (4) atypical short-chain dehydrogenase/reductase (SDR) family proteins, (5) C-terminal D1 processing endopeptidases, (6) tetratricopeptide repeat (TPR) proteins, (7) thiol/disulfide-modulating proteins, (8) peptidylprolyl isomerases (PPIases), (9) protein kinases, (10) protein phosphatases, (11) FtsH (filamentation temperature sensitive protein H) proteases, (12) Deg (Degradation of

periplasmic proteins) proteases, and (13) other auxiliary proteins with unique or unknown domain compositions (Table 1). As discussed below, many proteins that are involved in *de novo* PSII assembly also play roles in the repair and reassembly cycle of PSII.

Some of these factors, such as components of the thylakoid-protein-targeting complexes and the chloroplast-vesicle-transport system, are not specific for the assembly, stability, or repair of PSII. However, because thylakoid protein targeting and chloroplast vesicle transport are essential for translocation and accumulation of thylakoid membrane/lumen proteins, and because most of PSII subunits and assembly, stability, and repair factors are thylakoid membrane/lumen proteins, these two types of protein factors are included in this article.

Components of Thylakoid-Protein-Targeting Complexes

Four thylakoid transport and integration pathways have been identified to date (Cline and Dabney-Smith, 2008; Albinak et al., 2012). The cpSRP pathway and an unusual pathway that

TABLE 1 | Summary of proteins that influence the assembly, stability, and repair of PSII in Arabidopsis.

Name	Gene locus in <i>Synechocystis</i> sp. PCC 6803 ^a	Gene locus in Arabidopsis ^b	Full-length Size (kDa)	Mature size (kDa)	Location	Protein classification	Function	References ^c
cpSRP43	–	At2g47450	41	35	CS	Thylakoid protein targeting: cpSRP translocase	Insertion and assembly of PSII proteins such as D1, D2, and CP47, and LHClI subunits	Henry et al., 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Albinaki et al., 2012
cpSRP54	slr1531	At5g03940	61	53	CS	Thylakoid protein targeting: cpSRP translocase	Insertion and assembly of PSII proteins such as D1, D2, and CP47, and LHClI subunits	Henry et al., 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Albinaki et al., 2012; Walter et al., 2015
cpFtsY	slr2102	At2g45770	40	36	CS, TM	Thylakoid protein targeting: cpSRP translocase	Insertion and assembly of PSII proteins such as D1, D2, and CP47, and LHClI subunits	Henry et al., 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Albinaki et al., 2012; Walter et al., 2015
ALB3	slr1471	At2g28800	50	45	TM	Thylakoid protein targeting: cpSRP translocase	Insertion and assembly of PSII proteins such as D1, D2, and CP47, and LHClI subunits	Pasch et al., 2005; Henry et al., 2007; Ma et al., 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Walter et al., 2015
cpSecA1	sl0616	At4g01800	117	111	CS, TM	Thylakoid protein targeting: cpSec translocase	Insertion and assembly of PSII proteins such as PsbO	Cline and Theg, 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Albinaki et al., 2012
cpSecA2	–	At1g21650	203	203	CS, TM	Thylakoid protein targeting: cpSec translocase	Insertion and assembly of PSII proteins such as PsbO	Cline and Theg, 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Albinaki et al., 2012
cpSecE1	ssl3335	At4g14870	19	15	TM	Thylakoid protein targeting: cpSec translocase	Insertion and assembly of PSII proteins such as PsbO	Cline and Theg, 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Albinaki et al., 2012
cpSecE2	–	At4g38490	17	12	TM	Thylakoid protein targeting: cpSec translocase	Insertion and assembly of PSII proteins such as PsbO	Cline and Theg, 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Albinaki et al., 2012
cpSecY1	sl11814	At2g18710	59	51	TM	Thylakoid protein targeting: cpSec translocase	Insertion and assembly of PSII proteins such as D1 and PsbO	Cline and Theg, 2007; Henry et al., 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Albinaki et al., 2012; Walter et al., 2015
cpSecY2	–	At2g31530	65	61	TM	Thylakoid protein targeting: cpSec translocase	Insertion and assembly of PSII proteins such as D1 and PsbO	Cline and Theg, 2007; Henry et al., 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Albinaki et al., 2012; Walter et al., 2015
Tha4	slr1046	At5g28750	16	14	TM	Thylakoid protein targeting: cpTat translocase	Insertion and assembly of PSII proteins such as PsbP and PsbQ	Cline and Theg, 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Albinaki et al., 2012

(Continued)

TABLE 1 | Continued

Name	Gene locus in <i>Synechocystis</i> sp. PCC 6803 ^a	Gene locus in <i>Arabidopsis</i> ^b	Full-length Size (kDa)	Mature size (kDa)	Location	Protein classification	Function	References ^c
HCF106	sl1046	At5g52440	28	19	TM	Thylakoid protein targeting: cpTat translocase	Insertion and assembly of PSII proteins such as PsbP and PsbQ	Cline and Theg, 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Albriak et al., 2012
cpTatC	sl1094	At2g01110	37	34	TM	Thylakoid protein targeting: cpTat translocase	Insertion and assembly of PSII proteins such as PsbP and PsbQ	Cline and Theg, 2007; Schunemann, 2007; Cline and Dabney-Smith, 2008; Albriak et al., 2012
CPRabA5e	–	At1g05810	29	24	CS, TM	Chloroplast vesicle transport	Transport of PSII proteins such as LHCB1, LHCB3, and CP47, to and from thylakoids	Karim et al., 2014; Karim and Aronsson, 2014
CYO1/SCO2	–	At3g19220	21	17	TM	Chloroplast vesicle transport; thiol/disulfide-modulating protein	Chloroplast and thylakoid biogenesis; folding and transport of cysteine-containing proteins such as CP43, CP47, and LHCB1; stability of PSI-LHCI and PSII-LHCII supercomplexes	Shimada et al., 2007; Albrecht et al., 2008; Muranaka et al., 2012; Tanz et al., 2012
THF1/PSB29	sl1414	At2g20890	34	27	CE, CS, TM	Chloroplast vesicle transport	Thylakoid biogenesis; dynamics of PSII-LHCII supercomplexes	Wang et al., 2004; Keren et al., 2005; Huang et al., 2006, 2013; Shi et al., 2012; Yamatani et al., 2013
TerC	–	At5g12130	42	34	TM	Chloroplast vesicle transport	Thylakoid biogenesis; co-translational insertion of PSII proteins such as D1, D2, and CP43	Kwon and Cho, 2008; Schneider et al., 2014
VIPP1	sl10617	At1g65260	36	32	ICE, TM	Chloroplast vesicle transport	Thylakoid biogenesis; transport and/or co-translational insertion of photosynthetic proteins such as D1	Kroll et al., 2001; Zhang et al., 2012; Zhang and Sakamoto, 2012; Walter et al., 2015
PPL1	sl1418	At3g55330	26	18	TL	PSII subunit-like: PsbP-like	PSII repair	Ishihara et al., 2007
ELIP1	–	At3g22840	20	16	TM	PSII subunit-like: LHCP-like	Binding of chlorophyll and/or stability of pigment-binding proteins and complexes during photoinhibition?	Hutin et al., 2003; Casazza et al., 2005; Hedded et al., 2006; Rossini et al., 2006
ELIP2	–	At4g14690	20	16	TM	PSII subunit-like: LHCP-like	Binding of chlorophyll and/or stability of pigment-binding proteins and complexes during photoinhibition?	Hutin et al., 2003; Casazza et al., 2005; Hedded et al., 2006; Rossini et al., 2006
SEP3.1/LIL3.1	–	At4g17600	29	25	TM	PSII subunit-like: LHCP-like	Anchoring geranylgeranyl reductase to thylakoid membranes; stabilizing LHCII	Tanaka et al., 2010; Takahashi et al., 2014; Lohscheider et al., 2015

(Continued)

TABLE 1 | Continued

Name	Gene locus in <i>Synechocystis</i> sp. PCC 6803 ^a	Gene locus in <i>Arabidopsis</i> ^b	Full-length Size (kDa)	Mature size (kDa)	Location	Protein classification	Function	References ^c
SEP3.2/ LIL3.2	–	At5g47110	29	24	TM	PSII subunit-like: LHCP-like	Anchoring geranylgeranyl reductase to thylakoid membranes; stabilizing LHCII	Tanaka et al., 2010; Takahashi et al., 2014; Lohscheider et al., 2015
HCF173	sl11218	At1g16720	66	57	CS, TM	atypical SDR	Translational initiation of the <i>psbA</i> mRNA	Schult et al., 2007; Link et al., 2012
HCF244	slr0399	At4g35250	44	38	TM	atypical SDR	Translational initiation of the <i>psbA</i> mRNA	Link et al., 2012
CtpA1	slr0008	At3g57680	56	47	TL	C-terminal processing peptidase	C-terminal processing of D1 under high light	Yamamoto et al., 2001; Yin et al., 2008
CtpA2	–	At4g17740	56	46	TL	C-terminal processing peptidase	C-terminal processing of D1	Yamamoto et al., 2001; Che et al., 2013
LPA1/ PrtA	slr2048	At1g02910	50	47	TM	TPR	Biogenesis and assembly of the D1 protein	Peng et al., 2006
MET1	–	At1g55480	37	30	TM	TPR	Supercomplex formation in PSII repair	Ishikawa et al., 2005a; Bhuiyan et al., 2015
LQY1	–	At1g75690	16	12	TL	Thio/disulfide-modulating protein	Disassembly, folding, and/or reassembly of cysteine-containing PSII subunits and complexes and/or D1 synthesis and turnover, during PSII repair	Lu, 2011; Lu et al., 2011
PD16/ PDIL1-2	–	At1g77510	56	54	CS	Thio/disulfide-modulating protein	Regulation of D1 synthesis	Houston et al., 2005; Wittenberg et al., 2014
TRX-M1	slr0623	At1g03680	20	14	CS	Thio/disulfide-modulating protein	Assembly of CP47 into PSII	Cain et al., 2009; Wang et al., 2013
TRX-M2	–	At4g03520	20	13	CS	Thio/disulfide-modulating protein	Assembly of CP47 into PSII	Cain et al., 2009; Wang et al., 2013
TRX-M4	–	At3g15360	21	13	CS	Thio/disulfide-modulating protein	Assembly of CP47 into PSII	Cain et al., 2009; Wang et al., 2013
LTO1	slr0565	At4g35760	40	35	TM	Thio/disulfide-modulating protein	Disulfide bond formation in PsbO	Feng et al., 2011; Karamoko et al., 2011; Lu et al., 2013
RBD1	slr2033	At1g54500	22	16	TM	Thio/disulfide-modulating protein	PSII assembly and stability	Calderon et al., 2013

(Continued)

TABLE 1 | Continued

Name	Gene locus in <i>Synechocystis</i> sp. PCC 6803 ^a	Gene locus in <i>Arabidopsis</i> ^b	Full-length Size (kDa)	Mature size (kDa)	Location	Protein classification	Function	References ^c
CYP20-3/ROCA	slr1251	At3g62030	34	23	CS	PPiase	Repair and reassembly of PSII under high light; redox regulation during stress acclimation	Lippuner et al., 1994; Cai et al., 2008; Dominguez-Solis et al., 2008; Park et al., 2013; Speiser et al., 2015
CYP38/TLP40	slr0408	At3g01480	48	38-40	TL	PPiase	Inhibiting dephosphorylation of PSII subunits during the PSII repair; conversion of PSII core monomers to PSII supercomplexes	Fulgosi et al., 1998; Vener et al., 1999; Rokka et al., 2000; Fu et al., 2007; Sirpiö et al., 2008; Vasudevan et al., 2012
FKBP20-2	slr1761	At3g60370	27	20	TM, TL	PPiase	Formation of PSII-LHCII supercomplexes under normal and high light	Lima et al., 2006
STN7	–	At1g68830	63	59	TM	Protein kinase	Phosphorylation of LHClI; phosphorylation of D1, D2, CP43, and PsbH under low light	Bellaïflore et al., 2005; Bonardi et al., 2005; Tikkanen et al., 2008; Pesaresi et al., 2011; Tikkanen and Aro, 2012
STN8	–	At5g01920	55	50	TM	Protein kinase	Phosphorylation of D1, D2, CP43, and PsbH	Bonardi et al., 2005; Tikkanen et al., 2008; Pesaresi et al., 2011; Tikkanen and Aro, 2012; Nath et al., 2013
PBCP	–	At2g30170	32	30	CS, TM	Protein phosphatase	Dephosphorylation of D1, D2, CP43, and PsbH	Samol et al., 2012
TLP18.3	slr1390	At1g54780	31	18	TL, TM	Protein phosphatase	D1 degradation and PSII dimerization; dephosphorylation of PSII core proteins (e.g., D1 and D2)	Sirpiö et al., 2007; Wu et al., 2011
PPH1/TAP38	–	At4g27800	43	41	TM	Protein phosphatase	Dephosphorylation of LHClI	Pribil et al., 2010; Shapiguzov et al., 2010; Pesaresi et al., 2011
FtsH1	slr1390, slr0228, slr1604, slr1463 ^d	At1g50250	77	71	TM	FtsH protease	Degradation of photodamaged D1	Sakamoto et al., 2003; Yu et al., 2004, 2005; Zaltsman et al., 2005b
FtsH2/VAP2	–	At2g30950	74	69	TM	FtsH protease	Chloroplast biogenesis; thylakoid formation; degradation of photodamaged D1	Bailey et al., 2002; Sakamoto et al., 2002, 2003; Yu et al., 2004, 2005; Zaltsman et al., 2005a,b; Kato et al., 2007, 2009, 2012; Wagner et al., 2011
FtsH5/VAP1	–	At5g42270	75	69	TM	FtsH protease	Chloroplast biogenesis; thylakoid formation; degradation of photodamaged D1	Sakamoto et al., 2002, 2003; Yu et al., 2004, 2005; Zaltsman et al., 2005b; Kato et al., 2009; Wagner et al., 2011

(Continued)

TABLE 1 | Continued

Name	Gene locus in <i>Synechocystis</i> sp. PCC 6803 ^a	Gene locus in <i>Arabidopsis</i> ^b	Full-length Size (kDa)	Mature size (kDa)	Location	Protein classification	Function	References ^c
FtsH6		At5g15250	77	69	TM	FtsH protease	Degradation of LHCII during high-light acclimation and senescence?	Sakamoto et al., 2003; Yu et al., 2004; Zelisko et al., 2005; Wagner et al., 2011
FtsH8		At1g06430	73	69	TM	FtsH protease	Degradation of photodamaged D1	Sakamoto et al., 2003; Yu et al., 2004, 2005; Zaltsman et al., 2005b; Wagner et al., 2011
FtsH11		At5g53170	89	82	C(TM?), IMM	FtsH protease	Thermoprotection of the photosynthetic apparatus	Sakamoto et al., 2003; Yu et al., 2004; Urantowka et al., 2005; Chen et al., 2006b; Wagner et al., 2011
Deg1	slr1204, slr1679, slr1427 ^e	At3g27925	47	42	TL, TM,	Deg protease	Degradation of plastocyanin and PsbO, and photodamaged D1; integration of newly synthesized PSII subunits such as D1, D2, CP43, and CP47, into PSII complexes	Chassin et al., 2002; Huesgen et al., 2005; Kapri-Pardes et al., 2007; Sun et al., 2010b; Schuhmann and Adamska, 2012
Deg2		At2g47940	67	59	CS, TM	Deg protease	Stress-induced degradation of LHCB6; a minor protease in <i>in vivo</i> degradation of photodamaged D1	Haubühl et al., 2001; Huesgen et al., 2005, 2006; Lucinski et al., 2011b; Schuhmann and Adamska, 2012
Deg5		At4g18370	35	32	TL, TM	Deg protease	Degradation of photodamaged D1; wound-induced degradation of PsbF	Huesgen et al., 2005; Sun et al., 2007; Lucinski et al., 2011a; Kato et al., 2012; Schuhmann and Adamska, 2012
Deg7		At3g03380	120	120?	CS, TM	Deg protease	Degradation of photodamaged D1, D2, CP43, and CP47	Huesgen et al., 2005; Sun et al., 2010a; Schuhmann et al., 2011; Schuhmann and Adamska, 2012
Deg8		At5g39830	47	45	TL, TM	Deg protease	Degradation of photodamaged D1	Huesgen et al., 2005; Sun et al., 2007; Kato et al., 2012; Schuhmann and Adamska, 2012
HCF243	–	At3g15095	76	67	TM	Other ^f	Biogenesis, C-terminal processing, and assembly of D1; possibly biogenesis of D2	Zhang et al., 2011
PSB27-H1	slr1645	At1g03600	19	12	TL, TM	Other	C-terminal processing of D1 during PSII repair?	Chen et al., 2006a; Wei et al., 2010; Dietzel et al., 2011; Shi et al., 2012; Mabbitt et al., 2014
PSB27-H2/LPA19		At1g05385	22	15	TL, TM	Other	C-terminal processing during <i>de novo</i> PSII assembly	Wei et al., 2010; Shi et al., 2012; Mabbitt et al., 2014

(Continued)

TABLE 1 | Continued

Name	Gene locus in <i>Synechocystis</i> sp. PCC 6803 ^a	Gene locus in <i>Arabidopsis</i> ^b	Full-length Size (kDa)	Mature size (kDa)	Location	Protein classification	Function	References ^c
HCF136	slr2034	At5g23120	44	38	TL	Other	Assembly of PSII reaction-center complexes such as RC, RC47a, and RC47b	Meurer et al., 1998; Plücker et al., 2002; Mabbitt et al., 2014
PAM68	sl0933	At4g19100	24	20	TM	Other	Conversion of PSII minimal reaction-center complexes into larger PSII assembly intermediates; C-terminal processing of D1	Armbruster et al., 2010
PsbN/PBF1	smr0009	AtCg00700	4.7	4.7	TM	Other	Assembly of PSII minimal reaction-center complexes; regulation of PSII core and antenna protein phosphorylation	Krech et al., 2013; Torabi et al., 2014
PSB28	sl1398	At4g28660	22	14	TL	Other	Biogenesis of chlorophyll-binding proteins such as CP47, PsaA, and PsaB	Jung et al., 2008; Shi et al., 2012; Mabbitt et al., 2014
LPA2	–	At5g51545	20	~20	TM	Other	Synthesis and assembly of CP43	Ma et al., 2007; Cai et al., 2010
LPA3	–	At1g73060	40	34	CS, TM	Other	Synthesis and assembly of CP43	Cai et al., 2010
PSB33	–	At1g71500	32	25	TM	Other	Association of LHClI with PSII	Fristedt et al., 2015
HHL1	–	At1g67700	26	18	TM	Other	Reassembly of PSII core monomers and PSII-LHClI supercomplexes during PSII repair	Jin et al., 2014
MPH1	–	At5g07020	24	20	TM	Other	Assembly and/or stability of PSII core monomers and higher order PSII complexes under high light	Liu and Last, 2015a,b

C, chloroplast; CE, chloroplast envelope; CS, chloroplast stroma; ICE, inner chloroplast envelope; IMM, inner mitochondrial membranes; TL, thylakoid lumen; TM, thylakoid membranes.

^aThe gene loci in *Synechocystis* sp. PCC 6803 are included in this table only to distinguish between factors that are conserved in cyanobacteria and land plants and factors that are found in land plants but not in cyanobacteria. The hyphen indicates that the corresponding factor is either absent or not yet found in *Synechocystis* sp. PCC 6803.

^bThis article focuses on the identification and roles of PSII assembly, stability, and repair factors in *Arabidopsis*; therefore the gene loci for factors in other land plants are not listed in this table.

^cDetailed descriptions of factors involved in the assembly, stability, and repair of PSII in cyanobacteria can be found in Nickelsen and Hengstl (2013). Therefore, references for factors in cyanobacteria are not listed in this table.

^dThe *Synechocystis* sp. PCC 6803 genome contains four *FtsH* genes. Because there is no straightforward one-to-one correspondence between the 12 *Arabidopsis* *FtsH* genes and four cyanobacterial *FtsH* genes, the four cyanobacterial *FtsH* loci are listed together.

^eThe *Synechocystis* sp. PCC 6803 genome contains three *Deg* genes. Because there is no straightforward one-to-one correspondence between the 16 *Arabidopsis* *Deg* genes and three cyanobacterial *Deg* genes, the three cyanobacterial *Deg* loci are listed together.

^fFactors that do not fall in the 12 well-defined classifications (see Section Proteins that influence the assembly, stability, and repair of PSII) are classified as “other proteins” that influence the assembly, stability, and repair of PSII.

requires none of the known targeting apparatus are responsible for translocating thylakoid membrane proteins. LHCPs are translocated via the cpSRP pathway that requires the action of cpSRP43 (chloroplast Signal Recognition Particle protein of 43 kDa), cpSRP54 (cpSRP protein of 54 kDa), cpFtsY (chloroplast filamentation temperature sensitive protein Y), and ALB3 (Albino3) (Figures 1, 2). ALB3 was found to interact with PSII subunits D1, D2, and CP47 (Ossenbühl et al., 2004; Pasch et al., 2005; Göhre et al., 2006), consistent with the function of the cpSRP pathway in translocating PSII proteins such as D1, D2, and CP47 (Figures 1, 2). Most of the remaining thylakoid membrane proteins are inserted by the unusual pathway that requires none of the known targeting apparatus. The cpSecA-cpSecYE (cpSec means chloroplast secretory) pathway and the chloroplast twin-arginine translocation (cpTat) pathway are responsible for translocation of luminal proteins. In Arabidopsis, each cpSec component is encoded by two genes (Table 1). The cpTat pathway has three components: Thylakoid Assembly 4 (THA4), High Chlorophyll Fluorescence 106 (HCF106), and cpTat protein C (cpTatC) (Albiniak et al., 2012). PsbO and plastocyanin are substrates of the cpSec pathway while PsbP and PsbQ are substrates of the cpTat pathway (Figures 1, 2; Albiniak et al., 2012). Like their bacterial counterparts, components of the cpSRP and cpSec pathways may act in a modular fashion (Cline and Theg, 2007; Henry et al., 2007). For instance, cpSRP54 was identified in co-translational D1 insertion intermediates, along with cpSecY (chloroplast secretory translocase Y) and chloroplast ribosomes (Cline and Theg, 2007; Henry et al., 2007). Consistent with this finding, Walter et al. (2015) reported that cpSecY forms a complex with VIPP1 (Vesicle-Inducing Protein in Plastids 1) and cpSec components ALB3, cpFtsY, and cpSRP54 during co-translational integration of D1 (Figures 1, 2). How different thylakoid membrane and luminal proteins, including components of PSII complexes, are transported and integrated into thylakoids by the above-mentioned thylakoid-protein-targeting complexes can be found in a number of reviews (Schunemann, 2007; Cline and Dabney-Smith, 2008; Albiniak et al., 2012).

Components of the Putative Chloroplast-Vesicle-Transport System

PSII is located in the thylakoid membranes of oxygenic photosynthetic organisms; therefore thylakoid membrane biogenesis is essential to PSII assembly (Nickelsen and Zerges, 2013; Rast et al., 2015). It has been suggested that thylakoids could develop from invaginations of the inner envelope membrane via a vesicle-based transfer process and these vesicles are thought as a method of transporting lipids and proteins to and from thylakoids (Hooper et al., 1991; Westphal et al., 2001, 2003; Charuvi et al., 2012; Karim and Aronsson, 2014; Rast et al., 2015). Chloroplast vesicles are typically not observed when vesicular transport from the inner chloroplast envelope to thylakoids is continuous at ambient temperature (Morré et al., 1991). When vesicular transport is blocked by low temperature, chloroplast vesicles accumulate (Morré et al., 1991). A number of proteins have been implicated as part of

the chloroplast-vesicle-transport system, including CPRabA5e (chloroplast Rab GTPase A5e), CYO1/SCO2 (Shiyou 1/Snowy Cotyledon 2), THF1/PSB29 (Thylakoid Formation 1/PSII protein 29), TerC (Tellurite-resistance protein C), and VIPP1 (Karim and Aronsson, 2014; Rast et al., 2015).

CPRabA5e is a small Rab GTPase targeted to the chloroplast stroma and thylakoid membranes (Karim et al., 2014). Transfer DNA (T-DNA) insertions in the *CPRabA5e* gene cause a reduced amount of grana thylakoids in Arabidopsis leaves (Karim et al., 2014). After pre-incubation at 4°C, the *cpRabA5e* mutants have larger plastoglobules and an increased number of small vesicles, compared to the wild type. CPRabA5e was found to interact with a number of photosynthetic proteins, including PSI subunits H2 and K (PsaH2 and PsaK; Psa stands for Photosystem I), the PSII core subunit CP47, and LHC proteins LHCb1 and LHCb3 (Karim et al., 2014). These findings led to the hypothesis that CPRabA5e is involved in transport of photosynthetic proteins, such as LHCb1, LHCb3, and CP47 (Figures 1, 2), to developing thylakoids in young chloroplasts via vesicles (Karim et al., 2014).

CYO1/SCO2 is a zinc-finger-domain-containing thylakoid membrane protein with protein disulfide isomerase (PDI) activity (Shimada et al., 2007; Albrecht et al., 2008; Muranaka et al., 2012; Tanz et al., 2012). A loss-of-function mutation in the *CYO1/SCO2* gene results in globularly or normally shaped plastids with very big vesicles in Arabidopsis cotyledons (Tanz et al., 2012). A closer look at normally shaped chloroplasts from mutant cotyledons showed that small vesicles emerged from the inner chloroplast envelope, even at ambient temperature. These vesicles were not observed in chloroplasts from wild-type cotyledons used in the same study. These observations suggest that vesicular transport from the inner chloroplast envelope to developing thylakoids is blocked in the cotyledons of mutant seedlings and that the chloroplast-vesicle-transport system is important for thylakoid biogenesis rather than damage or programmed degradation of the thylakoid membrane system. CYO1/SCO2 was found to interact with PSI core subunits PsaA and PsaB, PSII core subunits CP43 and CP47, and LHCb1 (Muranaka et al., 2012; Tanz et al., 2012) and to co-migrate with Photosystem I-light-harvesting complex I (PSI-LHCI) and PSII-LHCII supercomplexes in blue native-polyacrylamide gel electrophoresis (BN-PAGE) (Shimada et al., 2007). Therefore, it is conceivable that CYO1/SCO2 participates in transport of photosynthetic proteins, such as CP43, CP47, and LHCb1 (Figures 1, 2), to and from thylakoids via chloroplast vesicles in cotyledons (Tanz et al., 2012). The association of CYO1/SCO2 with PSI-LHCI and PSII-LHCII supercomplexes also begs the question whether this protein has a role in stabilizing these supercomplexes (Figures 1, 2).

THF1/PSB29 is a coiled-coil-domain-containing protein targeted to the chloroplast envelope, chloroplast stroma, and thylakoid membranes (Wang et al., 2004; Huang et al., 2006). The *thf1/psb29* knockout mutant of Arabidopsis has a variegated phenotype in cotyledons and true leaves and the chloroplasts in the yellow sectors of *thf1/psb29* leaves lack normal thylakoids and accumulate chloroplast vesicles (Wang et al., 2004; Keren et al., 2005; Zhang et al., 2009). These data led to the hypothesis that THF1/PSB29 is required for organizing chloroplast vesicles

into mature thylakoids (Wang et al., 2004). THF1/PSB29 was thought to be involved in maintaining levels of FtsHs in plants because D1, a substrate of FtsHs, was found to be more stable in the *thf1/psb29* mutant than in the wild type (Zhang et al., 2009). However, it was later found that the stay-green phenotype of the *thf1/psb29* mutant is not due to reduced FtsH protease activity, because *fsh2/var2* (*var2* stands for *yellow variegated 2*) leaves turn yellow much faster than wild-type and *thf1/psb29* leaves during dark-induced senescence (Huang et al., 2013). THF1/PSB29 was found to interact with all six LHCb proteins (Huang et al., 2013); therefore it is possible that LHCb proteins are transported to thylakoid membranes via direct interaction with THF1/PSB29 (Figures 1, 2). In addition, it was proposed that THF1/PSB29 regulates the dynamics of PSII-LHCII supercomplexes during high-light stress and leaf senescence (Figure 2; Huang et al., 2013; Yamatani et al., 2013). The THF1/PSB29-deficient mutants of Arabidopsis and rice (*Oryza sativa*) also have a stay-green phenotype in pathogen-infected and dark-induced senescent leaves (Huang et al., 2013; Yamatani et al., 2013). In dark- and high-light-treated *thf1/psb29* Arabidopsis leaves, PSII-LHCII supercomplexes are highly unstable but a type of PSII-LHCII megacomplexes is retained (Huang et al., 2013). Consistent with a role in regulating the dynamics of PSII-LHCII supercomplexes, THF1/PSB29 was found to co-migrate with trimeric and monomeric LHCII in BN-PAGE (Huang et al., 2013).

TerC is an integral thylakoid membrane protein with eight transmembrane helices (Kwon and Cho, 2008; Schneider et al., 2014). T-DNA insertions in the *TerC* gene caused a pigment-deficient and seedling-lethal phenotype in Arabidopsis (Kwon and Cho, 2008; Schneider et al., 2014). This is accompanied with a substantial reduction or complete loss of thylakoid membranes and over-accumulation of chloroplast vesicles. Therefore, TerC was considered to be involved in thylakoid biogenesis and vesicle transport (Kwon and Cho, 2008). To further analyze the function of TerC, Schneider et al. (2014) generated an artificial microRNA-based knockdown allele *amiR-TerC* in Arabidopsis and found that the severe phenotype of the T-DNA mutants is likely due to the substantially reduced rates of synthesis and insertion of PSII proteins. In line with these observations, TerC was found to interact with PSII proteins D1, D2, and CP43 as well as PSII assembly factors ALB3, LPA1/PratA (Low PSII Accumulation 1/Processing-associated tetratricopeptide repeat protein A), LPA2 (Low PSII Accumulation 2), and PAM68 (Photosynthesis Affected Mutant 68) (Schneider et al., 2014). Taken together, it is reasonable to propose that TerC is also involved in co-translational insertion of PSII proteins (e.g., D1, D2, and CP43) into thylakoid membranes, in collaboration with other PSII assembly factors (Figures 1, 2; Schneider et al., 2014).

VIPP1 is homologous to the phage shock protein A in *Escherichia coli*, which is induced under various stress environments (Karim and Aronsson, 2014). VIPP1 is targeted to the inner chloroplast envelope and thylakoid membranes (Kroll et al., 2001). The *vipp1* knockdown and knockout mutants of Arabidopsis are pigment-deficient and semi-lethal (i.e., unable to grow photoautotrophically) (Kroll et al., 2001; Zhang et al.,

2012; Zhang and Sakamoto, 2012). Chloroplasts from the *vipp1* knockout and knockdown mutants are defective in thylakoid membrane formation and vesicle budding from inner envelope membranes. In addition, *vipp1* mutant chloroplasts are swollen due to damage in the chloroplast envelope and increases in the osmotic pressure in the chloroplast stroma (Zhang et al., 2012; Zhang and Sakamoto, 2012). VIPP1 was recently found in co-translational D1 insertion intermediates isolated from thylakoid membranes of pea (*Pisum sativum*) leaves, along with cpSecY, ALB3, cpFtsY, and cpSRP54 (Walter et al., 2015). Therefore, VIPP1 was proposed as a multifunctional protein that is involved in chloroplast vesicular transport, thylakoid biogenesis, and co-translational insertion of photosynthetic proteins (Figure 1).

PSII Subunit-Like Proteins, e.g., PsbP-Like and LHCP-Like Proteins

Five PSII subunit-like proteins, including one PsbP-like protein and four LHCP-like proteins, have been implicated to be involved in the assembly, stability, and/or repair of PSII complexes or subunits. PsbP-like proteins are in the same family as PsbP proteins (Bricker et al., 2013; Ifuku, 2014). Unlike PsbP proteins, whose primary function is water splitting and oxygen evolution, PsbP-like proteins are not part of the OEC. Arabidopsis has two PsbP-like proteins: PPL1 and PPL2 (Table 1). A T-DNA insertion in the *PPL1* gene led to increased sensitivity to high light and delayed recovery after photoinhibition (Ishihara et al., 2007). These data suggest that PPL1 is required for efficient repair of photodamaged PSII. Although PPL1 has not been shown to be associated with PSII in higher plants, its cyanobacterial homolog cyanoP (ssl1418) has been shown to be loosely associated with PSII and possess the same beta-sandwich fold and a well-conserved zinc-binding site as PsbP in higher plants (Ishikawa et al., 2005b; Summerfield et al., 2005; Michoux et al., 2010, 2014; Jackson et al., 2012). Further studies are needed to investigate the precise function of PPL1 in PSII repair. Unlike PPL1, PPL2 is required for the accumulation of chloroplast NAD(P)H dehydrogenase complex (Ishihara et al., 2007); therefore the function of PPL2 is not discussed in this review.

LHCP-like proteins are in the same superfamily as LHCPs (Heddad et al., 2012). Unlike LHCPs, whose primary function is light-harvesting, most LHCP-like proteins are involved in chlorophyll- and carotenoid-binding, assembly and stability of chlorophyll-protein complexes, and/or photoprotection. LHCP-like proteins can be classified into three subfamilies: three-helix, early light-induced proteins (ELIPs), two-helix, stress-enhanced proteins (SEPs), and one-helix, high-light-induced, small chlorophyll-binding-like proteins (OHPs/HLIPs/SCPs). ELIPs are restricted to green algae and land plants, SEPs are ubiquitously present in photosynthetic eukaryotes, OHP1 exists in cyanophages, cyanobacteria, and photosynthetic eukaryotes, and OHP2 exists in eukaryotes. The Arabidopsis genome encodes two ELIPs, six SEPs, and two OHPs (Heddad et al., 2012).

ELIPs appear to be associated with PSII under standard conditions in pea plants; they become associated with monomeric and trimeric LHCII under high light in Arabidopsis

(Adamska and Klopstech, 1991; Heddad et al., 2006). Heddad et al. (2006) showed that the relative amounts of ELIP transcripts and proteins increase as the light intensity increases. It was therefore proposed that ELIPs may be involved in photoprotection by binding free chlorophyll released during degradation of pigment-binding proteins or by stabilizing the assembly of pigment-binding proteins during photoinhibition (Adamska and Klopstech, 1991; Hutin et al., 2003). The potential function of ELIPs in stabilizing LHCII and PSII-LHCII supercomplexes under light stress is included in **Figure 2**. However, *elip1* and *elip2* single knockout mutants and the *elip1 elip2* double knockout mutant have a similar phenotype and light sensitivity as the wild-type Arabidopsis plants (Casazza et al., 2005; Rossini et al., 2006). Additional studies are needed to dissect the exact functions of ELIP proteins.

Among the six SEPs in Arabidopsis, the functions of SEP3.1/LIL3:1 (SEP stands for stress-enhanced protein; LIL stands for light-harvesting-like protein) and SEP3.2/LIL3:2 have been extensively studied (Tanaka et al., 2010; Takahashi et al., 2014; Lohscheider et al., 2015; Mork-Jansson et al., 2015). The anti-sense *sep3.1/lil3:1* mutant and the *sep3.1/lil3:1 sep3.2/lil3:2* double knockout mutant are deficient in chlorophyll and α -tocopherol biosynthesis (Tanaka et al., 2010; Lohscheider et al., 2015). The deficiency is due to a substantial reduction in the amount of chlorophyll and α -tocopherol biosynthetic enzyme geranylgeranyl reductase. In line with these findings, SEP3.1/LIL3:1 and SEP3.2/LIL3:2 were found to interact with geranylgeranyl reductase and their transmembrane domain was found to be important for the interaction (Tanaka et al., 2010; Takahashi et al., 2014). Therefore, SEP3.1/LIL3:1 and SEP3.2/LIL3:2 were proposed to be involved in chlorophyll and tocopherol biosynthesis by anchoring and stabilizing geranylgeranyl reductase to thylakoid membranes. In addition, SEP3.1/LIL3:1 and SEP3.2/LIL3:2 were found to accumulate with increasing light irradiance and they are associated with subcomplexes of LHCII (Lohscheider et al., 2015). Thus, it is also possible that SEP3.1/LIL3:1 and SEP3.2/LIL3:2 may function in stabilizing LHCII (**Figures 1, 2**).

The OHP genes have an expression pattern similar to the *ELIP* and *SEP* genes, whose expression is up-regulated upon high light (Mulo et al., 2008; Heddad et al., 2012). The function of OHP1 is not yet known, but OHP2 was showed to be associated with PSI under low or high light and was therefore proposed to play a role in photoprotection of PSI (Andersson et al., 2003). Recently, a cyanobacterial OHP family protein was found to bind chlorophyll *a* and β -carotene and possess an energy-dissipative conformation, suggesting that OHP family proteins may have a photoprotective role (Staleva et al., 2015).

Atypical SDR Family Proteins

Classic SDR family proteins have an intact cofactor-binding site (TGXXGXXG) and an intact catalytic tetrad (NSYK), which are required for their SDR activity (Persson et al., 2009). Unlike classic SDR family proteins, atypical SDR family proteins have no known enzyme activity because they have an altered glycine-rich cofactor-binding site and partially or completely lack the signature catalytic tetrad (Link et al., 2012). Two atypical SDR

family proteins have been found to be important for PSII: HCF173 (High Chlorophyll Fluorescence 173) and HCF244 (High Chlorophyll Fluorescence 244) (Schult et al., 2007; Link et al., 2012; Chidgey et al., 2014; Knoppová et al., 2014). Compared to HCF244, HCF173 is ~200 amino acids longer and its SDR domain is fragmented into two regions. HCF173 and HCF244 have the same subcellular localization: they are both predominantly associated with chloroplast membranes, with a small fraction located in the chloroplast stroma. Loss-of-function mutations in the *HCF173* or *HCF244* gene result in similar defects in Arabidopsis: a drastic reduction in D1 synthesis, inability to accumulate PSII subunits, substantial decreases in PSII activity, and a complete loss of photoautotrophy (Schult et al., 2007; Link et al., 2012). The *hcf173* and *hcf244* single mutants are able to grow on sucrose-supplemented media but they are pale green and much smaller than the wild type (Link et al., 2012). Polysome association experiments demonstrated that these defects are caused by reduced translation initiation of the *psbA* transcript (Schult et al., 2007; Link et al., 2012). The decrease in translation initiation is accompanied by a reduction in *psbA* mRNA stability. The *hcf173 hcf244* double mutant grown on sucrose-supplemented media is smaller than the single mutants, suggesting that simultaneous loss of HCF173 and HCF244 has an additive effect (Link et al., 2012). The function of HCF173 and HCF244 in translation initiation of the *psbA* transcript is included in **Figures 1, 2**. Some SDR family proteins, such as dihydroloipoamide acetyltransferases, glyceraldehyde-3-phosphate dehydrogenase, and lactate dehydrogenase, have evolved the capacity to bind RNA (Hentze, 1994; Nagy et al., 2000; Pioli et al., 2002; Bohne et al., 2013). Therefore, it is possible that HCF173 and HCF244 may act as RNA-binding proteins and facilitate translation initiation of the *psbA* mRNA (Link et al., 2012).

C-terminal Processing Peptidases

The PSII reaction-center protein D1 is often synthesized in the precursor form (pD1), with a C-terminal extension of 8–16 amino acids (Nixon et al., 1992; Anbudurai et al., 1994; Liao et al., 2000). In plants, this C-terminal extension is cleaved in a single step by C-terminal processing peptidase A (CtpA) at an early step of *de novo* PSII assembly or at the reassembly step of PSII repair (**Figures 1, 2**). CtpAs are serine endopeptidases with a serine/lysine catalytic dyad (Anbudurai et al., 1994; Liao et al., 2000; Yamamoto et al., 2001). Recombinant spinach (*Spinacia oleracea*) CtpA exhibited efficient proteolytic activity toward thylakoid membrane-embedded pD1 (Yamamoto et al., 2001). Arabidopsis has three CtpAs in the thylakoid lumen and the functions of CtpA1 and CtpA2 have been studied (Yin et al., 2008; Che et al., 2013).

Under normal growth conditions, T-DNA insertions in the Arabidopsis *CtpA1* gene do not cause changes to plant growth and morphology, PSII activity, or thylakoid membrane complex formation (Yin et al., 2008). Under high light, the *ctpA1* mutant displays retarded growth, accelerated D1 turnover, as well as increased photosensitivity and delayed recovery of PSII activity. Therefore, CtpA1 was proposed to be involved in D1 protein C-terminal processing in the PSII repair cycle (**Figure 2**).

Unlike the *ctpA1* mutant, the T-DNA mutant of the Arabidopsis *CtpA2* gene is lethal under normal light but is viable in sucrose-supplemented media under low light (Che et al., 2013). The viable *ctpA2* mutant displays a complete loss of the mature D1 protein, reduced levels of other PSII core proteins, a severely decreased level of PSII supercomplexes, and a substantial reduction or complete loss of PSII activity. pD1 and other PSII subunits in the viable *ctpA2* mutant are present in PSII monomers and PSII dimers but absent in PSII supercomplexes. These data suggest that CtpA2 is indispensable for C-terminal processing of D1 (Figure 1), which in itself is essential for *de novo* PSII assembly. A weak allele expressing ~2% of the wild-type level of CtpA2 appears to be normal under normal light but displays stunted growth and over-accumulation of pD1 under elevated light (Che et al., 2013). These data suggest that CtpA2 is also involved in C-terminal processing of D1 during high-light-induced PSII repair (Figure 2).

TPR Proteins

The TPR is a 34-amino acid repeated motif that ubiquitously exists among all organisms (Ishikawa et al., 2005a). Two TPR proteins have been found to be involved in PSII assembly and/or repair: LPA1/PratA and MET1 (Mesophyll-Enriched Thylakoid protein 1) (Ishikawa et al., 2005a; Peng et al., 2006; Bhuiyan et al., 2015).

LPA1/PratA is an intrinsic thylakoid membrane protein with two tandem TPR motifs and a double-pass transmembrane domain (Klinkert et al., 2004; Peng et al., 2006). T-DNA insertions in the Arabidopsis *LPA1/PratA* gene result in reduced growth, pale-green leaves, reduced PSII activity, reduced amounts of PSII proteins, reduced synthesis of D1 and D2, increased turnover of PSII core subunits D1, D2, CP43, and CP47, and inefficient assembly of PSII (Peng et al., 2006). The transcript levels of genes encoding PSII core subunits are unchanged in the mutants. LPA1/PratA was found to directly interact with D1 in a split-ubiquitin yeast-two-hybrid assay. These data suggest that LPA1/PratA has a role in biogenesis and assembly of D1 (Figures 1, 2). Consistent with this hypothesis, LPA1/PratA was identified in thylakoid-membrane-associated ribosome nascent chain fractions (Peng et al., 2006).

MET1 has an N-terminal PDZ domain and a C-terminal TPR motif, which are conserved across green algae and land plants (Ishikawa et al., 2005a; Bhuiyan et al., 2015). MET1 is peripherally attached to thylakoid membranes on the stromal side and it is enriched in stroma lamellae (Bhuiyan et al., 2015). T-DNA insertions in the Arabidopsis *MET1* gene do not cause obvious changes to the accumulation and assembly state of the photosynthetic apparatus under normal light (Bhuiyan et al., 2015). Under fluctuating light, the *met1* mutants demonstrate reduced growth, decreased PSII efficiency, a near-complete loss of PSII-LHCII supercomplexes, and increased amounts of unassembled CP43. Loss of MET1 also causes increased photosensitivity of PSII activity and an accelerated rate of D1 turnover under high light. MET1 was found to co-migrate with a series of PSII subcomplexes, such as PSII dimers, PSII

core monomers, CP43-less PSII monomers, and PSII reaction-center complexes (i.e., RC, RC47a, and RC47b), in BN-PAGE. Therefore, MET1 was proposed to be involved in supercomplex formation during PSII repair (Figure 2). In line with this hypothesis, MET1 was found to interact with the stromal loops of PSII core subunits CP43 and CP47 (Bhuiyan et al., 2015).

Thiol/Disulfide-Modulating Proteins

Thiol/disulfide modulation is important for regulating photosynthetic processes (Järvi et al., 2013; Karamoko et al., 2013). Three types of thiol/disulfide-modulating proteins have been found to be involved in the assembly, stability, function, and repair of PSII: protein disulfide isomerases (PDIases), protein disulfide reducing proteins, and protein thiol oxidizing proteins.

As mentioned above, thylakoid membrane protein CYO1/SCO2 was found to play a role in chloroplast and thylakoid biogenesis and vesicular transport of photosynthetic proteins to developing thylakoids in cotyledons (Shimada et al., 2007; Albrecht et al., 2008; Tanz et al., 2012). CYO1/SCO2 has a C₄-type zinc-finger domain with two conserved CXXCXGXG repeats, the signature domain for PDIase activity (Shimada et al., 2007). Recombinant CYO1/SCO2 is able to catalyze reduction of protein disulfide bonds and oxidative renaturation of reduced and denatured protein substrates, indicating that CYO1/SCO2 is a PDIase (Shimada et al., 2007). CYO1/SCO2 was found to interact with PsaA, PsaB, CP47, CP43, and LHCBI (Muranaka et al., 2012; Tanz et al., 2012). These five CYO1/SCO2-interacting proteins contain cysteine in the hydrophobic region(s). Therefore, CYO1/SCO2 may also participate in folding of cysteine-containing PSI and PSII subunits (Figures 1, 2), by forming transient disulfide bonds with its protein substrates via the CXXC motif (Jessop et al., 2007; Feige and Hendershot, 2011).

LQY1 (Low Quantum Yield of PSII 1) is another thylakoid membrane protein with PDIase activity. Full-length LQY1 has a chloroplast transit peptide, a transmembrane domain, and a C-terminal C₄-type zinc-finger domain with four conserved CXXCXGXG repeats (Lu et al., 2011). Recombinant LQY1 is able to catalyze oxidative renaturation of reduced and denatured protein substrates and reductive renaturation of oxidized protein substrates. T-DNA insertions in the Arabidopsis *LQY1* gene cause reduced efficiency of PSII photochemistry, increased sensitivity to high light, and increased accumulation of reactive oxygen species under high light. The *lqy1* mutants were found to accumulate fewer PSII-LHCII supercomplexes and have altered rates of high-light-induced D1 turnover and re-synthesis. The *lqy1* mutant phenotype can be suppressed by complementation of *lqy1* mutants with the wild-type *LQY1* gene (Lu, 2011). LQY1 is associated with the PSII core monomer and the CP43-less PSII monomer (a marker for ongoing PSII repair and reassembly) and it is most abundant in stroma-exposed thylakoid membranes, where important steps of PSII repair occurs (Lu et al., 2011). Under high light, LQY1 associated with PSII monomers increases at the expense of free LQY1 and LQY1 associated with smaller PSII complexes. Immunoprecipitation analysis showed that LQY1 interacts with PSII core subunits

CP47 and CP43, which contain three and four conserved cysteine residuals, respectively. Therefore, it was proposed that LQY1 is involved in PSII repair (Lu, 2011; Lu et al., 2011). It is possible that LQY1 participates in disassembly, folding, and/or reassembly of cysteine-containing PSII subunits and complexes and/or regulates D1 synthesis and turnover during PSII repair (Figure 2). These hypotheses require further investigation.

PDI6/PDIL1-2 (Protein Disulfide Isomerase 6/Protein Disulfide Isomerase-Like 1-2) contains two redox-active thioredoxin domains (with the WCGHC active site), two redox-inactive thioredoxin-like domains, and a C-terminal endoplasmic reticulum retention signal KDEL (Houston et al., 2005; Wittenberg et al., 2014). PDI6/PDIL1-2 is dual-targeted to chloroplasts and the endoplasmic reticulum; chloroplast-targeted PDI6 is located in the stroma (Wittenberg et al., 2014). Similar to CYO1/SCO2 and LQY1, recombinant PDI6/PDIL1-2 is capable of catalyzing oxidative renaturation of reduced and denatured protein substrates. Compared to wild-type *Arabidopsis*, the *pdi6-1* and *pdi6-2* knockdown mutants display increased resistance to high light, reduced photoinhibition, and an accelerated rate of D1 synthesis (Wittenberg et al., 2014). Therefore, it was proposed that PDI6/PDIL1-2 may function as an attenuator of D1 synthesis during PSII repair (Figure 2).

Thioredoxins are small proteins that contain a redox-active thioredoxin domain with the WCGHC active site (Cain et al., 2009). As enzymes, thioredoxins are active in the reduced form and are able to reduce disulfide bonds in protein substrates (Cain et al., 2009). Thioredoxins are important for regulating thiol/disulfide homeostasis inside chloroplasts (Cain et al., 2009). Three M-type thioredoxins (TRX-M1, M2, and M4) have been found to be involved in PSII biogenesis in chloroplasts (Wang et al., 2013). TRX-M1, M2, and M4 are associated with minor PSII assembly intermediate subcomplexes and they interact with PSII core subunits D1, D2, and CP47. Simultaneous inactivation of the three *Arabidopsis* TRX-M genes causes pale-green leaves, reduced PSII activity, decreased accumulation of PSII complexes, and increased accumulation of reactive oxygen species. PSII core proteins D1 and CP47 were found to be able to form redox-sensitive intermolecular disulfide bonds and concurrent loss of the three M-type thioredoxins interrupts the redox status of these PSII core subunits. According to these results, Wang et al. (2013) proposed that the three TRX-M proteins may assist incorporation of CP47 into PSII core complexes (Figure 1).

In addition to PDIsases and thioredoxins, a new type of proteins has been found to regulate thiol/disulfide homeostasis and they mainly act as oxidases by converting free thiols on protein substrates into disulfide bridges. One example is Lumen Thiol Oxidoreductase 1 (LTO1), a thylakoid membrane protein with an integral-membrane vitamin K epoxide reductase domain and a soluble disulfide-bond A oxidoreductase-like domain (Feng et al., 2011; Karamoko et al., 2011; Lu et al., 2013). Each of the two domains contains four conserved cysteine residues (a pair of cysteine residues in the CXXC motif and another pair of separate cysteine residues), which are critical for the disulfide-bond-forming activity of LTO1 (Feng et al., 2011). According to membrane topology analysis, Feng et al. (2011) proposed that the eight conserved cysteine residues are

positioned on the luminal side of thylakoid membranes. This led to the hypothesis that LTO1 is involved in formation of the intramolecular disulfide bond in PsbO (Figures 1, 2), which is located on the luminal side of thylakoid membranes (Karamoko et al., 2011). Consistent with this hypothesis, LTO1 was found to interact with PsbO1 and PsbO2 and catalyze formation of intramolecular disulfide bonds in recombinant PsbO (Karamoko et al., 2011). In line with these observations, the amounts of PsbO, PsbQ, and PsbQ are substantially reduced in the LTO1-deficient *Arabidopsis* mutants and the mutants display reduced efficiency of PSII photochemistry, increased accumulation of reactive oxygen species, a smaller plant size, and delayed growth (Karamoko et al., 2011; Lu et al., 2013).

RBD1 (rubredoxin 1) is a small iron-containing protein with a C-terminal transmembrane domain and a rubredoxin domain with two redox-active CXXC motifs (Calderon et al., 2013). Homologs of RBD1 have been found in thylakoid membranes but not plasma membranes of cyanobacteria and in thylakoid membranes of green algae and land plants (Shen et al., 2002; Calderon et al., 2013). The *rbd1* knockout mutants in the cyanobacterium *Synechocystis* sp. PCC 6803, the green alga *Chlamydomonas reinhardtii*, and the higher plant *Arabidopsis* display a substantial reduction or complete loss of PSII activity and photoautotrophy (Calderon et al., 2013). The amounts of PSII core subunits, such as D1, D2, and CP47, are reduced by 40–90% in these mutants while other components of the photosynthetic apparatus, such as PSI, cytochrome *b₆f* complex, and ATP synthase, are not affected. Based on these data, Calderon et al. (2013) proposed that RBD1 is required for the assembly and/or stability of PSII in oxygenic photosynthetic organisms. Further studies are needed to dissect the precise function of RBD1.

PPIases

Peptide bonds to proline have *cis* and *trans* conformations (Fischer et al., 1998; Ingelsson et al., 2009). Therefore, folding of proteins such as PSII subunits often involves *cis-trans* proline isomerization, which is catalyzed by PPIases (He et al., 2004; Romano et al., 2004; Ingelsson et al., 2009). Three PPIase families have been established according to their immunosuppressant ligand specificity: cyclophilins (CYPs), FK506 (tacrolimus)-binding proteins (FKBPs), and parvulins (Fischer et al., 1998; He et al., 2004; Ingelsson et al., 2009). Two CYPs and one FKBP have been found to be important for the assembly, stability, and/or repair of PSII (Järvi et al., 2015).

CYP20-3/ROC4 (20-kDa cyclophilin 3/rotamase cyclophilin 4) is localized in the chloroplast stroma (Lippuner et al., 1994). The *Arabidopsis* T-DNA insertion mutant of CYP20-3/ROC4 has a normal phenotype and normal PSII function under ambient light (Cai et al., 2008). Under high light, PSII in the *cyp20-3/roc4* mutant exhibits increased photosensitivity and delayed recovery, which caused growth retardation and leaf yellowing. Under high light, D1 degradation is not affected in the mutant but repair and reassembly of photodamaged PSII is impaired. According to the *cyp20-3/roc4* mutant phenotype, Cai et al. (2008) proposed that CYP20-3/ROC4 is involved in repair and reassembly of PSII under high light (Figure 2). The

PPIase activity of CYP20-3/ROC4 makes it a good candidate for catalyzing correct folding of PSII proteins during the repair process of PSII. CYP20-3/ROC4 was also reported to link light and redox signals to cysteine biosynthesis and stress acclimation (Dominguez-Solis et al., 2008; Park et al., 2013; Speiser et al., 2015).

CYP38/TLP40 (cyclophilin of 38 kDa/Thylakoid Lumen Protein of 40 kDa) is predominantly confined in the lumen of non-appressed thylakoids (Fulgosi et al., 1998). Full-length CYP38/TLP40 has a bipartite thylakoid lumen targeting transit peptide, a leucine zipper, a phosphatase-binding module, an acid region for protein-protein interaction, and a C-terminal cyclophilin-type PPIase domain (Fulgosi et al., 1998; Sirpiö et al., 2008). CYP38/TLP40 isolated from spinach leaves demonstrates *in vivo* PPIase activity and co-purification with a thylakoid membrane phosphatase (Fulgosi et al., 1998; Vener et al., 1999). In addition, it was reported that CYP38/TLP40 could be released from thylakoid membranes to the thylakoid lumen upon heat stress and the release is associated with activation of dephosphorylation of PSII subunits in thylakoid membranes (Rokka et al., 2000). Therefore, it was proposed that CYP38/TLP40 acts as a phosphatase inhibitor and regulates dephosphorylation of PSII subunits during PSII repair (Figure 2). Consistent with this hypothesis, T-DNA insertions in the Arabidopsis *CYP38/TLP40* gene cause increased phosphorylation of PSII core subunits and increased photosensitivity of PSII activity (Fu et al., 2007; Sirpiö et al., 2008). In addition, CYP38/TLP40 was found to interact with CP47 through its cyclophilin domain (Vasudevan et al., 2012). Dephosphorylation of PSII subunits during light acclimation is carried out by a type 2C protein phosphatase called PSII core phosphatase (PBCP) (Samol et al., 2012). Further studies are needed to investigate whether CYP38/TLP40 interacts with PBCP and inhibits its phosphatase activity. The *cyp38/tlp40* mutants display retarded growth, pale-green leaves, increased accumulation of PSII monomers, and decreased accumulation of PSII supercomplexes, even under low or ambient light (Fu et al., 2007; Sirpiö et al., 2008). In addition, CYP38/TLP40 was found to co-migrate with PSII core monomers in BN-PAGE (Sirpiö et al., 2008). Therefore, it is likely that CYP38 also functions in conversion of PSII core monomers into higher order PSII complexes (Figures 1, 2).

FKBP20-2 (20-kDa FK506-binding protein 2) is located in the thylakoid lumen (Lima et al., 2006). Recombinant FKBP20-2 demonstrates PPIase activity and the C-terminus of FKBP20-2 has a unique pair of cysteine residues which can be reduced by thioredoxin. T-DNA insertions in the Arabidopsis *FKBP20-2* gene cause smaller plant sizes, reduced chlorophyll contents, stunted growth, reduced PSII activity, increased accumulation of PSII monomers and PSII dimers, and decreased accumulation of PSII supercomplexes under normal light conditions (Lima et al., 2006). Under higher light, the difference in PSII activity between the mutant and the wild type is more pronounced. According to the mutant phenotype, Lima et al. (2006) proposed that FKBP20-2 functions in formation of PSII-LHCII supercomplexes under normal and high light (Figures 1, 2).

Protein Kinases

Multiple studies have shown that high light induces phosphorylation of PSII core proteins, such as D1, D2, CP43, and PsbH (Rintamaki et al., 1997; Vener et al., 2001), which facilitates migration of photodamaged PSII complexes from grana stacks to stroma lamellae (Tikkanen et al., 2008; Goral et al., 2010). Two serine/threonine protein kinases were found to be localized the thylakoid membranes: STN7 (state transition 7) and STN8 (state transition 8) (Bellafiore et al., 2005; Bonardi et al., 2005; Nath et al., 2013). Light-induced phosphorylation of PSII core proteins is carried out by STN8, and to a lesser degree under low light also by STN7 (Figure 2; Bonardi et al., 2005; Tikkanen et al., 2008; Nath et al., 2013). The primary role of STN7 is phosphorylation of LHCII proteins, which leads to displacement of LHCII from PSII to PSI (Bellafiore et al., 2005). Phosphorylation of PSII core proteins promotes unfolding of grana stacks and migration of photodamaged PSII complexes from grana stacks to stroma-exposed thylakoids. This allows easier access of membrane or membrane-associated proteases and co-translational integration of D1 and therefore facilitates repair of photodamaged PSII complexes and proteins (Bonardi et al., 2005; Tikkanen et al., 2008; Khatoon et al., 2009; Goral et al., 2010; Herbstová et al., 2012; Tikkanen and Aro, 2012).

Protein Phosphatases

While the migration of photodamaged PSII complexes is assisted by phosphorylation of PSII core proteins, dephosphorylation of D1 is necessary for efficient turnover of photodamaged D1 in stroma lamellae (Järvi et al., 2015). Two chloroplast protein phosphatases, PBCP and TLP18.3 (Thylakoid Lumen Protein of 18.3 kDa), have demonstrated *in vivo* or *in vitro* phosphatase activity toward PSII core proteins (Sirpiö et al., 2007; Wu et al., 2011; Samol et al., 2012). PBCP is a type 2C protein phosphatase predominantly found in the chloroplast stroma, with a minor fraction associated with thylakoid membranes (Samol et al., 2012). Compared to wild-type Arabidopsis, the PBCP-deficient mutants display delayed dephosphorylation of PSII core proteins (D1, D2, CP43, and PsbH) and normal dephosphorylation of LHCII proteins, upon exposure to far-red light, which favors PSI excitation and dephosphorylation of thylakoid proteins (Samol et al., 2012). Samol et al. (2012) concluded that PBCP is required for efficient dephosphorylation of PSII core proteins (Figure 2).

TLP18.3 is a thylakoid membrane protein with the N-terminal domain of unknown function located in the thylakoid lumen (Sirpiö et al., 2007). It was originally identified as an auxiliary protein involved in dimerization of PSII monomers and degradation of photodamaged D1 (Figures 1, 2). The TLP18.3-deficient Arabidopsis mutants do not show a clear visual phenotype under normal growth conditions but exhibit retarded growth under fluctuating light (Sirpiö et al., 2007), suggesting that TLP18.3 is more important to PSII repair than to *de novo* PSII assembly. Compared to the wild type, the TLP18.3-deficient mutants have fewer PSII dimers and more PSI monomers under normal and fluctuating light. In addition, the rate of high-light-induced D1 turnover is ~50% slower in

the TLP18.3-deficient mutants. Consistent with its dual roles in dimerization of PSII monomers, which occurs in grana stacks, and degradation of photodamaged D1, which takes place in stroma lamellae, TLP18.3 was found evenly distributed between grana stacks and stroma lamellae (Sirpiö et al., 2007). It was later found that the domain of unknown function in TLP18.3 possesses acid phosphatase activity toward synthetic phosphorylated oligopeptides that resemble the phosphorylation sites of PSII core proteins D1 and D2 (Wu et al., 2011). However, how the acid phosphatase activity of TLP18.3 is related to the role of TLP18.3 in PSII assembly and repair is still not clear (**Figure 2**): the phosphorylation sites of D1 are exposed to the stroma side while the acid phosphatase domain of TLP18.3 is located in the thylakoid lumen.

Dephosphorylation of LHCII proteins, such as LHCB1 and LHCB2, is carried out by PPH1/TAP38 (Protein Phosphatase 1/Thylakoid-Associated Phosphatase of 38 kDa) (Pribil et al., 2010; Shapiguzov et al., 2010). PPH1/TAP38 is a type 2C protein phosphatase with a C-terminal single-pass transmembrane domain; it is predominantly located in stroma lamellae and grana margins, where active dephosphorylation of LHCII and PSII core proteins occurs. Loss-of-function mutations in the Arabidopsis *PPH1/TAP38* gene causes decreased dephosphorylation of LHCII while overexpression of PPH1/TAP38 enhances dephosphorylation of LHCII (Pribil et al., 2010; Shapiguzov et al., 2010). The phosphorylation status of PSII core proteins is largely unaffected in the *pph1/tap38* mutants, suggesting that the primary function of PPH1/TAP38 is dephosphorylation of LHCII proteins. Recombinant PPH1/TAP38 is able to dephosphorylate LHCII directly, in an *in vitro* assay (Pribil et al., 2010). Reversible phosphorylation of LHCII is important for the movement of LHCII between PSII and PSI, according to the changes in the spectral composition of incident light. Therefore, the catalytic activities of LHCII kinase STN7 and phosphatase PPH1/TAP38 are important for balancing the light absorption capacity between PSI and PSII (Pesaresi et al., 2011).

FtsH Proteases

FtsH proteases are ubiquitous ATP-dependent, zinc metalloendopeptidases (Yu et al., 2004). FtsHs typically consist of an N-terminal double-pass transmembrane domain, an ATPase domain, and a C-terminal zinc-binding site. Crystal structures of the ATPase domain of bacterial FtsHs and single-particle electron cryo-microscopy analysis of cyanobacterial FtsHs showed that FtsHs exist as ringlike hexamers (Krzywdka et al., 2002; Niwa et al., 2002; Boehm et al., 2012b). Bacteria contain one *FtsH* gene and the FtsH protein forms homohexamers while cyanobacteria and eukaryotes have multiple *FtsH* genes and the FtsH proteins form heterohexamers (Mann et al., 2000; Krzywdka et al., 2002; Niwa et al., 2002; Zaltsman et al., 2005b; Boehm et al., 2012b). The Arabidopsis genome encodes 12 FtsH proteases; eight FtsHs (FtsH1, FtsH2/VAR2, FtsH5/VAR1—FtsH9, and FtsH12; VAR1 stands for Yellow Variegated 1) were verified experimentally to be chloroplast-targeted; FtsH11 was showed to be dual targeted to the chloroplast (possible thylakoid membranes) and the inner mitochondria membrane (Chen et al., 2000; Takechi et al., 2000; Sakamoto et al., 2002, 2003; Urantowka et al., 2005).

FtsH2/VAR2 and FtsH5/VAR1 were found to be localized to thylakoid membranes, with their catalytic domain facing the stromal side of the membrane (Chen et al., 2000; Sakamoto et al., 2003). Among the nine chloroplast- or dual-targeted FtsHs, the functions of FtsH1, FtsH2/VAR2, FtsH5/VAR1, FtsH6, FtsH8, and FtsH11 have been explored experimentally. FtsH and Deg proteases have been known to be involved in degradation of photodamaged D1. Early *in vitro* studies suggested that this is a two-step process including the initial cleavage at the stromal DE loop via Deg2 and the subsequent removal of the N-terminal fragment by FtsHs (Lindahl et al., 1996, 2000; Spetea et al., 1999; Haubühl et al., 2001). It was later proposed that FtsHs play a more important role than Deg proteases in D1 turnover (Silva et al., 2003; Nixon et al., 2005, 2010; Huesgen et al., 2009; Kato et al., 2012; Komenda et al., 2012a).

Loss-of-function mutations in the Arabidopsis *FtsH2/VAR2* or *FtsH5/VAR1* gene cause variegated leaves (Sakamoto et al., 2002, 2003; Yu et al., 2004, 2005; Zaltsman et al., 2005a,b; Kato et al., 2007, 2009, 2012; Wagner et al., 2011). The green sectors in the *ftsh2/var2* or *ftsh5/var1* mutants are formed by cells with normal chloroplasts and the white leaf sectors are formed by viable cells with undifferentiated plastids (Chen et al., 2000; Sakamoto et al., 2002; Kapri-Pardes et al., 2007). These data suggest that FtsH2/VAR2 and FtsH5/VAR1 are required for chloroplast biogenesis and thylakoid formation (Chen et al., 2000; Sakamoto et al., 2002; Zaltsman et al., 2005a,b; Kapri-Pardes et al., 2007).

Compared to wild-type leaves, the green leaf sectors of the *ftsh2/var2* or *ftsh5/var1* mutants demonstrate increased photosensitivity and delayed recovery of PSII activity (Zaltsman et al., 2005a,b). Chloroplasts in the green sectors of the *ftsh2/var2* or *ftsh5/var1* mutants accumulate fewer PSII supercomplexes, more PSII subcomplexes, and more reactive oxygen species than wild-type chloroplasts (Kato et al., 2009). These observations are due to the proteolytic activity of FtsHs toward photodamaged D1 (Bailey et al., 2002; Kato et al., 2009). Because the variegated phenotype complicates biochemical analyses, Kato et al. (2009) used another mutation, *fu-gaeril* (*fug1*), to suppress leaf variegation, and generated the non-variegated *ftsh2/var2 fug1* and *ftsh5/var1 fug1* plants. Compared to the *fug1* single mutant, photodamaged D1 is not replaced in the *ftsh2/var2 fug1* and *ftsh5/var1 fug1* mutants, under different light intensities. Taken together, these data show that FtsH2/VAR2 and FtsH5/VAR1 play an important role at the early stage of D1 turnover and not just in the subsequent removal of D1 degradation products (**Figure 2**; Bailey et al., 2002; Kato et al., 2009).

The phenotypes observed in the *ftsh2/var2* or *ftsh5/var1* mutants are absent in the *ftsh1*, *ftsh6*, *ftsh8*, and *ftsh11* single mutants under normal or high-light conditions (Sakamoto et al., 2003; Zaltsman et al., 2005b; Chen et al., 2006b; Wagner et al., 2011). These findings suggest that FtsH2/VAR2 and FtsH5/VAR1 play a dominant role in chloroplast biogenesis, thylakoid formation, and PSII repair (Sakamoto et al., 2003). Phylogenetic analysis showed that *FtsH1* and *FtsH5* are two duplicated genes (subunit type A), so are *FtsH2* and *FtsH8* (subunit type B) (Yu et al., 2004, 2005; Zaltsman et al., 2005b).

The phenotype of the *ftsh2/var2* mutant can be restored by overexpression of *FtsH8*; the *ftsh2/var2 ftsh8* double mutant is infertile; and the *ftsh1 ftsh5/var1* double mutant resembles the *ftsh2/var2 ftsh8* double mutant (Yu et al., 2004; Zaltsman et al., 2005b). These data suggest that FtsH1 and FtsH5/VAR1 are interchangeable, so are FtsH2/VAR2 and FtsH8, and that the presence of two types of FtsH subunits is necessary for chloroplast biogenesis, thylakoid formation, and PSII repair (**Figure 2**; Zaltsman et al., 2005b).

FtsH6 was reported to participate in degradation of LHCII in Arabidopsis leaves during high-light acclimation and senescence (Zelisko et al., 2005). Using an *in vitro* degradation system (i.e., isolated thylakoid membranes), Zelisko et al. (2005) showed that, compared to the wild type, the *ftsh6* knockout mutant has reduced degradation of LHCB1 after high-light acclimation and reduced degradation of LHCB3 after dark-induced senescence. However, *in vivo* degradation of LHCII proteins does not appear to be impaired in the *ftsh6* knockout mutants (Wagner et al., 2011). Under various conditions, including high-light acclimation and dark-induced senescence, the abundances of LHCB1 and LHCB3 in the *ftsh6* knockout mutants are not statistically different from those in the wild type. Further investigation is needed to understand the precise role of FtsH6.

FtsH11 was reported to be critical in thermoprotection of the photosynthetic apparatus (Chen et al., 2006b; Wagner et al., 2011). When exposed to temperatures above 30°C, which are permissive for wild-type Arabidopsis, the growth and development of the *ftsh11* mutants is arrested (Chen et al., 2006b). Compared to the wild type under the same high-temperature treatment, the *ftsh11* mutants have reduced levels of chlorophyll and reduced PSII activity. Consistent with the hypothesis that FtsH11 is involved in thermotolerance, the expression of the *FtsH11* gene is up-regulated by high temperature (Chen et al., 2006b).

Deg Proteases

Deg proteases are ubiquitous ATP-independent, serine endopeptidases (Schuhmann and Adamska, 2012). The Arabidopsis genome encodes 16 Deg proteases, five of which are peripherally attached to thylakoid membranes: two (Deg2 and Deg7) on the stroma side and three (Deg1, Deg5, and Deg8) on the lumenal side (Huesgen et al., 2005; Schuhmann and Adamska, 2012). These five chloroplast-localized Deg proteases have been proposed to be involved in degradation of photodamaged D1 (Schuhmann and Adamska, 2012). In addition to the trypsin-like protease domain, most Deg proteases, such as Deg1, Deg2, Deg7, and Deg8, have at least one PDZ domain for protein-protein interactions. It is conceivable that these chloroplast-localized PDZ domain-containing Deg proteases may act as chaperones and function in assembly of the photosynthetic apparatus (Sun et al., 2010b; Schuhmann and Adamska, 2012).

Arabidopsis RNA interference (RNAi) lines of Deg1 have a smaller plant size, increased sensitivity of PSII activity to high light, and increased accumulation of non-degraded (and

presumably photodamaged) D1 (Kapri-Pardes et al., 2007). The RNAi lines display decreased accumulation of the 16- and 5.2-kDa C-terminal degradation products of D1, which correspond to the cleavage products at the lumenal CD loop, and immediately after the transmembrane helix E, respectively. The addition of recombinant Deg1 into inside-out thylakoid membranes isolated from the Deg1-deficient plants induces formation of the 5.2-kDa C-terminal degradation product of D1. These data suggest that Deg1 is involved in degradation of photodamaged D1 (specifically, the cleavage at the lumenal CD loop immediately downstream of the transmembrane helix E) in PSII repair (**Figure 2**; Kapri-Pardes et al., 2007). Unfortunately, D1 turnover was not assessed in Deg1 RNAi lines. Deg1 is capable of degrading lumenal proteins plastocyanin and PsbO, suggesting that Deg1 may also acts as a general purpose endopeptidase in the thylakoid lumen (Chassin et al., 2002).

Because Deg1 has a PDZ domain for protein-protein interactions, Sun et al. (2010b) investigated whether Deg1 functions as a chaperone during assembly of PSII complexes (Sun et al., 2010b). Deg1 was found to co-migrate with D1 in BN-PAGE and pull down D1, D2, CP43, and CP47 in an immunoprecipitation assay. Recombinant Deg1 has the ability to fold reduced and denatured protein substrates in the presence of both reduced and oxidized glutathione. The RNAi lines of Deg1 display reduced accumulation of PSII complexes and normal accumulation of other photosynthetic complexes. In addition, assembly of newly synthesized PSII subunits into PSII dimers and PSII supercomplexes is hindered in the RNAi lines, although synthesis of the corresponding proteins in chloroplasts is not impaired. Based on these results, Sun et al. (2010b) proposed that Deg1 also acts as a chaperone and functions in the integration of newly synthesized PSII subunits, such as D1, D2, CP43, and CP47, into PSII complexes (**Figures 1, 2**). However, these experimental data do not rule out the possibility that Deg1 is directly involved in degradation of these PSII subunits.

Deg2 has a PDZ domain and a short hydrophobic segment, in addition to the trypsin-like protease domain (Haubühl et al., 2001). Recombinant Deg2 demonstrates proteolytic activity toward photodamaged D1 and is able to produce the 23-kDa N-terminal and the 10-kDa C-terminal degradation products, which correspond to the cleavage products at the stromal DE loop (Haubühl et al., 2001). Although recombinant Deg2 is proteolytically active, the *deg2* knockout mutants have the same plant morphology, PSII activity, and D1 turnover rate as wild-type Arabidopsis, under normal or elevated light (Huesgen et al., 2006). Therefore, it was proposed that Deg2 functions as a minor protease in *in vivo* degradation of photodamaged D1 (**Figure 2**). Deg2 was also reported to be involved in stress-induced degradation of LHCB6 (Luciński et al., 2011b).

Deg5 and Deg8 form heterohexamers and the Deg5-to-Deg8 ratio is ~1:1 (Sun et al., 2007). Deg5 is ~120 amino acids shorter than Deg1 and Deg8 and it does not contain any PDZ domain (Schuhmann and Adamska, 2012). Although Deg8 has a PDZ domain, it shows no chaperone activity (i.e., PDIase activity)

toward reduced and denatured protein substrates (Sun et al., 2010b). Recombinant Deg8 demonstrates proteolytic activity toward photodamaged D1 and is able to produce the 16-kDa N-terminal and the 18-kDa C-terminal degradation products, which correspond to the cleavage products at the luminal CD loop (Sun et al., 2007). Although only recombinant Deg8 is proteolytically active, the *deg5* and *deg8* single knockout mutants of Arabidopsis both display impaired degradation of newly synthesized D1 and the impairment is more pronounced in the *deg5 deg8* double mutant (Sun et al., 2007). The defect in D1 turnover in the mutants is reflected in PSII activity. PSII in the *deg5* and *deg8* single mutants exhibits increased sensitivity to high light and the sensitivity is more obvious in the *deg5 deg8* double mutant. Under normal light, the two single mutants and the double mutant have a normal phenotype. Based on these data, Sun et al. (2007) proposed that Deg5 and Deg8 act as heterohexameric endopeptidases and cleave photodamaged D1 at the luminal CD loop during PSII repair (Figure 2). Deg5 has also been reported to be involved in wounding-related disposal of PsbF (Luciński et al., 2011a).

Deg7 (1097 amino acids at full length) is twice as long as most Deg proteases; it has two trypsin-like protease domains (one active and one degenerated) and four PDZ domains (three active and one degenerated) (Schuhmann et al., 2011). The domain composition suggests that Deg7 is the result of a whole-gene duplication event followed by subsequent degeneration (Schuhmann et al., 2011). Deg7 forms homotrimers and the oligomerization is mediated through the degenerated protease domain (Schuhmann et al., 2011). Recombinant Deg7 demonstrates proteolytic activity toward photodamaged D1, D2, CP43, and CP47 (Sun et al., 2010a). Compared to wild-type Arabidopsis, the *deg7* null mutant displays retarded growth, reduced PSII activity, and reduced degradation of PSII core proteins D1, D2, CP43, and CP47 under high light. However, under normal light, there is no apparent difference in plant growth or morphology between the *deg7* mutant and the wild type. These data suggest that Deg7 is involved in cleavage of photodamaged PSII core proteins D1, D2, CP47, and CP43 from the stroma side during PSII repair (Figure 2; Sun et al., 2010a).

HCF243, PSB27-H1, and PSB27-H2/LPA19 in Biogenesis, C-Terminal Processing, and/or Assembly of D1

HCF243 (High Chlorophyll Fluorescence 243) is an intrinsic thylakoid membrane protein with no recognizable domain or motif (Zhang et al., 2011). A loss-of-function mutation in the Arabidopsis *HCF243* gene causes substantial reductions in accumulation of PSII core subunits (D1, D2, CP43, and CP47) and assembly of PSII complexes (Zhang et al., 2011). Unlike PSII core subunits, the amounts of extrinsic PSII subunits, PSII antenna proteins, and other non-PSII thylakoid membrane proteins are hardly affected in the *hcf243* mutant. In line with these observations, PSII activity is dramatically reduced in the *hcf243* mutant and the mutant has pale-green leaves and a much smaller plant size than the wild type. Pulse-labeling experiments

indicated that these defects are caused by the severely reduced synthesis of D1 and to a lesser degree of D2 (Zhang et al., 2011). Indeed, the *hcf243* mutant over-accumulates pD1, the D1 precursor with an unprocessed C-terminus. HCF243 was also found to interact with D1 *in vivo* (Zhang et al., 2011). These data suggest that HCF243 is involved in biogenesis, processing, and assembly of D1 and possible biogenesis of D2 as well (Figures 1, 2).

PSB27s (PSII protein 27s) are thylakoid lumen proteins peripherally attached to thylakoid membranes (Chen et al., 2006a; Wei et al., 2010). Cyanobacterial Psb27 was found to facilitate assembly of the OEC manganese and plays a role in PSII repair (Nowaczyk et al., 2006; Liu et al., 2011a,b; Komenda et al., 2012b). The Arabidopsis genome encodes two PSB27s: PSB27-H1 and PSB27-H2/LPA19 (LPA19 stands for Low PSII Accumulation 19). Under normal light, the PSII protein composition is not changed and PSII activity is only slightly reduced in the Arabidopsis T-DNA insertion mutant of PSB27-H1 (Chen et al., 2006a; Dietzel et al., 2011). However, the *psb27-H1* mutant has reduced amounts of PSII-LHCII supercomplexes, suggesting that PSB27-H1 is required for the formation and stability of PSII-LHCII supercomplexes (Figure 1; Dietzel et al., 2011). Under high light, PSII activity and the amount of D1 decrease much faster in the *psb27-H1* mutant than in the wild type. In addition, the *psb27-H1* mutant displays delayed recovery of PSII activity after photoinhibition, suggesting that PSB27-H1 is involved in the repair cycle of photodamaged PSII. Unlike *psb27-H1*, the *psb27-H2/lpa19* mutants of Arabidopsis have pale-green leaves, a smaller plant size, and reduced PSII activity even under normal light conditions (Wei et al., 2010). Pulse-labeling experiments showed that C-terminal processing of D1 is impaired in the *psb27-H2/lpa19* null mutants. PSB27-H2/LPA19 was found to specifically interact with the soluble C-terminus of precursor and mature D1. It was concluded that PSB27-H2/LPA19 functions in C-terminal processing of D1 (Figure 1; Wei et al., 2010).

HCF136, PAM68, and PsbN/PBF1 in Assembly OF PSII Reaction Complexes

HCF136 (High Chlorophyll Fluorescence 136) is a luminal protein found in stroma lamellae; it contains no recognizable domain or motif (Meurer et al., 1998). The cyanobacterial homolog of HCF136, YCF48 (hypothetical chloroplast reading frame number 48), was found to interact with pD1, and to a lesser degree, partially processed and unassembled D1, but not with mature and unassembled D1 or D2, in a split-ubiquitin yeast-two-hybrid assay (Komenda et al., 2008). In addition, higher-plant HCF136 was found to be associated with the PSII precomplex D2-Cyt *b*₅₅₉ and PSII reaction-center complexes RC, RC47a, and RC47b (Plücker et al., 2002). The HCF136-deficient Arabidopsis mutant is able to accumulate PSI and cytochrome *b*_{6f} complex proteins, but unable to accumulate PSII proteins, perform PSII activity, or grow photoautotrophically (Meurer et al., 1998; Plücker et al., 2002). Pulse-labeling studies showed that the *hcf136* mutant is defective in biogenesis of PSII minimal reaction-center complexes, not in biosynthesis

of PSII proteins (Meurer et al., 1998; Plücker et al., 2002). Therefore, it was proposed that HCF136 is involved in assembly of PSII reaction-center complexes such as RC, RC47a, and RC47b (Figure 1). Consistent with this hypothesis, HCF136 was found to interact with another PSII assembly factor, PAM68 (Armbruster et al., 2010).

PAM68 in vascular plants is an integral thylakoid membrane protein with an acidic domain and a double-pass transmembrane domain (Armbruster et al., 2010). PAM68 was found to be associated with LMM complexes that are formed at an early step of PSII assembly (Armbruster et al., 2010). These complexes contain D1, D2, and LPA1/PratA, which may correspond to PSII minimal reaction-center complexes. PAM68 was found to interact with a number of PSII core subunits (D1, D2, CP43, CP47, PsbH, and PsbI) and PSII assembly factors (ALB3, HCF136, and LPA1/PratA, and LPA2). The PAM68-deficient *Arabidopsis* mutants have pale-green leaves, drastically reduced PSII activity, and severely retarded growth, under normal growth conditions. Consistent with these observations, the *pam68* mutants have severely reduced amounts of PSII core subunits, and they over-accumulate PSII reaction-center complexes at the expense of higher order PSII complexes. PAM68 was therefore proposed to be necessary for converting PSII minimal reaction-center complexes into larger PSII complexes (Figure 1; Armbruster et al., 2010). The *pam68* mutants were also found to over-accumulate pD1 (Armbruster et al., 2010), suggesting that PAM68 is also necessary for efficient C-terminal processing of D1 (Figures 1, 2).

PsbN is a small thylakoid membrane protein encoded by the plastid genome (Krech et al., 2013; Torabi et al., 2014). It was originally thought as a PSII subunit (Ikeuchi et al., 1989); however, two recent studies showed that PsbN is not a PSII subunit, but an assembly factor of PSII (Krech et al., 2013; Torabi et al., 2014). Thus, Krech et al. (2013) proposed a new name, Photosystem Biogenesis Factor 1 (PBF1) for this protein. Tobacco (*Nicotiana tabacum*) homoplasmic $\Delta psbN/pbf1$ mutants have pale-green leaves and slow autotrophic growth (Krech et al., 2013). Pulse-labeling and two-dimensional gel electrophoresis showed that formation of PSII pre-complexes, e.g., pD1-PsbI and D2-Cyt b_{559} , is not affected in the $\Delta psbN/pbf1$ mutants but assembly of PSII minimal reaction-center complexes and higher order PSII complexes is hampered (Torabi et al., 2014). It was concluded that PsbN/PBF1 is involved in formation of PSII minimal reaction-center complexes (Figure 1). The $\Delta psbN/pbf1$ mutants are extremely sensitive to light, even at relatively low light and PSII in the $\Delta psbN/pbf1$ mutants is unable to recover from photoinhibition (Torabi et al., 2014). These data indicate that PsbN/PBF1 may also function in PSII repair, which occurs in stroma lamellae. Consistent with this hypothesis, PsbN was found to be predominantly located in stroma lamellae (Torabi et al., 2014). Loss of PsbN changes the phosphorylation status of PSII core proteins and LHCI proteins, which is important for the migration of photodamaged PSII complexes from grana stacks to stroma-exposed thylakoids, and for balancing the light absorption capacity between PSI and PSII, respectively. Therefore, it is possible that PsbN is involved in PSII repair by regulating the

phosphorylation status of PSII core proteins and LHCI proteins (Figure 2).

PSB28 in Biogenesis of Chlorophyll-Binding Proteins Such as CP47, PsaA, and PsaB

PSB28 (PSII protein 28) is a small protein predominantly localized in the stroma/cytoplasm (Shi et al., 2012). In *Synechocystis* sp. PCC 6803, a small fraction of Psb28 was found to be associated with unassembled CP47, RC47, and PSII monomers at the stroma side (Dobáková et al., 2009; Boehm et al., 2012a). Deletion of this protein in *Synechocystis* sp. PCC 6803 cause a decreased amount of unassembled CP47 and increased amounts of PSII minimal reaction-center complexes and unassembled D1 (Dobáková et al., 2009). In addition, the $\Delta psb28$ mutant exhibits reduced synthesis of CP47 and PsaA/PsaB heterodimers. Therefore, Psb28 was considered to function in biogenesis and assembly of chlorophyll-containing proteins such as CP47, PsaA, and PsaB in cyanobacteria (Dobáková et al., 2009). Little is known about the role of PSB28 in photosynthetic eukaryotes except that PSB28 does exist in higher plants and that the absence of PSB28 results in a pale-green phenotype in rice (Jung et al., 2008; Mabbitt et al., 2014). However, because PSB28 is evolutionary conserved, it is reasonable to predict that PSB28 in higher plants may also function in biogenesis and assembly of chlorophyll-containing proteins such as CP47 (Figure 1).

LPA2 and LPA3 in Synthesis and Assembly of CP43

LPA2 is a small intrinsic thylakoid membrane protein with a C-terminal double-pass transmembrane domain (Ma et al., 2007). LPA3 (Low PSII Accumulation 3) is a chloroplast protein without any transmembrane domain or any other recognizable domain or motif. However, sub-chloroplast fractionation revealed that LPA3 could be located in chloroplast stroma or associated with thylakoid membranes (Cai et al., 2010). Although LPA2 and LPA3 are not homologous, they were identified as two auxiliary proteins assisting incorporation of CP43 into PSII via interaction with cpSRP translocase ALB3 (Ma et al., 2007; Cai et al., 2010). LPA2 and LPA3 were found to interact with CP43, ALB3, and each other (Cai et al., 2010). Pulse-labeling experiments showed that assembly from CP43-less reaction-center complexes to PSII monomers and formation of PSII-LHCI supercomplexes are distinctively slower in the *lpa2* and *lpa3* single mutants than in wild-type *Arabidopsis* (Ma et al., 2007; Cai et al., 2010). Synthesis of CP43 is greatly reduced in the *lpa2* and *lpa3* single mutants while synthesis of other PSII core subunits D1, D2, and CP47 is comparable between the single mutants and the wild type (Ma et al., 2007; Cai et al., 2010). LPA2 and LPA3 appear to be functional redundant because the *lpa2 lpa3* double mutant has no detectable amounts of D1, D2, CP43, and CP47 and is seedling-lethal. Taken together, LPA2 and LPA3 were proposed to be involved in synthesis and assembly of CP43 (Figures 1, 2; Ma et al., 2007; Cai et al., 2010).

PSB33 in Association of LHCII with PSII

PSB33 (PSII protein 33) is a thylakoid membrane protein with an N-terminal Rieske-type domain exposed to the stroma side, a double-pass transmembrane domain, and a C-terminal partial chlorophyll-binding domain (Fristedt et al., 2015). PSB33 was found to co-migrate with PSII-LHCII supercomplexes, PSII dimers, PSII monomers, and CP43-less PSII monomers in BN-PAGE (Fristedt et al., 2015). Loss-of-function mutations in the Arabidopsis *PSB33* gene cause reduced amounts of PSII-LHCII supercomplexes, and increased amounts of PSII dimers, lower state transition, lower non-photochemical quenching, increased photosensitivity, and retarded growth (Fristedt et al., 2015). According to these data, Fristedt et al. (2015) proposed that PSB33 may mediate the association of LHCII with PSII core complexes (Figure 1) and balance the light absorption capacity between PSII and PSI. It is likely that PSB33 also functions in attaching LHCII to PSII during reassembly of repaired PSII (Figure 2), because the defect in PSII activity in the *psb33* mutants is more pronounced under higher light (Fristedt et al., 2015).

HHL1 and MPH1 in Protection of PSII from Photodamage

HHL1 (Hypersensitive to High Light 1) is a thylakoid membrane protein with a single-pass transmembrane domain, and a C-terminal partial von Willebrand factor type A domain, which is known to mediate protein-protein interactions (Jin et al., 2014). Under high light, the HHL1-deficient mutants have lower efficiency of PSII photochemistry, lower amounts of PSII core subunits and PSII-LHCII supercomplexes, and higher amounts of reactive oxygen species, than wild-type Arabidopsis (Jin et al., 2014). Many of these defects become milder under normal light. Therefore, it was proposed that HHL1 is involved in the repair and reassembly cycle of photodamaged PSII. Consistent with this hypothesis, HHL1 was found in both grana stacks and stroma lamellae, and PSII core subunits in thylakoid membranes isolated from HHL1-deficient plants were found to be less stable than those isolated from wild-type plants (Jin et al., 2014). Jin et al. (2014) also observed *in vivo* and *in vitro* interaction between HHL1 and LQY1, another protein involved in PSII repair. Because the majority of HHL1 is associated with PSII core monomers, it is likely that HHL1 “collaborates” with LQY1 and assists the reassembly of PSII core monomers and PSII-LHCII supercomplexes during PSII repair (Figure 2). In line with this hypothesis, the *hhl1 lqy1* double mutant is more sensitive to high light than the single mutants (Jin et al., 2014).

MPH1 (Maintenance of PSII under High light 1) is a proline-rich intrinsic thylakoid membrane protein with a single-pass transmembrane domain; it is present in grana stacks, grana margins, and stroma lamellae (Liu and Last, 2015a,b). Under normal light, PSII activity and the composition of PSII complexes

in MPH1-deficient mutants are similar to those in wild-type Arabidopsis. Under high light, the *mph1* mutants have lower efficiency of PSII photochemistry, and lower amounts of PSII-LHCII supercomplexes, PSII dimers, and PSII core monomers, than the wild type. Therefore, it was proposed that MPH1 has a role in protection and/or stabilization of PSII under high light (Liu and Last, 2015a,b). Consistent with this hypothesis, MPH1 was found to change its association under different light intensities: under normal light, the majority of MPH1 is associated with PSII core monomers; under high light, the majority of MPH1 becomes associated with PSII-LHCII supercomplexes (Liu and Last, 2015a). Because of the interactions between MPH1 and different PSII complexes under different light irradiance and the reductions in the amounts of PSII monomers and higher order PSII complexes, it is likely that MPH1 is involved in the assembly and/or stability of PSII core monomers and higher order PSII complexes under high light (Figure 2).

CONCLUDING REMARKS

Photosynthesis directly or indirectly provides chemical energy for nearly all life forms on earth. Due to the importance of photosynthesis, the structure, biogenesis, and maintenance of the photosynthetic apparatus have long been one of the major focuses of research. The combination of proteomics, X-ray crystallography, and single-particle electron cryo-microscopy approaches has led to a comprehensive understanding of the structure and subunit composition of PSII. In addition, significant progresses have been made in the identification and functional studies of protein factors that are involved in *de novo* assembly and/or the repair and reassembly cycle of PSII. The inclusion of thylakoid protein trafficking/targeting systems and enzymes that catalyze important enzymatic steps, along with various assembly/stability factors allows a more comprehensive view of recent advances in this field. However, additional efforts are in great need to (1) dissect the precise functions of understudied assembly/stability factors or enzymes; and (2) build a protein interactome network that would provide a systems view of the interplay among different assembly/stability factors, enzymes, thylakoid protein trafficking/targeting systems, PSII assembly and repair complexes, and PSII subunits.

AUTHOR CONTRIBUTIONS

This review was written entirely by YL.

ACKNOWLEDGMENTS

This work was supported by the US National Science Foundation Grant MCB-1244008.

REFERENCES

- Adamska, I., and Kloppstech, K. (1991). Evidence for an association of the early light-inducible protein (ELIP) of pea with Photosystem II. *Plant Mol. Biol.* 16, 209–223. doi: 10.1007/BF00020553
- Albiniak, A. M., Baglieri, J., and Robinson, C. (2012). Targeting of lumenal proteins across the thylakoid membrane. *J. Exp. Bot.* 63, 1689–1698. doi: 10.1093/jxb/err444
- Albrecht, V., Ingenfeld, A., and Apel, K. (2008). *Snowy cotyledon 2*: the identification of a zinc finger domain protein essential for chloroplast development in cotyledons but not in true leaves. *Plant Mol. Biol.* 66, 599–608. doi: 10.1007/s11103-008-9291-y
- Anbudurai, P. R., Mor, T. S., Ohad, I., Shestakov, S. V., and Pakrasi, H. B. (1994). The *ctpA* gene encodes the C-terminal processing protease for the D1 protein of the Photosystem II reaction center complex. *Proc. Natl. Acad. Sci. U.S.A.* 91, 8082–8086. doi: 10.1073/pnas.91.17.8082
- Andersson, U., Heddad, M., and Adamska, I. (2003). Light stress-induced one-helix protein of the chlorophyll *a/b*-binding family associated with Photosystem I. *Plant Physiol.* 132, 811–820. doi: 10.1104/pp.102.019281
- Armbruster, U., Zühlke, J., Rengstl, B., Kreller, R., Makarenko, E., Ruhle, T., et al. (2010). The *Arabidopsis* thylakoid protein PAM68 is required for efficient D1 biogenesis and Photosystem II assembly. *Plant Cell* 22, 3439–3460. doi: 10.1105/tpc.110.077453
- Aro, E.-M., Suorsa, M., Rokka, A., Allahverdiyeva, Y., Paakkarinen, V., Saleem, A., et al. (2005). Dynamics of Photosystem II: a proteomic approach to thylakoid protein complexes. *J. Exp. Bot.* 56, 347–356. doi: 10.1093/jxb/eri041
- Bailey, S., Thompson, E., Nixon, P. J., Horton, P., Mullineaux, C. W., Robinson, C., et al. (2002). A critical role for the Var2 FtsH homologue of *Arabidopsis thaliana* in the Photosystem II repair cycle *in vivo*. *J. Biol. Chem.* 277, 2006–2011. doi: 10.1074/jbc.M105878200
- Bellafore, S., Barneche, F., Peltier, G., and Rochaix, J. D. (2005). State transitions and light adaptation require chloroplast thylakoid protein kinase STN7. *Nature* 433, 892–895. doi: 10.1038/nature03286
- Bhuiyan, N. H., Friso, G., Poliakov, A., Ponnala, L., and van Wijk, K. J. (2015). MET1 is a thylakoid-associated TPR protein involved in Photosystem II supercomplex formation and repair in *Arabidopsis*. *Plant Cell* 27, 262–285. doi: 10.1105/tpc.114.132787
- Boehm, M., Yu, J., Krynicka, V., Barker, M., Tichy, M., Komenda, J., et al. (2012b). Subunit organization of a *Synechocystis* hetero-oligomeric thylakoid FtsH complex involved in Photosystem II repair. *Plant Cell* 24, 3669–3683. doi: 10.1105/tpc.112.100891
- Boehm, M., Yu, J., Reisinger, V., Beckova, M., Eichacker, L. A., Schlodder, E., et al. (2012a). Subunit composition of CP43-less Photosystem II complexes of *Synechocystis* sp. PCC 6803: implications for the assembly and repair of Photosystem II. *Phil. Trans. R. Soc. B* 367, 3444–3454. doi: 10.1098/rstb.2012.0066
- Bohne, A.-V., Schwarz, C., Schottkowski, M., Lidschreiber, M., Piotrowski, M., Zerges, W., et al. (2013). Reciprocal regulation of protein synthesis and carbon metabolism for thylakoid membrane biogenesis. *PLoS Biol.* 11:e1001482. doi: 10.1371/journal.pbio.1001482
- Bonardi, V., Pesaresi, P., Becker, T., Schleiff, E., Wagner, R., Pfannschmidt, T., et al. (2005). Photosystem II core phosphorylation and photosynthetic acclimation require two different protein kinases. *Nature* 437, 1179–1182. doi: 10.1038/nature04016
- Bricker, T., Roose, J. L., Zhang, P., and Frankel, L. (2013). The PsbP family of proteins. *Photosynth. Res.* 116, 235–250. doi: 10.1007/s11120-013-9820-7
- Bricker, T. M., Roose, J. L., Fagerlund, R. D., Frankel, L. K., and Eaton-Rye, J. J. (2012). The extrinsic proteins of Photosystem II. *Biochim. Biophys. Acta* 1817, 121–142. doi: 10.1016/j.bbabi.2011.07.006
- Cai, W., Ma, J., Chi, W., Zou, M., Guo, J., Lu, C., et al. (2010). Cooperation of LPA3 and LPA2 is essential for Photosystem II assembly in *Arabidopsis*. *Plant Physiol.* 154, 109–120. doi: 10.1104/pp.110.159558
- Cai, W., Ma, J., Guo, J., and Zhang, L. (2008). Function of ROC4 in the efficient repair of photodamaged Photosystem II in *Arabidopsis*. *Photochem. Photobiol.* 84, 1343–1348. doi: 10.1111/j.1751-1097.2008.00448.x
- Cain, P., Hall, M., Schröder, W. P., Kieselbach, T., and Robinson, C. (2009). A novel extended family of stromal thioredoxins. *Plant Mol. Biol.* 70, 273–281. doi: 10.1007/s11103-009-9471-4
- Calderon, R. H., García-Cerdán, J. G., Malnoë, A., Cook, R., Russell, J. J., Gaw, C., et al. (2013). A Conserved rubredoxin is necessary for Photosystem II accumulation in diverse oxygenic photoautotrophs. *J. Biol. Chem.* 288, 26688–26696. doi: 10.1074/jbc.M113.487629
- Casazza, A. P., Rossini, S., Rosso, M. G., and Soave, C. (2005). Mutational and expression analysis of ELIP1 and ELIP2 in *Arabidopsis thaliana*. *Plant Mol. Biol.* 58, 41–51. doi: 10.1007/s11103-005-4090-1
- Charuvi, D., Kiss, V., Nevo, R., Shimoni, E., Adam, Z., and Reich, Z. (2012). Gain and loss of photosynthetic membranes during plastid differentiation in the shoot apex of *Arabidopsis*. *Plant Cell* 24, 1143–1157. doi: 10.1105/tpc.111.094458
- Chassin, Y., Kapri-Pardes, E., Sinvany, G., Arad, T., and Adam, Z. (2002). Expression and characterization of the thylakoid lumen protease DegP1 from *Arabidopsis*. *Plant Physiol.* 130, 857–864. doi: 10.1104/pp.007922
- Che, Y., Fu, A., Hou, X., McDonald, K., Buchanan, B. B., Huang, W., et al. (2013). C-terminal processing of reaction center protein D1 is essential for the function and assembly of Photosystem II in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 110, 16247–16252. doi: 10.1073/pnas.1313894110
- Chen, H., Zhang, D., Guo, J., Wu, H., Jin, M., Lu, Q., et al. (2006a). A Psb27 homologue in *Arabidopsis thaliana* is required for efficient repair of photodamaged Photosystem II. *Plant Mol. Biol.* 61, 567–575. doi: 10.1007/s11103-006-0031-x
- Chen, J., Burke, J. J., Velten, J., and Xin, Z. (2006b). FtsH11 protease plays a critical role in *Arabidopsis* thermotolerance. *Plant J.* 48, 73–84. doi: 10.1111/j.1365-313X.2006.02855.x
- Chen, M., Choi, Y., Voytas, D. F., and Rodermel, S. (2000). Mutations in the *Arabidopsis* VAR2 locus cause leaf variegation due to the loss of a chloroplast FtsH protease. *Plant J.* 22, 303–313. doi: 10.1046/j.1365-313x.2000.00738.x
- Chidgey, J. W., Linhartová, M., Komenda, J., Jackson, P. J., Dickman, M. J., Canniffe, D. P., et al. (2014). A cyanobacterial chlorophyll synthase-HliD complex associates with the Ycf39 protein and the YidC/Alb3 insertase. *Plant Cell* 26, 1267–1279. doi: 10.1105/tpc.114.124495
- Cline, K., and Dabney-Smith, C. (2008). Plastid protein import and sorting: different paths to the same compartments. *Curr. Opin. Plant Biol.* 11, 585–592. doi: 10.1016/j.pbi.2008.10.008
- Cline, K., and Theg, S. (2007). “The Sec and Tat protein translocation pathways in chloroplasts,” in *The Enzymes: Molecular Machines Involved in Protein Transport across Cellular Membranes*, eds R. E. Dalbey, C. M. Koehler, and F. Tamanoi (Waltham, MA: Academic Press), 463–492.
- da Fonseca, P., Morris, E. P., Hankamer, B., and Barber, J. (2002). Electron crystallographic study of Photosystem II of the cyanobacterium *Synechococcus elongatus*. *Biochemistry* 41, 5163–5167. doi: 10.1021/bi0120650
- Dall’Osto, L., Bressan, M., and Bassi, R. (2015). Biogenesis of light harvesting proteins. *Biochim. Biophys. Acta* 1847, 861–871. doi: 10.1016/j.bbabi.2015.02.009
- Dietzel, L., Bräutigam, K., Steiner, S., Schöffler, K., Lepetit, B., Grimm, B., et al. (2011). Photosystem II supercomplex remodeling serves as an entry mechanism for state transitions in *Arabidopsis*. *Plant Cell* 23, 2964–2977. doi: 10.1105/tpc.111.087049
- Dobáková, M., Sobotka, R., Tichy, M., and Komenda, J. (2009). Psb28 protein is involved in the biogenesis of the Photosystem II inner antenna CP47 (PsbB) in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Physiol.* 149, 1076–1086. doi: 10.1104/pp.108.130039
- Dominguez-Solis, J. R., He, Z. Y., Lima, A., Ting, J., Buchanan, B. B., and Luan, S. (2008). A cyclophilin links redox and light signals to cysteine biosynthesis and stress responses in chloroplasts. *Proc. Natl. Acad. Sci. U.S.A.* 105, 16386–16391. doi: 10.1073/pnas.0808204105
- Feige, M. J., and Hendershot, L. M. (2011). Disulfide bonds in ER protein folding and homeostasis. *Curr. Opin. Cell Biol.* 23, 167–175. doi: 10.1016/j.ceb.2010.10.012
- Feng, W.-K., Wang, L., Lu, Y., and Wang, X.-Y. (2011). A protein oxidase catalysing disulfide bond formation is localized to the chloroplast thylakoids. *FEBS J.* 278, 3419–3430. doi: 10.1111/j.1742-4658.2011.08265.x
- Fischer, G., Tradler, T., and Zarnt, T. (1998). The mode of action of peptidyl prolyl *cis/trans* isomerases *in vivo*: binding vs. catalysis. *FEBS Lett.* 426, 17–20. doi: 10.1016/S0014-5793(98)00242-7
- Fristed, R., Herdean, A., Blaby-Haas, C. E., Mamedov, F., Merchant, S. S., Last, R. L., et al. (2015). PHOTOSYSTEM II PROTEIN33, a protein conserved

- in the plastid lineage, is associated with the chloroplast thylakoid membrane and provides stability to Photosystem II supercomplexes in *Arabidopsis*. *Plant Physiol.* 167, 481–492. doi: 10.1104/pp.114.253336
- Fu, A., He, Z. Y., Cho, H. S., Lima, A., Buchanan, B. B., and Luan, S. (2007). A chloroplast cyclophilin functions in the assembly and maintenance of Photosystem II in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U.S.A.* 104, 15947–15952. doi: 10.1073/pnas.0707851104
- Fulgosi, H., Vener, A. V., Altschmied, L., Herrmann, R. G., and Andersson, B. (1998). A novel multi-functional chloroplast protein: identification of a 40 kDa immunophilin-like protein located in the thylakoid lumen. *EMBO J.* 17, 1577–1587. doi: 10.1093/emboj/17.6.1577
- Göhre, V., Ossenbühl, F., Crèvecoeur, M., Eichacker, L. A., and Rochaix, J.-D. (2006). One of two Alb3 proteins is essential for the assembly of the photosystems and for cell survival in *Chlamydomonas*. *Plant Cell* 18, 1454–1466. doi: 10.1105/tpc.105.038695
- Goral, T. K., Johnson, M. P., Brain, A. P., Kirchoff, H., Ruban, A. V., and Mullineaux, C. W. (2010). Visualizing the mobility and distribution of chlorophyll proteins in higher plant thylakoid membranes: effects of photoinhibition and protein phosphorylation. *Plant J.* 62, 948–959. doi: 10.1111/j.1365-3113x.2010.04207.x
- Haubühl, K., Andersson, B., and Adamska, I. (2001). A chloroplast DegP2 protease performs the primary cleavage of the photodamaged D1 protein in plant Photosystem II. *EMBO J.* 20, 713–722. doi: 10.1093/emboj/20.4.713
- He, Z., Li, L., and Luan, S. (2004). Immunophilins and parvulins. Superfamily of peptidyl prolyl isomerases in *Arabidopsis*. *Plant Physiol.* 134, 1248–1267. doi: 10.1104/pp.103.031005
- Heddad, M., Engelken, J., and Adamska, I. (2012). “Light stress proteins in viruses, cyanobacteria and photosynthetic eukaryota,” in *Photosynthesis*, eds J. J. Eaton-Rye, B. C. Tripathy, and T. D. Sharkey (Dordrecht: Springer), 299–317.
- Heddad, M., Norén, H., Reiser, V., Dunaeva, M., Andersson, B., and Adamska, I. (2006). Differential expression and localization of early light-induced proteins in *Arabidopsis*. *Plant Physiol.* 142, 75–87. doi: 10.1104/pp.106.081489
- Henry, R., Goforth, R., and Schunemann, D. (2007). “Chloroplast SRP/FtsY and Alb3 in protein integration into the thylakoid membrane,” in *The Enzymes: Molecular Machines Involved in Protein Transport across Cellular Membranes*, eds R. E. Dalbey, C. M. Koehler, and F. Tamanoi (Waltham, MA: Academic Press), 493–521.
- Hentze, M. W. (1994). Enzymes as RNA-binding proteins: a role for (di)nucleotide-binding domains? *Trends Biochem. Sci.* 19, 101–103. doi: 10.1016/0968-0004(94)90198-8
- Herbstová, M., Tietz, S., Kinzel, C., Turkina, M. V., and Kirchoff, H. (2012). Architectural switch in plant photosynthetic membranes induced by light stress. *Proc. Natl. Acad. Sci. U.S.A.* 109, 20130–20135. doi: 10.1073/pnas.1214265109
- Hooper, J. K., Boyd, C. O., and Paavola, L. G. (1991). Origin of thylakoid membranes in *Chlamydomonas reinhardtii* *y-1* at 38°C. *Plant Physiol.* 96, 1321–1328. doi: 10.1104/pp.96.4.1321
- Hooper, J. K., Eggink, L. L., and Chen, M. (2007). Chlorophylls, ligands and assembly of light-harvesting complexes in chloroplasts. *Photosynth. Res.* 94, 387–400. doi: 10.1007/s11120-007-9181-1
- Houston, N. L., Fan, C., Xiang, Q.-Y., Schulze, J.-M., Jung, R., and Boston, R. S. (2005). Phylogenetic analyses identify 10 classes of the protein disulfide isomerase family in plants, including single-domain protein disulfide isomerase-related proteins. *Plant Physiol.* 137, 762–778. doi: 10.1104/pp.104.056507
- Huang, J., Taylor, J. P., Chen, J. G., Uhrig, J. F., Schnell, D. J., Nakagawa, T., et al. (2006). The plastid protein THYLAKOID FORMATION1 and the plasma membrane G-protein GPA1 interact in a novel sugar-signaling mechanism in *Arabidopsis*. *Plant Cell* 18, 1226–1238. doi: 10.1105/tpc.105.037259
- Huang, W., Chen, Q., Zhu, Y., Hu, F., Zhang, L., Ma, Z., et al. (2013). *Arabidopsis* thylakoid formation 1 is a critical regulator for dynamics of PSII-LHCII complexes in leaf senescence and excess light. *Mol. Plant* 6, 1673–1691. doi: 10.1093/mp/sst069
- Huesgen, P. F., Schuhmann, H., and Adamska, I. (2005). The family of Deg proteases in cyanobacteria and chloroplasts of higher plants. *Physiol. Plant.* 123, 413–420. doi: 10.1111/j.1399-3054.2005.00458.x
- Huesgen, P. F., Schuhmann, H., and Adamska, I. (2006). Photodamaged D1 protein is degraded in *Arabidopsis* mutants lacking the Deg2 protease. *FEBS Lett.* 580, 6929–6932. doi: 10.1016/j.febslet.2006.11.058
- Huesgen, P. F., Schuhmann, H., and Adamska, I. (2009). Deg/HtrA proteases as components of a network for Photosystem II quality control in chloroplasts and cyanobacteria. *Res. Microbiol.* 160, 726–732. doi: 10.1016/j.resmic.2009.08.005
- Hutin, C., Nussaume, L., Moise, N., Moya, I., Kloppstech, K., and Havaux, M. (2003). Early light-induced proteins protect *Arabidopsis* from photooxidative stress. *Proc. Natl. Acad. Sci. U.S.A.* 100, 4921–4926. doi: 10.1073/pnas.0736939100
- Ifuku, K. (2014). The PsbP and PsbQ family proteins in the photosynthetic machinery of chloroplasts. *Plant Physiol. Biochem.* 81, 108–114. doi: 10.1016/j.plaphy.2014.01.001
- Ikeuchi, M., Koike, H., and Inoue, Y. (1989). N-terminal sequencing of low-molecular-mass components in cyanobacterial Photosystem II core complex: two components correspond to unidentified open reading frames of plant chloroplast DNA. *FEBS Lett.* 253, 178–182. doi: 10.1016/0014-5793(89)80954-8
- Ingelsson, B., Shapiguzov, A., Kieselbach, T., and Vener, A. V. (2009). Peptidyl prolyl isomerase activity in chloroplast thylakoid lumen is a dispensable function of immunophilins in *Arabidopsis thaliana*. *Plant Cell Physiol.* 50, 1801–1814. doi: 10.1093/pcp/pcp1122
- Ishihara, S., Takabayashi, A., Ido, K., Endo, T., Ifuku, K., and Sato, F. (2007). Distinct functions for the two PsbP-like proteins PPL1 and PPL2 in the chloroplast thylakoid lumen of *Arabidopsis*. *Plant Physiol.* 145, 668–679. doi: 10.1104/pp.107.105866
- Ishikawa, A., Tanaka, H., Kato, C., Iwasaki, Y., and Asahi, T. (2005a). Molecular characterization of the *ZKT* gene encoding a protein with PDZ, K-Box, and TPR motifs in *Arabidopsis*. *Biosci. Biotechnol. Biochem.* 69, 972–978. doi: 10.1271/bbb.69.972
- Ishikawa, Y., Schroder, W. P., and Funk, C. (2005b). Functional analysis of the PsbP-like protein (sll1418) in *Synechocystis* sp. PCC 6803. *Photosynth. Res.* 84, 257–262. doi: 10.1007/s11120-005-0477-8
- Jackson, S. A., Hinds, M. G., and Eaton-Rye, J. J. (2012). Solution structure of CyanoP from *Synechocystis* sp. PCC 6803: new insights on the structural basis for functional specialization amongst PsbP family proteins. *Biochim. Biophys. Acta* 1817, 1331–1338. doi: 10.1016/j.bbabi.2012.02.032
- Jansson, S. (1999). A guide to the Lhc genes and their relatives in *Arabidopsis*. *Trends Plant Sci.* 4, 236–240. doi: 10.1016/S1360-1385(99)01419-3
- Järvi, S., Gollan, P. J., and Aro, E.-M. (2013). Understanding the roles of the thylakoid lumen in photosynthesis regulation. *Front. Plant Sci.* 4:434. doi: 10.3389/fpls.2013.00434
- Järvi, S., Suorsa, M., and Aro, E.-M. (2015). Photosystem II repair in plant chloroplasts—Regulation, assisting proteins and shared components with Photosystem II biogenesis. *Biochim. Biophys. Acta* 1847, 900–909. doi: 10.1016/j.bbabi.2015.01.006
- Jessop, C. E., Chakravarthi, S., Garbi, N., Hämmerling, G. J., Lovell, S., and Bulleid, N. J. (2007). ERp57 is essential for efficient folding of glycoproteins sharing common structural domains. *EMBO J.* 26, 28–40. doi: 10.1038/sj.emboj.7601505
- Jin, H., Liu, B., Luo, L., Feng, D., Wang, P., Liu, J., et al. (2014). HYPERSENSITIVE TO HIGH LIGHT1 interacts with LOW QUANTUM YIELD OF PHOTOSYSTEM III and functions in protection of Photosystem II from photodamage in *Arabidopsis*. *Plant Cell* 26, 1213–1229. doi: 10.1105/tpc.113.122424
- Jung, K.-H., Lee, J., Dardick, C., Seo, Y.-S., Cao, P., Canlas, P., et al. (2008). Identification and functional analysis of light-responsive unique genes and gene family members in rice. *PLoS Genet.* 4:e1000164. doi: 10.1371/journal.pgen.1000164
- Kapri-Pardes, E., Naveh, L., and Adam, Z. (2007). The thylakoid lumen protease Deg1 is involved in the repair of Photosystem II from photoinhibition in *Arabidopsis*. *Plant Cell* 19, 1039–1047. doi: 10.1105/tpc.106.046573
- Karamoko, M., Cline, S., Redding, K., Ruiz, N., and Hamel, P. P. (2011). Lumen Thiol Oxidoreductase1, a disulfide bond-forming catalyst, is required for the assembly of Photosystem II in *Arabidopsis*. *Plant Cell* 23, 4462–4475. doi: 10.1105/tpc.111.089680
- Karamoko, M., Gabilly, S., and Hamel, P. P. (2013). Operation of trans-thylakoid thiol-metabolizing pathways in photosynthesis. *Front. Plant Sci.* 4:476. doi: 10.3389/fpls.2013.00476

- Karim, S., Alezzawi, M., Garcia-Petit, C., Solymosi, K., Khan, N. Z., Lindquist, E., et al. (2014). A novel chloroplast localized Rab GTPase protein CPRabA5e is involved in stress, development, thylakoid biogenesis and vesicle transport in *Arabidopsis*. *Plant Mol. Biol.* 84, 675–692. doi: 10.1007/s11103-013-0161-x
- Karim, S., and Aronsson, H. (2014). The puzzle of chloroplast vesicle transport – involvement of GTPases. *Front. Plant Sci.* 5:472. doi: 10.3389/fpls.2014.00472
- Kashino, Y., Lauber, W. M., Carroll, J. A., Wang, Q. J., Whitmarsh, J., Satoh, K., et al. (2002). Proteomic analysis of a highly active Photosystem II preparation from the cyanobacterium *Synechocystis* sp. PCC 6803 reveals the presence of novel polypeptides. *Biochemistry* 41, 8004–8012. doi: 10.1021/bi026012+
- Kato, Y., Miura, E., Ido, K., Ifuku, K., and Sakamoto, W. (2009). The variegated mutants lacking chloroplastic FtsHs are defective in D1 degradation and accumulate reactive oxygen species. *Plant Physiol.* 151, 1790–1801. doi: 10.1104/pp.109.146589
- Kato, Y., Miura, E., Matsushima, R., and Sakamoto, W. (2007). White leaf sectors in *yellow variegated2* are formed by viable cells with undifferentiated plastids. *Plant Physiol.* 144, 952–960. doi: 10.1104/pp.107.099002
- Kato, Y., Sun, X., Zhang, L., and Sakamoto, W. (2012). Cooperative D1 degradation in the Photosystem II repair mediated by chloroplastic proteases in *Arabidopsis*. *Plant Physiol.* 159, 1428–1439. doi: 10.1104/pp.112.199042
- Keren, N., Ohkawa, H., Welsh, E. A., Liberton, M., and Pakrasi, H. B. (2005). Psb29, a conserved 22-kD protein, functions in the biogenesis of Photosystem II complexes in *Synechocystis* and *Arabidopsis*. *Plant Cell* 17, 2768–2781. doi: 10.1105/tpc.105.035048
- Khan, N. Z., Lindquist, E., and Aronsson, H. (2013). New putative chloroplast vesicle transport components and cargo proteins revealed using a bioinformatics approach: an *Arabidopsis* model. *PLoS ONE* 8:e59898. doi: 10.1371/journal.pone.0059898
- Khatoon, M., Inagawa, K., Pospisil, P., Yamashita, A., Yoshioka, M., Lundin, B., et al. (2009). Quality control of Photosystem II: Thylakoid unstacking is necessary to avoid further damage to the D1 protein and to facilitate D1 degradation under light stress in spinach thylakoids. *J. Biol. Chem.* 284, 25343–25352. doi: 10.1074/jbc.M109.007740
- Klinkert, B., Ossenbuhl, F., Sikorski, M., Berry, S., Eichacker, L., and Nickelsen, J. (2004). Prata, a periplasmic tetratricopeptide repeat protein involved in biogenesis of Photosystem II in *Synechocystis* sp. PCC 6803. *J. Biol. Chem.* 279, 44639–44644. doi: 10.1074/jbc.M405393200
- Knoppová, J., Sobotka, R., Tich, M., Yu, J., Konik, P., Halada, P., et al. (2014). Discovery of a chlorophyll binding protein complex involved in the early steps of Photosystem II assembly in *Synechocystis*. *Plant Cell* 26, 1200–1212. doi: 10.1105/tpc.114.123919
- Komenda, J., Knoppová, J., Kopečná, J., Sobotka, R., Halada, P., Yu, J., et al. (2012b). The Psb27 assembly factor binds to the CP43 complex of Photosystem II in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Physiol.* 158, 476–486. doi: 10.1104/pp.111.184184
- Komenda, J., Nickelsen, J., Tich, M., Prášil, O., Eichacker, L. A., and Nixon, P. J. (2008). The cyanobacterial homologue of HCF136/YCF48 is a component of an early Photosystem II assembly complex and is important for both the efficient assembly and repair of Photosystem II in *Synechocystis* sp. PCC 6803. *J. Biol. Chem.* 283, 22390–22399. doi: 10.1074/jbc.M801917200
- Komenda, J., Sobotka, R., and Nixon, P. J. (2012a). Assembling and maintaining the Photosystem II complex in chloroplasts and cyanobacteria. *Curr. Opin. Plant Biol.* 15, 245–251. doi: 10.1016/j.pbi.2012.01.017
- Krech, K., Fu, H.-Y., Thiele, W., Ruf, S., Schöttler, M. A., and Bock, R. (2013). Reverse genetics in complex multigene operons by co-transformation of the plastid genome and its application to the open reading frame previously designated *psbN*. *Plant J.* 75, 1062–1074. doi: 10.1111/tj.12256
- Kroll, D., Meierhoff, K., Bechtold, N., Kinoshita, M., Westphal, S., Vothknecht, U. C., et al. (2001). *VIPPI1*, a nuclear gene of *Arabidopsis thaliana* essential for thylakoid membrane formation. *Proc. Natl. Acad. Sci. U.S.A.* 98, 4238–4242. doi: 10.1073/pnas.061500998
- Krzywdka, S., Brzozowski, A. M., Verma, C., Karata, K., Ogura, T., and Wilkinson, A. J. (2002). The crystal structure of the AAA domain of the ATP-dependent protease FtsH of *Escherichia coli* at 1.5 Å resolution. *Structure* 10, 1073–1083. doi: 10.1016/S0969-2126(02)00806-7
- Kwon, K.-C., and Cho, M. H. (2008). Deletion of the chloroplast-localized *AtTerC* gene product in *Arabidopsis thaliana* leads to loss of the thylakoid membrane and to seedling lethality. *Plant J.* 55, 428–442. doi: 10.1111/j.1365-313X.2008.03523.x
- Liao, D. I., Qian, J., Chisholm, D. A., Jordan, D. B., and Diner, B. A. (2000). Crystal structures of the Photosystem II D1 C-terminal processing protease. *Nat. Struct. Biol.* 7, 749–753. doi: 10.1038/78973
- Lima, A., Lima, S., Wong, J. H., Phillips, R. S., Buchanan, B. B., and Luan, S. (2006). A redox-active FKBP-type immunophilin functions in accumulation of the Photosystem II supercomplex in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U.S.A.* 103, 12631–12636. doi: 10.1073/pnas.0605452103
- Lindahl, M., Spetea, C., Hundal, T., Oppenheim, A. B., Adam, Z., and Andersson, B. (2000). The thylakoid FtsH protease plays a role in the light-induced turnover of the Photosystem II D1 protein. *Plant Cell* 12, 419–432. doi: 10.1105/tpc.12.3.419
- Lindahl, M., Tabak, S., Cseke, L., Pichersky, E., Andersson, B., and Adam, Z. (1996). Identification, characterization, and molecular cloning of a homologue of the bacterial FtsH protease in chloroplasts of higher plants. *J. Biol. Chem.* 271, 29329–29334. doi: 10.1074/jbc.271.46.29329
- Link, S., Engelmann, K., Meierhoff, K., and Westhoff, P. (2012). The atypical short-chain dehydrogenases HCF173 and HCF244 are jointly involved in translational initiation of the *psbA* mRNA of *Arabidopsis*. *Plant Physiol.* 160, 2202–2218. doi: 10.1104/pp.112.205104
- Lippuner, V., Chou, I. T., Scott, S. V., Ettinger, W. F., Theg, S. M., and Gasser, C. S. (1994). Cloning and characterization of chloroplast and cytosolic forms of cyclophilin from *Arabidopsis thaliana*. *J. Biol. Chem.* 269, 7863–7868.
- Liu, H., Huang, R. Y.-C., Chen, J., Gross, M. L., and Pakrasi, H. B. (2011b). Psb27, a transiently associated protein, binds to the chlorophyll binding protein CP43 in Photosystem II assembly intermediates. *Proc. Natl. Acad. Sci. U.S.A.* 108, 18536–18541. doi: 10.1073/pnas.1111597108
- Liu, H., Roose, J. L., Cameron, J. C., and Pakrasi, H. B. (2011a). A genetically tagged Psb27 protein allows purification of two consecutive Photosystem II (PSII) assembly intermediates in *Synechocystis* 6803, a cyanobacterium. *J. Biol. Chem.* 286, 24865–24871. doi: 10.1074/jbc.M111.246231
- Liu, J., and Last, R. L. (2015a). A land plant-specific thylakoid membrane protein contributes to Photosystem II maintenance in *Arabidopsis thaliana*. *Plant J.* 82, 731–743. doi: 10.1111/tj.12845
- Liu, J., and Last, R. L. (2015b). MPH1 is a thylakoid membrane protein involved in protecting Photosystem II from photodamage in land plants. *Plant Signal. Behav.* 10:e1076602. doi: 10.1080/15592324.2015.1076602
- Liu, L.-N., Chen, X.-L., Zhang, Y.-Z., and Zhou, B.-C. (2005). Characterization, structure and function of linker polypeptides in phycobilisomes of cyanobacteria and red algae: an overview. *Biochim. Biophys. Acta* 1708, 133–142. doi: 10.1016/j.bbabi.2005.04.001
- Liu, Z., Yan, H., Wang, K., Kuang, T., Zhang, J., Gui, L., et al. (2004). Crystal structure of spinach major light-harvesting complex at 2.72 Å resolution. *Nature* 428, 287–292. doi: 10.1038/nature02373
- Lohscheider, J. N., Rojas-Stutz, M. C., Rothbart, M., Andersson, U., Funck, D., Mendgen, K., et al. (2015). Altered levels of LIL3 isoforms in *Arabidopsis* lead to disturbed pigment-protein assembly and chlorophyll synthesis, chlorotic phenotype and impaired photosynthetic performance. *Plant Cell Environ.* 38, 2115–2127. doi: 10.1111/pce.12540
- Lu, Y. (2011). The occurrence of a thylakoid-localized small zinc finger protein in land plants. *Plant Signal. Behav.* 6, 1181–1185. doi: 10.4161/psb.6.12.18022
- Lu, Y., Hall, D. A., and Last, R. L. (2011). A small zinc finger thylakoid protein plays a role in maintenance of Photosystem II in *Arabidopsis thaliana*. *Plant Cell* 23, 1861–1875. doi: 10.1105/tpc.111.085456
- Lu, Y., Wang, H.-R., Li, H., Cui, H.-R., Feng, Y.-G., and Wang, X.-Y. (2013). A chloroplast membrane protein LTO1/AtVKOR involving in redox regulation and ROS homeostasis. *Plant Cell Rep.* 32, 1427–1440. doi: 10.1007/s00299-013-1455-9
- Luciński, R., Misztal, L., Samardakiewicz, S., and Jackowski, G. (2011a). Involvement of Deg5 protease in wounding-related disposal of PsbF apoprotein. *Plant Physiol. Biochem.* 49, 311–320. doi: 10.1016/j.plaphy.2011.01.001
- Luciński, R., Misztal, L., Samardakiewicz, S., and Jackowski, G. (2011b). The thylakoid protease Deg2 is involved in stress-related degradation of the Photosystem II light-harvesting protein Lhcb6 in *Arabidopsis thaliana*. *New Phytol.* 192, 74–86. doi: 10.1111/j.1469-8137.2011.03782.x

- Ma, J., Peng, L., Guo, J., Lu, Q., Lu, C., and Zhang, L. (2007). LPA2 is required for efficient assembly of Photosystem II in *Arabidopsis thaliana*. *Plant Cell* 19, 1980–1993. doi: 10.1105/tpc.107.050526
- Mabbitt, P. D., Wilbanks, S. M., and Eaton-Rye, J. J. (2014). Structure and function of the hydrophilic Photosystem II assembly proteins: Psb27, Psb28 and Ycf48. *Plant Physiol. Biochem.* 81, 96–107. doi: 10.1016/j.plaphy.2014.02.013
- Mann, N. H., Novac, N., Mullineaux, C. W., Newman, J., Bailey, S., and Robinson, C. (2000). Involvement of an FtsH homologue in the assembly of functional Photosystem I in the cyanobacterium *Synechocystis* sp. PCC 6803. *FEBS Lett.* 479, 72–77. doi: 10.1016/S0014-5793(00)01871-8
- Meurer, J., Plücker, H., Kowallik, K. V., and Westhoff, P. (1998). A nuclear-encoded protein of prokaryotic origin is essential for the stability of Photosystem II in *Arabidopsis thaliana*. *EMBO J.* 17, 5286–5297. doi: 10.1093/emboj/17.18.5286
- Michoux, F., Boehm, M., Bialek, W., Takasaka, K., Maghlaoui, K., Barber, J., et al. (2014). Crystal structure of CyanoQ from the thermophilic cyanobacterium *Thermosynechococcus elongatus* and detection in isolated Photosystem II complexes. *Photosynth. Res.* 122, 57–67. doi: 10.1007/s11120-014-0010-z
- Michoux, F., Takasaka, K., Boehm, M., Nixon, P. J., and Murray, J. W. (2010). Structure of CyanoP at 2.8 Å: implications for the evolution and function of the PsbP subunit of Photosystem II. *Biochemistry* 49, 7411–7413. doi: 10.1021/bi1011145
- Mork-Jansson, A. E., Gargano, D., Kmiec, K., Furnes, C., Shevela, D., and Eichacker, L. A. (2015). Lil3 dimerization and chlorophyll binding in *Arabidopsis thaliana*. *FEBS Lett.* 589, 3064–3070. doi: 10.1016/j.febslet.2015.08.023
- Morré, D. J., Sellén, G., Sundqvist, C., and Sandelius, A. S. (1991). Stromal low temperature compartment derived from the inner membrane of the chloroplast envelope. *Plant Physiol.* 97, 1558–1564. doi: 10.1104/pp.97.4.1558
- Mulo, P., Sirpiö, S., Suorsa, M., and Aro, E. M. (2008). Auxiliary proteins involved in the assembly and sustenance of Photosystem II. *Photosynth. Res.* 98, 489–501. doi: 10.1007/s11120-008-9320-3
- Muranaka, A., Watanabe, S., Sakamoto, A., and Shimada, H. (2012). *Arabidopsis* cotyledon chloroplast biogenesis factor CYO1 uses glutathione as an electron donor and interacts with PSI (A1 and A2) and PSII (CP43 and CP47) subunits. *J. Plant Physiol.* 169, 1212–1215. doi: 10.1016/j.jplph.2012.04.001
- Nagy, E., Henics, T., Eckert, M., Miseta, A., Lightowers, R. N., and Kellermayer, M. (2000). Identification of the NAD(+)–binding fold of glyceraldehyde-3-phosphate dehydrogenase as a novel RNA-binding domain. *Biochem. Biophys. Res. Commun.* 275, 253–260. doi: 10.1006/bbrc.2000.3246
- Nath, K., Poudyal, R. S., Eom, J.-S., Park, Y. S., Zulfugarov, I. S., Mishra, S. R., et al. (2013). Loss-of-function of OsSTN8 suppresses the Photosystem II core protein phosphorylation and interferes with the Photosystem II repair mechanism in rice (*Oryza sativa*). *Plant J.* 76, 675–686. doi: 10.1111/tj.12331
- Nickelsen, J., and Rengstl, B. (2013). Photosystem II assembly: from cyanobacteria to plants. *Annu. Rev. Plant Biol.* 64, 609–635. doi: 10.1146/annurev-arplant-050312-120124
- Nickelsen, J., and Zerges, W. (2013). Thylakoid biogenesis has joined the new era of bacterial cell biology. *Front. Plant Sci.* 4:458. doi: 10.3389/fpls.2013.00458
- Nield, J., and Barber, J. (2006). Refinement of the structural model for the Photosystem II supercomplex of higher plants. *Biochim. Biophys. Acta* 1757, 353–361. doi: 10.1016/j.bbabi.2006.03.019
- Niwa, H., Tsuchiya, D., Makyio, H., Yoshida, M., and Morikawa, K. (2002). Hexameric ring structure of the ATPase domain of the membrane-integrated metalloprotease FtsH from *Thermus thermophilus* HB8. *Structure* 10, 1415–1423. doi: 10.1016/S0969-2126(02)00855-9
- Nixon, P. J., Barker, M., Boehm, M., de Vries, R., and Komenda, J. (2005). FtsH-mediated repair of the Photosystem II complex in response to light stress. *J. Exp. Bot.* 56, 357–363. doi: 10.1093/jxb/eri021
- Nixon, P. J., Michoux, F., Yu, J., Boehm, M., and Komenda, J. (2010). Recent advances in understanding the assembly and repair of Photosystem II. *Ann. Bot.* 106, 1–16. doi: 10.1093/aob/mcq059
- Nixon, P. J., Trost, J. T., and Diner, B. A. (1992). Role of the carboxy terminus of polypeptide D1 in the assembly of a functional water-oxidizing manganese cluster in Photosystem II of the cyanobacterium *Synechocystis* sp. PCC 6803: assembly requires a free carboxyl group at C-terminal position 344. *Biochemistry* 31, 10859–10871. doi: 10.1021/bi00159a029
- Nowaczyk, M. M., Hebel, R., Schlodder, E., Meyer, H. E., Warscheid, B., and Rögnér, M. (2006). Psb27, a cyanobacterial lipoprotein, is involved in the repair cycle of Photosystem II. *Plant Cell* 18, 3121–3131. doi: 10.1105/tpc.106.042671
- Ossenbühl, F., Göhre, V., Meurer, J., Krieger-Liszskay, A., Roicha, J.-D., and Eichacker, L. A. (2004). Efficient assembly of Photosystem II in *Chlamydomonas reinhardtii* requires Alb3.1p, a homolog of *Arabidopsis* ALBINO3. *Plant Cell* 16, 1790–1800. doi: 10.1105/tpc.023226
- Park, S. W., Li, W., Viehhauser, A., He, B., Kim, S., Nilsson, A. K., et al. (2013). Cyclophilin 20-3 relays a 12-oxo-phytyldienoic acid signal during stress responsive regulation of cellular redox homeostasis. *Proc. Natl. Acad. Sci. U.S.A.* 110, 9559–9564. doi: 10.1073/pnas.1218872110
- Pasch, J. C., Nickelsen, J., and Schünemann, D. (2005). The yeast split-ubiquitin system to study chloroplast membrane protein interactions. *Appl. Microbiol. Biotechnol.* 69, 440–447. doi: 10.1007/s00253-005-0029-3
- Peng, L. W., Ma, J. F., Chi, W., Guo, J. K., Zhu, S. Y., Lu, Q. T., et al. (2006). LOW PSII ACCUMULATION1 is involved in efficient assembly of Photosystem II in *Arabidopsis thaliana*. *Plant Cell* 18, 955–969. doi: 10.1105/tpc.105.037689
- Persson, B., Kallberg, Y., Bray, J. E., Bruford, E., Dellaporta, S. L., Favia, A. D., et al. (2009). The SDR (short-chain dehydrogenase/reductase and related enzymes) nomenclature initiative. *Chem. Biol. Interact.* 178, 94–98. doi: 10.1016/j.cbi.2008.10.040
- Pesaresi, P., Pribil, M., Wunder, T., and Leister, D. (2011). Dynamics of reversible protein phosphorylation in thylakoids of flowering plants: the roles of STN7, STN8 and TAP38. *Biochim. Biophys. Acta* 1807, 887–896. doi: 10.1016/j.bbabi.2010.08.002
- Pioli, P. A., Hamilton, B. J., Connolly, J. E., Brewer, G., and Rigby, W. F. (2002). Lactate dehydrogenase is an AU-rich element-binding protein that directly interacts with AUF1. *J. Biol. Chem.* 277, 35738–35745. doi: 10.1074/jbc.M204002200
- Plücker, H., Müller, B., Grohmann, D., Westhoff, P., and Eichacker, L. A. (2002). The HCF136 protein is essential for assembly of the Photosystem II reaction center in *Arabidopsis thaliana*. *FEBS Lett.* 532, 85–90. doi: 10.1016/S0014-5793(02)03634-7
- Pribil, M., Pesaresi, P., Hertle, A., Barbato, R., and Leister, D. (2010). Role of plastid protein phosphatase TAP38 in LHCI dephosphorylation and thylakoid electron flow. *PLoS Biol.* 8:e1000288. doi: 10.1371/journal.pbio.1000288
- Rast, A., Heinz, S., and Nickelsen, J. (2015). Biogenesis of thylakoid membranes. *Biochim. Biophys. Acta* 1847, 821–830. doi: 10.1016/j.bbabi.2015.01.007
- Rintamaki, E., Salonen, M., Suoranta, U. M., Carlberg, I., Andersson, B., and Aro, E. M. (1997). Phosphorylation of light-harvesting complex II and Photosystem II core proteins shows different irradiance-dependent regulation *in vivo*: application of phosphothreonine antibodies to analysis of thylakoid phosphoproteins. *J. Biol. Chem.* 272, 30476–30482. doi: 10.1074/jbc.272.48.30476
- Rokka, A., Aro, E.-M., Herrmann, R. G., Andersson, B., and Vener, A. V. (2000). Dephosphorylation of Photosystem II reaction center proteins in plant photosynthetic membranes as an immediate response to abrupt elevation of temperature. *Plant Physiol.* 123, 1525–1536. doi: 10.1104/pp.123.4.1525
- Rokka, A., Suorsa, M., Saleem, A., Battchikova, N., and Aro, E. M. (2005). Synthesis and assembly of thylakoid protein complexes: multiple assembly steps of Photosystem II. *Biochem. J.* 388, 159–168. doi: 10.1042/BJ20042098
- Romano, P. G., Horton, P., and Gray, J. E. (2004). The *Arabidopsis* cyclophilin gene family. *Plant Physiol.* 134, 1268–1282. doi: 10.1104/pp.103.022160
- Rossini, S., Casazza, A. P., Engelmann, E. C. M., Havaux, M., Jennings, R. C., and Soave, C. (2006). Suppression of both ELIP1 and ELIP2 in *Arabidopsis* does not affect tolerance to photoinhibition and photooxidative stress. *Plant Physiol.* 141, 1264–1273. doi: 10.1104/pp.106.083055
- Sakamoto, W., Tamura, T., Hanba-Tomita, Y., Sodmergen, and Murata, M. (2002). The *VARI1* locus of *Arabidopsis* encodes a chloroplastic FtsH and is responsible for leaf variegation in the mutant alleles. *Genes Cells* 7, 769–780. doi: 10.1046/j.1365-2443.2002.00558.x
- Sakamoto, W., Zaltsman, A., Adam, Z., and Takahashi, Y. (2003). Coordinated regulation and complex formation of YELLOW VARIEGATED1 and YELLOW VARIEGATED2, chloroplastic FtsH metalloproteases involved in the repair cycle of Photosystem II in *Arabidopsis* thylakoid membranes. *Plant Cell* 15, 2843–2855. doi: 10.1105/tpc.017319
- Samol, I., Shapiguzov, A., Ingelsson, B., Fucile, G., Crèvecoeur, M., Vener, A. V., et al. (2012). Identification of a Photosystem II phosphatase

- involved in light acclimation in *Arabidopsis*. *Plant Cell* 24, 2596–2609. doi: 10.1105/tpc.112.095703
- Schneider, A., Steinberger, I., Strissel, H., Kunz, H.-H., Manavski, N., Meurer, J., et al. (2014). The *Arabidopsis* Tellurite resistance C protein together with ALB3 is involved in Photosystem II protein synthesis. *Plant J.* 78, 344–356. doi: 10.1111/tpj.12474
- Schuhmann, H., and Adamska, I. (2012). Deg proteases and their role in protein quality control and processing in different subcellular compartments of the plant cell. *Physiol. Plant.* 145, 224–234. doi: 10.1111/j.1399-3054.2011.01533.x
- Schuhmann, H., Mogg, U., and Adamska, I. (2011). A new principle of oligomerization of plant DEG7 protease based on interactions of degenerated protease domains. *Biochem. J.* 435, 167–174. doi: 10.1042/BJ20101613
- Schult, K., Meierhoff, K., Paradies, S., Toller, T., Wolff, P., and Westhoff, P. (2007). The nuclear-encoded factor HCF173 is involved in the initiation of translation of the *psbA* mRNA in *Arabidopsis thaliana*. *Plant Cell* 19, 1329–1346. doi: 10.1105/tpc.106.042895
- Schunemann, D. (2007). Mechanisms of protein import into thylakoids of chloroplasts. *Biol. Chem.* 388, 907–915. doi: 10.1515/BC.2007.111
- Shapiguzov, A., Ingelsson, B., Samol, I., Andres, C., Kessler, F., Rochaix, J. D., et al. (2010). The PPH1 phosphatase is specifically involved in LHClI dephosphorylation and state transitions in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 107, 4782–4787. doi: 10.1073/pnas.0913810107
- Shen, G., Zhao, J., Reimer, S. K., Antonkine, M. L., Cai, Q., Weiland, S. M., et al. (2002). Assembly of Photosystem I: I. Inactivation of the *rubA* gene encoding a membrane-associated rubredoxin in the cyanobacterium *Synechococcus* sp. PCC 7002 causes a loss of Photosystem I activity. *J. Biol. Chem.* 277, 20343–20354. doi: 10.1074/jbc.M201103200
- Shi, L.-X., Hall, M., Funk, C., and Schröder, W. P. (2012). Photosystem II, a growing complex: updates on newly discovered components and low molecular mass proteins. *Biochim. Biophys. Acta* 1817, 13–25. doi: 10.1016/j.bbabi.2011.08.008
- Shimada, H., Mochizuki, M., Ogura, K., Froehlich, J. E., Osteryoung, K. W., Shirano, Y., et al. (2007). *Arabidopsis* cotyledon-specific chloroplast biogenesis factor CYO1 is a protein disulfide isomerase. *Plant Cell* 19, 3157–3169. doi: 10.1105/tpc.107.051714
- Silva, P., Thompson, E., Bailey, S., Kruse, O., Mullineaux, C. W., Robinson, C., et al. (2003). FtsH is involved in the early stages of repair of Photosystem II in *Synechocystis* sp. PCC 6803. *Plant Cell* 15, 2152–2164. doi: 10.1105/tpc.012609
- Sirpiö, S., Allahverdiyeva, Y., Suorsa, M., Paakkarinen, V., Vainonen, J., Battchikova, N., et al. (2007). TLP18.3, a novel thylakoid lumen protein regulating Photosystem II repair cycle. *Biochem. J.* 406, 415–425. doi: 10.1042/BJ20070460
- Sirpiö, S., Khrouchtchova, A., Allahverdiyeva, Y., Hansson, M., Fristedt, R., Vener, A. V., et al. (2008). AtCYP38 ensures early biogenesis, correct assembly and sustenance of Photosystem II. *Plant J.* 55, 639–651. doi: 10.1111/j.1365-313X.2008.03532.x
- Speiser, A., Haberland, S., Watanabe, M., Wirtz, M., Dietz, K. J., Saito, K., et al. (2015). The significance of cysteine synthesis for acclimation to high light conditions. *Front. Plant Sci.* 5:776. doi: 10.3389/fpls.2014.00776
- Spetea, C., Hundal, T., Lohmann, F., and Andersson, B. (1999). GTP bound to chloroplast thylakoid membranes is required for light-induced, multienzyme degradation of the Photosystem II D1 protein. *Proc. Natl. Acad. Sci. U.S.A.* 96, 6547–6552. doi: 10.1073/pnas.96.11.6547
- Staleva, H., Komenda, J., Shukla, M. K., Slouf, V., Kana, R., Polivka, T., et al. (2015). Mechanism of photoprotection in the cyanobacterial ancestor of plant antenna proteins. *Nat. Chem. Biol.* 11, 287–291. doi: 10.1038/nchembio.1755
- Suga, M., Akita, F., Hirata, K., Ueno, G., Murakami, H., Nakajima, Y., et al. (2015). Native structure of Photosystem II at 1.95 Å resolution viewed by femtosecond X-ray pulses. *Nature* 517, 99–103. doi: 10.1038/nature13991
- Summerfield, T., Winter, R., and Eaton-Rye, J. (2005). Investigation of a requirement for the PsbP-like protein in *Synechocystis* sp. PCC 6803. *Photosynth. Res.* 84, 263–268. doi: 10.1007/s11120-004-6431-3
- Sun, X., Fu, T., Chen, N., Guo, J., Ma, J., Zou, M., et al. (2010a). The stromal chloroplast Deg7 protease participates in the repair of Photosystem II after photoinhibition in *Arabidopsis*. *Plant Physiol.* 152, 1263–1273. doi: 10.1104/pp.109.150722
- Sun, X., Ouyang, M., Guo, J., Ma, J., Lu, C., Adam, Z., et al. (2010b). The thylakoid protease Deg1 is involved in Photosystem-II assembly in *Arabidopsis thaliana*. *Plant J.* 62, 240–249. doi: 10.1111/j.1365-313X.2010.04140.x
- Sun, X., Peng, L., Guo, J., Chi, W., Ma, J., Lu, C., et al. (2007). Formation of DEG5 and DEG8 complexes and their involvement in the degradation of photodamaged Photosystem II reaction center D1 protein in *Arabidopsis*. *Plant Cell* 19, 1347–1361. doi: 10.1105/tpc.106.049510
- Takahashi, K., Takabayashi, A., Tanaka, A., and Tanaka, R. (2014). Functional analysis of light-harvesting-like protein 3 (LIL3) and its light-harvesting chlorophyll-binding motif in *Arabidopsis*. *J. Biol. Chem.* 289, 987–999. doi: 10.1074/jbc.M113.525428
- Takechi, K., Sodmergen, Murata, M., Motoyoshi, F., and Sakamoto, W. (2000). The *YELLOW VARIEGATED (VAR2)* locus encodes a homologue of FtsH, an ATP-dependent protease in *Arabidopsis*. *Plant Cell Physiol.* 41, 1334–1346. doi: 10.1093/pcp/pcd067
- Tanaka, R., Rothbart, M., Oka, S., Takabayashi, A., Takahashi, K., Shibata, M., et al. (2010). LIL3, a light-harvesting-like protein, plays an essential role in chlorophyll and tocopherol biosynthesis. *Proc. Natl. Acad. Sci. U.S.A.* 107, 16721–16725. doi: 10.1073/pnas.1004699107
- Tanz, S. K., Kilian, J., Johnsson, C., Apel, K., Small, I., Harter, K., et al. (2012). The SCO2 protein disulphide isomerase is required for thylakoid biogenesis and interacts with LHCB1 chlorophyll a/b binding proteins which affects chlorophyll biosynthesis in *Arabidopsis* seedlings. *Plant J.* 69, 743–754. doi: 10.1111/j.1365-313X.2011.04833.x
- Thornton, L. E., Ohkawa, H., Roose, J. L., Kashino, Y., Keren, N., and Pakrasi, H. B. (2004). Homologs of plant PsbP and PsbQ proteins are necessary for regulation of Photosystem II activity in the cyanobacterium *Synechocystis* 6803. *Plant Cell* 16, 2164–2175. doi: 10.1105/tpc.104.023515
- Tikkanen, M., and Aro, E. M. (2012). Thylakoid protein phosphorylation in dynamic regulation of Photosystem II in higher plants. *Biochim. Biophys. Acta* 1817, 232–238. doi: 10.1016/j.bbabi.2011.05.005
- Tikkanen, M., Nurmi, M., Kangasjärvi, S., and Aro, E. M. (2008). Core protein phosphorylation facilitates the repair of photodamaged Photosystem II at high light. *Biochim. Biophys. Acta* 1777, 1432–1437. doi: 10.1016/j.bbabi.2008.08.004
- Torabi, S., Umate, P., Manavski, N., Plöching, M., Kleinknecht, L., Bogireddi, H., et al. (2014). PsbN is required for assembly of the Photosystem II reaction center in *Nicotiana tabacum*. *Plant Cell* 26, 1183–1199. doi: 10.1105/tpc.113.120444
- Umeha, Y., Kawakami, K., Shen, J. R., and Kamiya, N. (2011). Crystal structure of oxygen-evolving Photosystem II at a resolution of 1.9 Å. *Nature* 473, 55–U65. doi: 10.1038/nature09913
- Uniacke, J., and Zerges, W. (2007). Photosystem II assembly and repair are differentially localized in *Chlamydomonas*. *Plant Cell* 19, 3640–3654. doi: 10.1105/tpc.107.054882
- Urantowka, A., Knorpp, C., Olczak, T., Kolodziejczak, M., and Janska, H. (2005). Plant mitochondria contain at least two *i*-AAA-like complexes. *Plant Mol. Biol.* 59, 239–252. doi: 10.1007/s11103-005-8766-3
- Vasudevan, D., Fu, A., Luan, S., and Swaminathan, K. (2012). Crystal structure of *Arabidopsis* cyclophilin38 reveals a previously uncharacterized immunophilin fold and a possible autoinhibitory mechanism. *Plant Cell* 24, 2666–2674. doi: 10.1105/tpc.111.093781
- Vener, A. V., Harms, A., Sussman, M. R., and Vierstra, R. D. (2001). Mass spectrometric resolution of reversible protein phosphorylation in photosynthetic membranes of *Arabidopsis thaliana*. *J. Biol. Chem.* 276, 6959–6966. doi: 10.1074/jbc.M009394200
- Vener, A. V., Rokka, A., Fulgosi, H., Andersson, B., and Herrmann, R. G. (1999). A cyclophilin-regulated PP2A-like protein phosphatase in thylakoid membranes of plant chloroplasts. *Biochemistry* 38, 14955–14965. doi: 10.1021/bi990971v
- Wagner, R., Aigner, H., Pruzinska, A., Jankapaa, H. J., Jansson, S., and Funk, C. (2011). Fitness analyses of *Arabidopsis thaliana* mutants depleted of FtsH metalloproteases and characterization of three FtsH6 deletion mutants exposed to high light stress, senescence and chilling. *New Phytol.* 191, 449–458. doi: 10.1111/j.1469-8137.2011.03684.x
- Walter, B., Hristou, A., Nowaczyk, M. M., and Schunemann, D. (2015). *In vitro* reconstitution of co-translational D1 insertion reveals a role of the cpSec-Alb3 translocase and Vipp1 in Photosystem II biogenesis. *Biochem. J.* 468, 315–324. doi: 10.1042/BJ20141425
- Wang, P., Liu, J., Liu, B., Feng, D., Da, Q., Shu, S., et al. (2013). Evidence for a role of chloroplastic m-type thioredoxins in the biogenesis of Photosystem

- II in *Arabidopsis*. *Plant Physiol.* 163, 1710–1728. doi: 10.1104/pp.113.228353
- Wang, Q., Sullivan, R. W., Kight, A., Henry, R. L., Huang, J., Jones, A. M., et al. (2004). Deletion of the chloroplast-localized *Thylakoid formation1* gene product in *Arabidopsis* leads to deficient thylakoid formation and variegated leaves. *Plant Physiol.* 136, 3594–3604. doi: 10.1104/pp.104.049841
- Wei, L., Guo, J., Ouyang, M., Sun, X., Ma, J., Chi, W., et al. (2010). LPA19, a Psb27 homolog in *Arabidopsis thaliana*, facilitates D1 protein precursor processing during PSII biogenesis. *J. Biol. Chem.* 285, 21391–21398. doi: 10.1074/jbc.M110.105064
- Westphal, S., Soll, J., and Vothknecht, U. C. (2001). A vesicle transport system inside chloroplasts. *FEBS Lett.* 506, 257–261. doi: 10.1016/S0014-5793(01)02931-3
- Westphal, S., Soll, J., and Vothknecht, U. C. (2003). Evolution of chloroplast vesicle transport. *Plant Cell Physiol.* 44, 217–222. doi: 10.1093/pcp/pcg023
- Wittenberg, G., Levitan, A., Klein, T., Dangoor, I., Keren, N., and Danon, A. (2014). Knockdown of the *Arabidopsis thaliana* chloroplast protein disulfide isomerase 6 results in reduced levels of photoinhibition and increased D1 synthesis in high light. *Plant J.* 78, 1003–1013. doi: 10.1111/tpj.12525
- Wu, H. Y., Liu, M. S., Lin, T. P., and Cheng, Y. S. (2011). Structural and functional assays of AtTLP18.3 identify its novel acid phosphatase activity in thylakoid lumen. *Plant Physiol.* 157, 1015–1025. doi: 10.1104/pp.111.184739
- Yamamoto, Y., Inagaki, N., and Satoh, K. (2001). Overexpression and characterization of carboxyl-terminal processing protease for precursor D1 protein: regulation of enzyme-substrate interaction by molecular environments. *J. Biol. Chem.* 276, 7518–7525. doi: 10.1074/jbc.M008877200
- Yamatani, H., Sato, Y., Masuda, Y., Kato, Y., Morita, R., Fukunaga, K., et al. (2013). NYC4, the rice ortholog of *Arabidopsis THF1*, is involved in the degradation of chlorophyll – protein complexes during leaf senescence. *Plant J.* 74, 652–662. doi: 10.1111/tpj.12154
- Yin, S., Sun, X., and Zhang, L. (2008). An *Arabidopsis ctpA* homologue is involved in the repair of Photosystem II under high light. *Chin. Sci. Bull.* 53, 1021–1026. doi: 10.1007/s11434-008-0153-4
- Yu, F., Park, S., and Rodermel, S. R. (2004). The *Arabidopsis* FtsH metalloprotease gene family: interchangeability of subunits in chloroplast oligomeric complexes. *Plant J.* 37, 864–876. doi: 10.1111/j.1365-313X.2003.02014.x
- Yu, F., Park, S., and Rodermel, S. R. (2005). Functional redundancy of AtFtsH metalloproteases in thylakoid membrane complexes. *Plant Physiol.* 138, 1957–1966. doi: 10.1104/pp.105.061234
- Zaltsman, A., Feder, A., and Adam, Z. (2005a). Developmental and light effects on the accumulation of FtsH protease in *Arabidopsis* chloroplasts – implications for thylakoid formation and Photosystem II maintenance. *Plant J.* 42, 609–617. doi: 10.1111/j.1365-313X.2005.02401.x
- Zaltsman, A., Ori, N., and Adam, Z. (2005b). Two types of FtsH protease subunits are required for chloroplast biogenesis and Photosystem II repair in *Arabidopsis*. *Plant Cell* 17, 2782–2790. doi: 10.1105/tpc.105.035071
- Zelisko, A., Garcia-Lorenzo, M., Jackowski, G., Jansson, S., and Funk, C. (2005). AtFtsH6 is involved in the degradation of the light-harvesting complex II during high-light acclimation and senescence. *Proc. Natl. Acad. Sci. U.S.A.* 102, 13699–13704. doi: 10.1073/pnas.0503472102
- Zhang, D., Zhou, G., Liu, B., Kong, Y., Chen, N., Qiu, Q., et al. (2011). HCF243 encodes a chloroplast-localized protein involved in the D1 protein stability of the *Arabidopsis* Photosystem II complex. *Plant Physiol.* 157, 608–619. doi: 10.1104/pp.111.183301
- Zhang, L., Kato, Y., Otters, S., Vothknecht, U. C., and Sakamoto, W. (2012). Essential role of VIPP1 in chloroplast envelope maintenance in *Arabidopsis*. *Plant Cell* 24, 3695–3707. doi: 10.1105/tpc.112.103606
- Zhang, L., and Sakamoto, W. (2012). Possible function of VIPP1 in thylakoids: protection but not formation? *Plant Signal. Behav.* 8:e22860. doi: 10.4161/psb.22860
- Zhang, L., Wei, Q., Wu, W., Cheng, Y., Hu, G., Hu, F., et al. (2009). Activation of the heterotrimeric G protein alpha-subunit GPA1 suppresses the ftsH-mediated inhibition of chloroplast development in *Arabidopsis*. *Plant J.* 58, 1041–1053. doi: 10.1111/j.1365-313X.2009.03843.x

Conflict of Interest Statement: The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Lu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.