



# Nitrate transporters in leaves and their potential roles in foliar uptake of nitrogen dioxide<sup>†</sup>

Yanbo Hu<sup>1\*</sup>, Victoria Fernández<sup>2</sup> and Ling Ma<sup>3\*</sup>

<sup>1</sup> College of Life Science, Northeast Forestry University, Harbin, China

<sup>2</sup> Forest Genetics and Ecophysiology Research Group, School of Forest Engineering, Technical University of Madrid, Madrid, Spain

<sup>3</sup> School of Forestry, Northeast Forestry University, Harbin, China

## Edited by:

Ebrahim Hadavi, Islamic Azad University, Iran

## Reviewed by:

Abdullahil Baque, Sher-e-Bangla Agricultural University, Bangladesh  
Brian Grout, University of Copenhagen, Denmark

## \*Correspondence:

Yanbo Hu and Ling Ma, Northeast Forestry University, 26 Hexing Road, Xiangfang District, Harbin 150040, P. R. China  
e-mail: [huybnfu@yahoo.com](mailto:huybnfu@yahoo.com)

<sup>†</sup> This paper was dedicated to Professor Hartmut K. Lichtenthaler on the occasion of his 80th birthday.

While plant roots are specialized organs for the uptake and transport of water and nutrients, the absorption of gaseous or liquid mineral elements by aerial plant parts has been recognized since more than one century. Nitrogen (N) is an essential macronutrient which generally absorbed either as nitrate ( $\text{NO}_3^-$ ) or ammonium ( $\text{NH}_4^+$ ) by plant roots. Gaseous nitrogen pollutants like N dioxide ( $\text{NO}_2$ ) can also be absorbed by plant surfaces and assimilated via the  $\text{NO}_3^-$  assimilation pathway. The subsequent  $\text{NO}_3^-$  flux may induce or repress the expression of various  $\text{NO}_3^-$ -responsive genes encoding for instance, the transmembrane transporters,  $\text{NO}_3^-/\text{NO}_2^-$  (nitrite) reductase, or assimilatory enzymes involved in N metabolism. Based on the existing information, the aim of this review was to theoretically analyze the potential link between foliar  $\text{NO}_2$  absorption and N transport and metabolism. For such purpose, an overview of the state of knowledge on the  $\text{NO}_3^-$  transporter genes identified in leaves or shoots of various species and their roles for  $\text{NO}_3^-$  transport across the tonoplast and plasma membrane, in addition to the process of phloem loading is briefly provided. It is assumed that a  $\text{NO}_2$ -induced accumulation of  $\text{NO}_3^-/\text{NO}_2^-$  may alter the expression of such genes, hence linking transmembrane  $\text{NO}_3^-$  transporters and foliar uptake of  $\text{NO}_2$ . It is likely that *NRT1/NRT2* gene expression and species-dependent apoplastic buffer capacity may be also related to the species-specific foliar  $\text{NO}_2$  uptake process. It is concluded that further work focusing on the expression of *NRT1* (*NRT1.1*, *NRT1.7*, *NRT1.11*, and *NRT1.12*), *NRT2* (*NRT2.1*, *NRT2.4*, and *NRT2.5*) and chloride channel family genes (*CLCa* and *CLCd*) may help us elucidate the physiological and metabolic response of plants fumigated with  $\text{NO}_2$ .

**Keywords:** chloride channel gene, nitrate reductase, nitrate transporter, nitrogen dioxide, signal transmission

## INTRODUCTION

Nitrate ( $\text{NO}_3^-$ ) is the most common form of nitrogen used by plants for growth and development (Bertoni, 2012). Despite the major role of plant roots is absorbing and transporting water and mineral elements, there is abundant evidence showing that nutrients can also be taken up by aerial plant parts (e.g., leaves, fruits and stems) (Eichert and Fernández, 2012; Fernández and Brown, 2013). Foliar applied  $\text{NO}_3^-$  may be absorbed and assimilated efficiently as shown in several studies carried out with different plant species (e.g., Stiegler et al., 2011; Uscola et al., 2014). Gaseous air pollutants like nitrogen dioxide ( $\text{NO}_2$ ) can also be deposited into plant leaves and be taken up mainly through stomata (Eichert and Fernández, 2012).  $\text{NO}_2$  molecules may dissolve in the aqueous phase of the apoplastic space being consequently transformed into nitrate ( $\text{NO}_3^-$ ) and/or nitrite ( $\text{NO}_2^-$ ) by chemical reactions. Thereafter,  $\text{NO}_2$ -derived  $\text{NO}_3^-$  may be transported across the plasma membrane by  $\text{NO}_3^-$  transporters and reach the cytoplasm for further incorporation into cellular N-compounds and/or storage in vacuoles (Hawkesford et al., 2012). The  $\text{NO}_3^-$  stored in the vacuoles may be exported to compensate for the consumption of  $\text{NO}_3^-$  in the metabolic pool (De Angeli et al., 2006),

which suggests that vacuolar  $\text{NO}_3^-$  may largely serve as N buffer for transport processes (Hawkesford et al., 2012).  $\text{NO}_3^-$ /proton antiporters, which may be encoded by chloride channel family (CLC) genes, are responsible for  $\text{NO}_3^-$  influx into plant vacuoles (Geelen et al., 2000).  $\text{NO}_3^-$  transmembrane transporters may be expected to play a role after the uptake of exogenous  $\text{NO}_2$  or  $\text{NO}_3^-$  by the foliage. Several  $\text{NO}_3^-$  transporters identified in leaves have been demonstrated to be closely correlated with e.g., stomatal opening (Guo et al., 2003),  $\text{NO}_3^-$  reductase activity (Loqué et al., 2003), accumulation and remobilization of  $\text{NO}_3^-$  (De Angeli et al., 2006; Fan et al., 2009; Lv et al., 2009). Thereby, such physiological processes may significantly influence and also be affected by the foliar uptake of  $\text{NO}_2$  or  $\text{NO}_3^-$ .

Recently some  $\text{NO}_3^-$  transporter genes were detected in leaves including several members of plant *NRT1* family genes (e.g., *AtNRT1.1*, *AtNRT1.4*, *AtNRT1.7*, *AtNRT1.11*, and *AtNRT1.12*), *NRT2* family genes (e.g., *AtNRT2.1*, *AtNRT2.3*, *AtNRT2.4*, *AtNRT2.5*, *AtNRT2.6*, *AtNRT2.7*, *NpNRT2.1* and *ZmNrt2.1*), and CLC family genes such as *AtCLCa* and *AtCLCd* (Orsel et al., 2002; Guo et al., 2003; Chopin et al., 2007; Fan et al., 2009; Hsu and Tsay, 2013) (**Figure 1**). These genes show various expression levels



Previous studies on foliar uptake of NO<sub>2</sub> mainly focused on the deposition pathways, metabolic processes associated with NO<sub>2</sub>-derived NO<sub>3</sub><sup>-</sup> (Hu, 2011; Hu et al., 2014), and downstream products of NO<sub>2</sub>-N assimilation (Nussbaum et al., 1993; Weber et al., 1995). The current state of knowledge on the potential plant responses to NO<sub>2</sub> exposure is summarized in **Table 1**. However, the relationship between NO<sub>3</sub><sup>-</sup> transmembrane transporters and foliar NO<sub>2</sub> uptake has received only limited scientific attention so far. Foliar uptake of NO<sub>2</sub> seems to be species-specific and concentration-dependent (Hu and Sun, 2010). Expression of the genes encoding leaf NO<sub>3</sub><sup>-</sup> transporters also appears to be species-specific (Ono et al., 2000; Orsel et al., 2002). The contribution of various expression patterns of transporter genes to species-specific NO<sub>2</sub> uptake is currently unknown. Given the N transport mechanisms described above, a potential relationship between foliar NO<sub>2</sub> uptake (substomatal build-up of NO<sub>2</sub> and the subsequent reduction, storage, and remobilization of NO<sub>3</sub><sup>-</sup>) and NO<sub>3</sub><sup>-</sup>-responsive genes encoding the transmembrane transporters and NO<sub>3</sub><sup>-</sup>/NO<sub>2</sub><sup>-</sup> reductases may be hypothesized. For validating such hypothesis, future work focusing on the relationship between organ-specific expression of NRT1/NRT2 genes and species-specific NO<sub>2</sub> uptake should be carried out.

### SUBSTOMATAL BUILD-UP OF NO<sub>2</sub> MAY DISTURB APOPLASTIC pH AND NO<sub>3</sub><sup>-</sup> TRANSPORTERS

The apoplast is defined as the area within the plant tissues which is beyond the cell plasma membrane, and includes the cell wall, middle lamella, xylem and gas and water filled intercellular spaces (Sattelmacher, 2001). The leaf apoplastic space plays a role in ion exchange and as a diffusion barrier (Sattelmacher, 2001). Estimates of the volume of leaf water in the apoplast vary from 10 to 35% of total leaf water (Speer and Kaiser, 1991; Wardlaw, 2005). Dissolution of NO<sub>2</sub> in the apoplast may produce H<sup>+</sup> and NO<sub>3</sub><sup>-</sup>/NO<sub>2</sub><sup>-</sup>. Foliar NO<sub>2</sub> uptake is calculated to yield at most 0.22 mol excess H<sup>+</sup> per mol N (Raven, 1988). Therefore, a build-up of NO<sub>2</sub> in the leaf substomatal cavities may lead to apoplastic pH disturbances. Among other factors, the resulting apoplastic pH changes will depend on NO<sub>2</sub> concentration, root N supply and plant N status. The supply of NO<sub>3</sub><sup>-</sup> via the root system significantly increased the leaf apoplastic pH of *Phaseolus vulgaris* and *Helianthus annuus*, whereas the depletion of NO<sub>3</sub><sup>-</sup> in nutrient solution led to lower leaf apoplastic pH values in *Zea mays* (Mühling and Lauchli, 2001). NH<sub>4</sub>NO<sub>3</sub> nutrition did not change the leaf apoplastic pH in sunflower (Kosegarten et al., 1999). Moreover, foliar NH<sub>4</sub><sup>+</sup> fertilization may either lead to apoplastic alkalization (Felle and Hanstein, 2002) or acidification (Mühling and Lauchli, 2001). When supplying NH<sub>4</sub>Cl (1 mM) via the root to soybean plants, low concentrations of NO<sub>2</sub> (0.2–0.25 μL·L<sup>-1</sup>) significantly increased the leaf apoplastic pH (Qiao and Murray, 1997), whereas under a higher root NO<sub>3</sub><sup>-</sup> dose (5 mM), high concentrations of NO<sub>2</sub> (1.1 μL·L<sup>-1</sup>) increased the acidity of the leaves (Qiao and Murray, 1998). Apoplastic pH is an important factor affecting plasmalemma proton pumps (Hoffmann et al., 1992; Sattelmacher, 2001). Apoplastic alkalization or acidification may induce plasma membrane depolarization or hyperpolarization (Hedrich et al., 2001). This may further modulate the deactivation or activation of membrane-bound

proton-transporting enzymes, and the corresponding ion channel regulation for co-transport of anions (Savchenko et al., 2000). Wipfel et al. (2010) found that the fluctuation of apoplastic pH had a regulatory effect on plant sucrose transporters. Based on the above information, it can be reckoned that the apoplastic pH changes caused by NO<sub>2</sub> may repress or induce NO<sub>3</sub><sup>-</sup>-responsive genes encoding the transmembrane transporters.

Some NO<sub>3</sub><sup>-</sup> transporter genes (such as *AtNRT1.1* and *ZmNrt2.1*) in leaves are NO<sub>3</sub><sup>-</sup>-inducible, while others such as *AtNRT2.5* are NO<sub>3</sub><sup>-</sup>-repressible (Okamoto et al., 2003). Low-affinity transporter systems (*NRT1* family) may significantly contribute to NO<sub>3</sub><sup>-</sup> uptake at external NO<sub>3</sub><sup>-</sup> concentrations above 250 μM. However, high-affinity transporters (*NRT2* family) including the constitutive (cHATS Km = 6–20 mM) and inducible HATS (Km = 20–100 mM), are active at low external concentrations of 0–0.5 mM (Crawford and Glass, 1998; Quaggiotti et al., 2003; Hawkesford et al., 2012). When analyzing the *AtNRT1.7* NO<sub>3</sub><sup>-</sup> transporter gene in *Arabidopsis*, Fan et al. (2009) applied 50 mM K<sup>15</sup>NO<sub>3</sub> to carry out measurements on distal parts of the rosette leaf. The <sup>15</sup>N-NO<sub>3</sub><sup>-</sup> tracing assay showed that the percentage of total <sup>15</sup>N in the leaves ranged from 0 to 10% for wild-type plants, and between 5 and 15% for the *nrt1.7* mutants. The percentage of NO<sub>3</sub><sup>-</sup>-<sup>15</sup>N was in the range of the NO<sub>2</sub>-derived reduced N of wild herbaceous plants (from 0.98 to 10.1%) and woody plants (0.15–12.7%) for the 217 taxa fumigated with 4.0 ± 0.1 μmol·mol<sup>-1</sup> NO<sub>2</sub> (Morikawa et al., 1998). Accordingly, the content of NO<sub>2</sub>-derived reduced N ranged from 0.25 to 5.72 mg N·g<sup>-1</sup> dry weight for wild herbaceous plants, and 0.04–6.57 mg N·g<sup>-1</sup> dry weight for woody plants. This comparison suggests that the amounts of NO<sub>2</sub>-derived NO<sub>3</sub><sup>-</sup> in leaves are in the range of the NO<sub>3</sub><sup>-</sup> concentrations which may induce the two types of transporter systems (i.e., high and low affinity).

From the reasoning provided above, it can be reckoned that substomatal build-up of NO<sub>2</sub> may lead to concentration-dependent changes of apoplastic pH and NO<sub>3</sub><sup>-</sup> concentration. Such pH fluctuations may influence NO<sub>3</sub><sup>-</sup> transmembrane transport by the induction or repression of the transporters and transporter gene expression, and may provide some sort of feedback regulation on the uptake of NO<sub>2</sub> by the foliage. For example, apoplastic mesophyll signals have been recently found to induce rapid stomatal responses in *Commelina communis* (Fujita et al., 2013). In response to NO<sub>2</sub> fumigation, multiple physiological and metabolic responses may occur which could either ultimately favor or inhibit the process of symplastic N uptake (**Table 1**). The multi-responses of NO<sub>3</sub><sup>-</sup> transporters to the substomatal build-up of NO<sub>2</sub> may partially contribute to species-specific NO<sub>2</sub> uptake, but future studies with different plant species shall be carried out for clarifying this complex issue.

### NITRATE TRANSPORTERS ARE POSSIBLY INVOLVED IN THE REDUCTION AND ACCUMULATION OF NO<sub>2</sub>-DERIVED NO<sub>3</sub><sup>-</sup>

In leaf cytoplasm, NO<sub>2</sub>-derived NO<sub>3</sub><sup>-</sup> has at least two fates: (i) assimilation into amino acids, and (ii) accumulation in vacuole (Hawkesford et al., 2012). The metabolic pathway will depend on the external NO<sub>3</sub><sup>-</sup> concentration and leaf N demand (Stulen et al., 1998). NO<sub>2</sub>-derived NO<sub>3</sub><sup>-</sup> will be assimilated mainly through the NO<sub>3</sub><sup>-</sup> assimilation pathway (Morikawa et al., 1998). NO<sub>3</sub><sup>-</sup>

**Table 1 | Physiological and metabolic responses of plant organs to nitrogen dioxide (NO<sub>2</sub>) exposure.**

Plant organ	Action site	Physiological function of exogenous NO <sub>2</sub> on plants		References
		Low NO <sub>2</sub> concentration (e.g., 40–60 nL.l <sup>-1</sup> )	High NO <sub>2</sub> concentration (e.g., 1–4 μL.l <sup>-1</sup> )	
Leaf	Stomata	Stimulation on stomatal aperture and stomatal conductance <sup>[1.3]</sup> ; Reduced stomatal density <sup>[1.2]</sup>	Stomatal closure and declined stomatal conductance <sup>[1.1]</sup>	[1.1]Qiao and Murray, 1998; [1.2]Siegwolf et al., 2001; [1.3]Takagi and Gyokusen, 2004
	Apoplast	Increase in the malondialdehyde (MDA) level and superoxide dismutase (SOD) at 0.5 μL.L <sup>-1</sup> NO <sub>2</sub> <sup>[2.1]</sup>	Acidity of apoplast <sup>[1.1]</sup> ; Induced expression of germin-like proteins (RmGLP2) <sup>[2.2]</sup> ; Decline in MDA content and SOD activity <sup>[2.3]</sup> ; Decline in ASA <sup>[2.1]</sup>	[2.1]Ma et al., 2007; [2.2]Kondo et al., 2008; [2.3]Chen et al., 2010
	Chloroplast	Increase in NR, NiR <sup>[3.4]</sup> , photosynthetic rate <sup>[3.5]</sup> , and chlorophyll content, etc.	Decline in chlorophyll content, ratio of Fv/Fm <sup>[2.3]</sup> , and apparent photosynthesis <sup>[3.1]</sup> ; Accumulation of NO <sub>3</sub> <sup>-</sup> and NO <sub>2</sub> <sup>-</sup> <sup>[3.2]</sup> as well as increase in NR and NiR <sup>[3.2]</sup> ; Inhibition of NR <sup>[3.3]</sup>	[3.1]Srivastava et al., 1974; [3.2]Yoneyama et al., 1979; [3.3]Hisamatsu et al., 1988; [3.4]Weber et al., 1995 [3.5]Schmutz et al., 1995
	Mitochondria/ Peroxisome		Inhibition of dark respiration and apparent photorespiration <sup>[3.1,4.2]</sup> ; Protrusions from both plastids and mitochondria of <i>Phaseolus vulgaris</i> exposed to NO <sub>2</sub> (10 ml.l <sup>-1</sup> ) <sup>[4.1]</sup>	[4.1]Dolzmann and Ullrich, 1966; [4.2]Carlson, 1983
	In developing or maturing leaves	Increased leaf area <sup>[1.2]</sup> ; NO <sub>2</sub> -N incorporation into free amino acids such as Glu, Asp and Gln <sup>[3.4;5.3]</sup> ; Stimulation on cell proliferation and enlargement as well as up-regulation of the related genes, such as ARGOS, GRF5, and KLU <sup>[5.4]</sup>	NO <sub>2</sub> -N incorporation into free amino acids such as glutamine, glutamic acid, γ-amino butyric acid and aspartic acid <sup>[5.1]</sup> ; NO <sub>2</sub> led to swollen thylakoids and a reduction in the number of grana stacks <sup>[5.2]</sup>	[5.1]Yoneyama and Sasakawa, 1979; [5.2]Schiffgens-Gruber and Lutz, 1992 [5.3]Nussbaum et al., 1993; [5.4]Takahashi et al., 2014
Stems	Xylem	Enlarged width of xylem in the main stem of Poplar trees <sup>[3.5]</sup>	stem growth significantly decreased by NO <sub>2</sub> at 1.0 μL.l <sup>-1</sup> <sup>[6.1]</sup>	[6.1]Eastham and Ormrod, 1986
	Phloem	NO <sub>2</sub> -N incorporation into free amino acids of bark of Norway spruce <sup>[5.3]</sup>	NO <sub>2</sub> -N incorporation into free amino acids such as serine, asparagine and glutamine <sup>[6.2]</sup>	[6.2]Wellburn, 1990
Roots		NO <sub>2</sub> -N incorporation into free amino acids in Norway spruce roots <sup>[5.3]</sup>	Decrease in root/shoot ratio, dry matter production, concentration of soluble sugars in roots, root respiration of kidney bean plants <sup>[7.1]</sup> Decrease in root nitrate uptake in sunflower plants <sup>[7.2]</sup> and soybean plants <sup>[1.1]</sup> , increase in the ammonium concentration in roots of soybean plants at 1.1 μL.l <sup>-1</sup> NO <sub>2</sub> <sup>[1.1]</sup>	[7.1]Ito et al., 1985; [7.2]Okano et al., 1985
Flowers		Acceleration of flowering time and increase in flower number <sup>[5.4;8.1]</sup>		[8.1]Takahashi et al., 2011;
Fruits		Increased fruit yield <sup>[8.1]</sup> or grain yield (the number and weight of grain) and protein stored (at NO <sub>2</sub> of 170 nL.l <sup>-1</sup> ) <sup>[9.1]</sup>		[9.1]Murray et al., 1994

reductase (NR) is considered as a key rate-limiting enzyme of NO<sub>2</sub>-N assimilation (Hawkesford et al., 2012). A linear correlation was found between NR activity, NO<sub>2</sub> concentration and amounts of N incorporated into amino acids (Sparks et al., 2001). However, high levels of NO<sub>2</sub> fumigation resulted into a loss of NR activity or a rapid inactivation of the leaf NR (Takeuchi et al., 1985; Hisamatsu et al., 1988), and NO<sub>3</sub><sup>-</sup> accumulation (Ma et al., 2007). This down-regulation of the NR may be ascribed to at least one of the following phenomena: (i) a high NO<sub>2</sub> concentration will inhibit the activities of glutamine synthetase and glutamate synthase, which leads to NH<sub>4</sub><sup>+</sup> accumulation and subsequently brings about a loss of NR activity (Orebamjo and Stewart, 1975; Padidam et al., 1991), and (ii) a high NO<sub>2</sub>-induced stomatal closure may lead to a rapid NR inactivation due to a low CO<sub>2</sub> availability. High NO<sub>2</sub> rapidly induced stomatal closure (Qiao and Murray, 1998). Stomatal closure may trigger a chain reaction wherein the lower CO<sub>2</sub> availability will lead to the subsequent leaf NR activity decrease (Kaiser and Forster, 1989). *NIA1* and *NIA2* genes encode the two isoforms of the NR apoprotein (Wilkinson and Crawford, 1993). Recent reports show a close relationship between the expression of *NIA1/NIA2* genes and *NRT1/NRT2* genes. Addition of external NO<sub>3</sub><sup>-</sup> strongly induced the expression of genes encoding the NR (*AtNIA1* and *AtNIA2*) and the transmembrane transporters (*AtNRT1.1*, *AtNRT2.1*, *NpNRT2.1*; Fraiser et al., 2001; Jonassen et al., 2009). In contrast, high levels of external NO<sub>3</sub><sup>-</sup> caused a down-regulation of *AtNRT1.1* and *AtNIA1* through a pathway of NO<sub>2</sub><sup>-</sup>-induced repression (Loqué et al., 2003). This down-regulation of the *AtNRT1.1* gene is associated with a decrease in the NO<sub>3</sub><sup>-</sup> influx. Earlier studies showed that high NO<sub>2</sub> concentration fumigation under light or dark conditions resulted in leaf NO<sub>2</sub><sup>-</sup> accumulation (Yoneyama and Sasakawa, 1979; Yu et al., 1988). Thus, we may assume that a high NO<sub>2</sub>-caused NO<sub>2</sub><sup>-</sup> accumulation may lead to a negative feedback regulation on leaf NO<sub>2</sub> uptake through the down-regulation of the *NRT1.1* gene and the subsequent repression of the NO<sub>3</sub><sup>-</sup> influx. Moreover, studies on NR mutants showed that *AtNRT1.1*, *AtNRT1.7*, *NpNRT2.1*, and *AtNIA1* are up-regulated in NR-deficient mutants (*NIA2*- and/or *MoCo* biosynthesis-deficient mutants) (Lejay et al., 1999; Vidmar et al., 2000; Fan et al., 2009). Under NR-repressible or -deficient conditions, this up-regulation of the transporter genes may be beneficial to an exportation of excess NO<sub>3</sub><sup>-</sup> in the leaf (Fan et al., 2009). Moreover, Jonassen et al. (2009) demonstrated that the bZIP transcription factors *HY5* and *HYH* regulate positively *NIA2* gene and negatively *NRT1.1* gene. However, *HY5* and *HYH* appear to be mediated by light but not by external NO<sub>3</sub><sup>-</sup>.

Excess NO<sub>3</sub><sup>-</sup> may be accumulated in leaf vacuole (Hawkesford et al., 2012). The H<sup>+</sup>/NO<sub>3</sub><sup>-</sup> antiport across tonoplast is responsible for NO<sub>3</sub><sup>-</sup> influx and H<sup>+</sup>/NO<sub>3</sub><sup>-</sup> symport for NO<sub>3</sub><sup>-</sup> efflux (Figure 1). The flux direction will depend on the requirements and conditions of the cell (Schumaker and Sze, 1987). Three members of the chloride channel family (CLC) genes *AtClCa* (De Angeli et al., 2006), *AtClCc* (Harada et al., 2004) and *AtClCd* (Lv et al., 2009) have been identified in the leaf tonoplast. De Angeli et al. (2009) demonstrated that adenosine triphosphate (ATP) induces a negative regulation on *AtClCa* activity. NO<sub>2</sub> fumigation significantly increased ATP amounts of *Lolium perenne*

and *Phleum pratense*, the amounts increasing with raising NO<sub>2</sub> concentrations (Wellburn et al., 1981). This may be due to the formation of free radicals in response to NO<sub>2</sub> fumigation, which may damage photosynthetic membranes and hence alter the proton gradients to which ATP formation is linked (Wellburn et al., 1981). Yoneyama and Sasakawa (1979) found that 8 ppm NO<sub>2</sub> fumigation under dark conditions resulted in NO<sub>2</sub><sup>-</sup> accumulation in spinach leaves. High doses of NO<sub>2</sub><sup>-</sup> resulted in the peroxidation of lipid constituents of chloroplastic membrane (Ezzine and Ghorbel, 2006). Chen et al. (2010) found that leaf uptake of NO<sub>2</sub> reduced the rate of photosynthesis and increased the malondialdehyde (MDA) concentration, may be due to a competition for nicotinamide adenine dinucleotide phosphate (NADPH) between the processes of NO<sub>2</sub><sup>-</sup> reduction vs. carbon assimilation, and the generation of reactive oxygen species (ROS) (Sabaratnam and Gupat, 1988; Shimazaki et al., 1992).

### TRANSMEMBRANE TRANSPORTERS IN LEAVES MEDIATING NO<sub>3</sub><sup>-</sup> SIGNALING

NO<sub>2</sub> fumigation may significantly disrupt plant morphology and physiology by, for instance, changing the shoot to root ratio, stomatal, and gas exchange dynamics, or modifying root N uptake (Qiao and Murray, 1997, 1998; Table 1). The exposure of plants to NO<sub>2</sub> increased the total content of soluble free amino acids in leaves and shoots (Nussbaum et al., 1993). Most of the amino acids may be used locally for the synthesis of e.g., Chlorophyll and Rubisco during rapid vegetative growth, or be ultimately designated for e.g., filling pods (Imsande and Touraine, 1994). NO<sub>3</sub><sup>-</sup> assimilation products (protein/nucleic acids and amino acids/amides) can also be transferred into roots under soil N deficit (Wellburn, 1990). Under a low NO<sub>3</sub><sup>-</sup> supply, gaseous NO<sub>2</sub> may change the amino acid ratio of the xylem. For example, the amount of serine, asparagine and glutamine were high in the xylem of plants exposed to atmospheric NO<sub>2</sub>, whereas arginine, cysteine, valine and lysine were high in the control plants (Wellburn, 1990; Table 1). Moreover, NO<sub>2</sub> treatment increased phloem transport of organic N and inhibited the rate of xylem N translocation.

NO<sub>2</sub>-N metabolism and the mobilization of metabolic products will trigger various signaling pathways that regulate the physiological and metabolic processes. The dissolution of NO<sub>2</sub> and subsequent reduction can result in root NO<sub>3</sub><sup>-</sup> uptake changes. The xylem is part of leaf apoplast (Felle and Hanstein, 2002). Thus, NO<sub>2</sub>-caused apoplastic pH changes may serve as a signal to modify the uptake of NO<sub>3</sub><sup>-</sup> via the root system. This NO<sub>3</sub><sup>-</sup> signaling pathway has been explained by Qiao and Murray (1998). Moreover, NO<sub>3</sub><sup>-</sup> reduction can produce malate (Touraine et al., 1988); the organic acid needs to be membrane transported to be loaded into the phloem. A transport of malate from leaves to roots can serve as another signal to control root uptake of NO<sub>3</sub><sup>-</sup> (Touraine et al., 1992). This signaling pathway has been reported by Imsande and Touraine (1994). Tonoplast transport of malate plays an important role in physiological regulation in NO<sub>3</sub><sup>-</sup> nutrition (Hawkesford et al., 2012). At the cellular level, NO<sub>3</sub><sup>-</sup> accumulation led to increased expression of genes encoding organic acid synthesis (*PPC*, cytosolic PK, *CS*, *ICDH-1*) and accumulation of malate and a-oxoglutarat. In contrast, leaf

malate supply can inhibit *NIA* expression, affecting both the *NIA* transcript level and the activity (Müller et al., 2001).

NO<sub>3</sub><sup>-</sup> itself may serve as a signaling molecule (Scheible et al., 1997). NO<sub>3</sub><sup>-</sup> addition to the growing media can induce or repress the expression of various genes encoding e.g., NO<sub>3</sub><sup>-</sup> transporters, NO<sub>3</sub><sup>-</sup>/NO<sub>2</sub><sup>-</sup> reductase, ferredoxin reductase, and the enzymes in the pentose phosphate pathway, or iron or sulfate transport and metabolism (Wang et al., 2003; Marschner, 2012). The expression of NO<sub>3</sub><sup>-</sup>-responsive genes [such as NADH-specific and NAD(P)H-bispecific NR genes] is dependent upon NO<sub>3</sub><sup>-</sup> flux but not on the NO<sub>3</sub><sup>-</sup> amount stored in the tissue (Gojon et al., 1991). Excess external NO<sub>3</sub><sup>-</sup> may be stored in several vacuoles and recirculated after storage (Hawkesford et al., 2012). NO<sub>3</sub><sup>-</sup> remobilization may occur among different organs, for instance, from older leaves to younger leaves during the vegetative stage or from leaves to seeds during the reproductive stage (Schiltz et al., 2005; Hawkesford et al., 2012). N remobilization is rate-limited by the transport of NO<sub>3</sub><sup>-</sup> across tonoplast of vacuole, plasma membrane of mesophyll cell, plasma membrane of companion cell and sieve element, and phloem loading (Fan et al., 2009). CLC genes (*AtClCa*, *AtClCc*, and *AtClCd*) are required for the transport of NO<sub>3</sub><sup>-</sup> across tonoplast in vacuole. Disruption of one of these genes will influence the flux of NO<sub>3</sub><sup>-</sup> in vacuoles (De Angeli et al., 2006). *AtNRT2.4* showed a strong induction in a low NO<sub>3</sub><sup>-</sup> provision. Orsel et al. (2004) suggested that *AtNRT2.4* and *AtNRT2.5* participate in the transport of NO<sub>3</sub><sup>-</sup> from stored pools (vacuoles) to cytoplasm. Moreover, *AtNRT2.7* was also involved in this type of NO<sub>3</sub><sup>-</sup> flux; this gene could play roles in leaf balance between the amount of NO<sub>3</sub><sup>-</sup> used for assimilation and that re-absorbed for further transport (Orsel et al., 2002). Four NRT1 family genes (*AtNRT1.4*, *AtNRT1.7*, *AtNRT1.11*, and *AtNRT1.12*) participate in the phloem- and/or xylem-loading of NO<sub>3</sub><sup>-</sup> (Figure 1). *AtNRT1.4* was expressed predominantly in the leaf petiole and involved in petiole NO<sub>3</sub><sup>-</sup> accumulation (Chiu et al., 2004). The mutation of the *AtNRT1:4* resulted in significant changes of NO<sub>3</sub><sup>-</sup> content in leaf petiole and the lamina. Furthermore, the deficiency of *AtNRT1.4* can alter leaf development. *NRT1.7* was expressed in the phloem of the leaf minor vein and mediated the remobilization of excess NO<sub>3</sub><sup>-</sup> from older leaves to younger leaves (Fan et al., 2009). Compared with the wild-type plants, the *nrt1.7* null mutants accumulated a higher amount of NO<sub>3</sub><sup>-</sup> in the older leaves and decreased the NO<sub>3</sub><sup>-</sup> content of phloem exudates from older leaves. The newly identified NO<sub>3</sub><sup>-</sup> transporters (*NRT1.11* and *NRT1.12*) were expressed in the companion cells of the major vein (Hsu and Tsay, 2013). They play roles in xylem-to-phloem transfer for redistributing NO<sub>3</sub><sup>-</sup> into developing leaves. Moreover, several NRT2 genes may also be involved in the N remobilization. For example, *ZmNrt2.1* plays a potential role in NO<sub>3</sub><sup>-</sup> loading from the xylem or in its compartmentation (Quaggiotti et al., 2003).

## CONCLUSION

The substomatal build-up of NO<sub>2</sub> and subsequent NO<sub>3</sub><sup>-</sup> metabolism may lead to apoplastic alkalization or acidification and to NO<sub>3</sub><sup>-</sup>/NO<sub>2</sub><sup>-</sup> concentration fluctuations in the leaf apoplast and symplast (e.g., cytoplasm and vacuole), which depend on NO<sub>2</sub> concentration and root N supply. These changes

will cause complex responses of NO<sub>3</sub><sup>-</sup>-responsive genes encoding NO<sub>3</sub><sup>-</sup> transporters and NO<sub>3</sub><sup>-</sup>/NO<sub>2</sub><sup>-</sup> reductase. For example, addition of external NO<sub>3</sub><sup>-</sup> produced a strong induction on NR genes (such as *AtNIA1* and *AtNIA2*) and the transporter genes (such as *AtNRT1.1*, *AtNRT2.1*, and *NpNRT2.1*). However, excess NO<sub>2</sub><sup>-</sup> significantly inhibited the expression of *AtNRT1.1* and *AtNIA1*, and disturbed CLC family genes by regulating the generation of ATP. This down-regulation of the *NRT1.1* gene is associated with a decrease in NO<sub>3</sub><sup>-</sup> influx. Moreover, *AtNRT2.4*, *AtNRT2.5*, and *AtNRT2.7* may participate in the transfer of NO<sub>3</sub><sup>-</sup> from stored pools (vacuoles) to cytoplasm. *AtNRT1.4*, *AtNRT1.7*, *AtNRT1.11*, and *AtNRT1.12* are involved in the phloem- and/or xylem-loading of NO<sub>3</sub><sup>-</sup>. Thus, these genes are suggested to play rate-limiting roles in foliar uptake of NO<sub>2</sub>. Further work is proposed to investigate the relationship between organ specificity of NRT1/NRT2 gene expression and species-specific NO<sub>2</sub> uptake.

In practical terms, a high rate of low concentration NO<sub>2</sub> absorption by the foliage may be positive for preserving an adequate plant N status. However, a high NO<sub>2</sub> concentration may alter leaf apoplast chemistry, leading to the accumulation of NO<sub>3</sub><sup>-</sup> and NO<sub>2</sub><sup>-</sup>, and providing signals which may negatively affect plant N nutrition. These factors are however closely linked with leaf NO<sub>3</sub><sup>-</sup> transporters and may also interact with the foliar uptake processes (e.g., by promoting stomatal closure). Thereby, a low NO<sub>2</sub> concentration may act as a positive regulation signal (Takahashi et al., 2014) by stimulating the leaf NO<sub>3</sub><sup>-</sup> transporters, and enhancing NO<sub>3</sub><sup>-</sup> transport and distribution. In contrast, a high NO<sub>2</sub> concentration in relation to a high rate of foliar NO<sub>2</sub> absorption, may repress the expression of NO<sub>3</sub><sup>-</sup> transporters and enzymes, which may protect the cells or organelles from NO<sub>2</sub> damage.

## AUTHOR CONTRIBUTIONS

Yanbo Hu organized and wrote the original manuscript; Yanbo Hu and Victoria Fernandez discussed and revised the revising manuscript and approved the final version; Ling Ma collected the information for Table 1, and went through the manuscript.

## ACKNOWLEDGMENTS

This work was supported by grants from “Fundamental Research Funds for the Central Universities” (Grant No. 2572014CA24, DL10BB24) and “National Natural Science Foundation” (Grant No. 31300506).

## REFERENCES

- Bertoni, G. (2012). A nitrate transporter for both roots and shoots. *Plant Cell* 24, 1. doi: 10.1105/tpc.112.240110
- Carlson, R. W. (1983). Interaction between SO<sub>2</sub> and NO<sub>2</sub> and their effects on photosynthetic properties of soybean *Glycine max*. *Environ. Pollut. Ecol. Biol.* 32, 11–38. doi: 10.1016/0143-1471(83)90071-5
- Chen, Z. M., Chen, Y. X., Du, G. J., Wu, X. L., and Li, F. (2010). Effects of 60-day NO<sub>2</sub> fumigation on growth, oxidative stress and antioxidative response in *Cinnamomum camphora* seedlings. *J. Zhejiang Univ. Sci. B* 11, 190–199. doi: 10.1631/jzus.B0910350
- Chiu, C. C., Lin, C. S., Hsia, A. P., Su, R. C., Lin, H. L., and Tsay, Y. F. (2004). Mutation of a nitrate transporter, *AtNRT1:4*, results in a reduced petiole nitrate content and altered leaf development. *Plant Cell Physiol.* 45, 1139–1148. doi: 10.1093/pcp/pch143

- Chopin, F., Orsel, M., Dorbe, M. F., Chardon, F., Truong, H. N., Miller, A. J., et al. (2007). The Arabidopsis AtNRT2.7 nitrate transporter controls nitrate content in seeds. *Plant Cell* 19, 1590–1602. doi: 10.1105/tpc.107.050542
- Crawford, N. M., and Glass, A. D. M. (1998). Molecular and physiological aspects of nitrate uptake in plants. *Trends Plant Sci.* 10, 389–395. doi: 10.1016/S1360-1385(98)01311-9
- De Angeli, A., Monachello, D., Ephritikhine, G., Frachisse, J. M., Thomine, S., Gambale, F., et al. (2006). The nitrate/proton antiporter AtCLCa mediates nitrate accumulation in plant vacuoles. *Nature* 442, 939–942. doi: 10.1038/nature05013
- De Angeli, A., Moran, O., Wege, S., Filleur, S., Ephritikhine, G., Thomine, S., et al. (2009). ATP binding to the C terminus of the Arabidopsis thaliana nitrate/proton antiporter, AtCLCa, regulates nitrate transport into plant vacuoles. *J. Biol. Chem.* 284, 26526–26532. doi: 10.1074/jbc.M109.005132
- Dolzmann, P., and Ullrich, H. (1966). Einige Beobachtungen über Beziehungen zwischen Chloroplasten und Mitochondrien im Palisadenparenchym von *Phaseolus vulgaris*. *Zeitschrift für Pflanzenphysiologie* 55, 165–180.
- Eastham, A. M., and Ormrod, D. P. (1986). Visible injury and growth responses of young cuttings of *Populus Canadensis* and *P. nigra* to nitrogen dioxide and sulphur dioxide. *Can J. For. Res.* 16, 1289–1292. doi: 10.1139/x86-228
- Eichert, T., and Fernández, V. (2012). “Uptake and release of elements by leaves and other aerial plant parts,” in *Marschner’s Mineral Nutrition of Higher Plants*, ed P. Marschner (Oxford: Academic Press), 71–84.
- Ezzine, M., and Ghorbel, M. H. (2006). Physiological and biochemical responses resulting from nitrite accumulation in tomato. (*Lycopersicon esculentum* Mill. cv. Ibiza F1). *J. Plant Physiol.* 163, 1032–1039. doi: 10.1016/j.jplph.2005.07.013
- Fan, S. C., Lin, C. S., Hsu, P. K., Lin, S. H., and Tsay, Y. F. (2009). The Arabidopsis nitrate transporter NRT1.7, expressed in phloem, is responsible for source-to-sink remobilization of nitrate. *Plant Cell* 21, 2750–2761. doi: 10.1105/tpc.109.067603
- Felle, H. H., and Hanstein, S. (2002). The apoplastic pH of the substomatal cavity of *Vicia faba* leaves and its regulation responding to different stress factors. *J. Exp. Bot.* 53, 73–82. doi: 10.1093/jxb/53.366.73
- Fernández, V., and Brown, P. H. (2013). From plant surface to plant metabolism: the uncertain fate of foliar-applied nutrients. *Front. Plant Sci.* 4:289. doi: 10.3389/fpls.2013.00289
- Fraïsis, V., Dorbe, M. F., and Daniel-Vedele, F. (2001). Identification and expression analyses of two genes encoding putative low-affinity nitrate transporters from *Nicotiana glauca*. *Plant Mol. Biol.* 45, 181–190. doi: 10.1023/A:1006426616760
- Fujita, T., Noguchi, K., and Terashima, I. (2013). Apoplastic mesophyll signals induce rapid stomatal responses to CO<sub>2</sub> in *Commelinacommunis*. *New Phytol.* 199, 395–406. doi: 10.1111/nph.12261
- Geelen, D., Lurin, C., Bouchez, D., Frachisse, J. M., Lelièvre, F., Courtial, B., et al. (2000). Disruption of putative anion channel gene AtCLC-a in Arabidopsis suggests a role in the regulation of nitrate content. *Plant J.* 21, 259–267. doi: 10.1046/j.1365-313x.2000.00680.x
- Gojon, A., Wakrim, R., Passama, L., and Robin, P. (1991). Regulation of NO<sub>3</sub><sup>-</sup> assimilation by anion availability in excised soybean leaves. *Plant Physiol.* 96, 398–405. doi: 10.1104/pp.96.2.398
- Guo, F. Q., Young, J., and Crawford, N. M. (2003). The nitrate transporter AtNRT1.1 (CHL1) functions in stomatal opening and contributes to drought susceptibility in Arabidopsis. *Plant Cell* 15, 107–117. doi: 10.1105/tpc.006312
- Harada, H., Kuromori, T., Hirayama, T., Shinozaki, K., and Leigh, R. A. (2004). Quantitative trait loci analysis of nitrate storage in Arabidopsis leading to an investigation of the contribution of the anion channel gene, AtCLC-c, to variation in nitrate levels. *J. Exp. Bot.* 55, 2005–2014. doi: 10.1093/jxb/erh224
- Hawkesford, M., Horst, W., Kichey, T., Lambers, H., Schjoerring, J., Skrumsager, I., et al. (2012). “Functions of Macronutrients,” in *Marschner’s Mineral Nutrition of Higher Plants*, 3rd Edn., eds P. Marschner (London: Elsevier), 135–178.
- Hedrich, R., Neimainis, S., Savchenko, G., Felle, H. H., Kaiser, W. M., and Heber, U. (2001). Changes in apoplastic pH and membrane potential in leaves in relation to stomatal responses to CO<sub>2</sub>, malate, abscisic acid or interruption of water supply. *Planta* 213, 594–601. doi: 10.1007/s004250100524
- Hisamatsu, S., Nihira, J., Takeuchi, Y. C., Satoh, S., and Kondo, N. (1988). NO<sub>2</sub> suppression of light-induced nitrate reductase in squash cotyledons. *Plant Cell Physiol.* 29, 395–401.
- Hoffmann, B., Plänker, R., and Mengel, K. (1992). Measurements of pH in the apoplast of sunflower leaves by means of fluorescence. *Physiol. Plant.* 84, 146–153. doi: 10.1111/j.1399-3054.1992.tb08777.x
- Hsu, P. K., and Tsay, Y. F. (2013). Two phloem nitrate transporters, NRT1.11 and NRT1.12, are important for redistributing xylem-borne nitrate to enhance plant growth. *Plant Physiol.* 163, 844–856. doi: 10.1104/pp.113.226563
- Hu, Y. B. (2011). NO<sub>2</sub>-driven NO<sub>3</sub><sup>-</sup> metabolism in leaves. *Insci. J.* 1, 90–101. doi: 10.5640/insc.010290
- Hu, Y. B., Bellaloui, N., Sun, G. Y., Tigabu, M., and Wang, J. H. (2014). Exogenous sodium sulfide improves morphological and physiological responses of a hybrid Populus species to nitrogen dioxide. *J. Plant Physiol.* 171, 868–875. doi: 10.1016/j.jplph.2013.10.018
- Hu, Y. B., and Sun, G. Y. (2010). Leaf nitrogen dioxide uptake coupling apoplastic chemistry, carbon/sulfur assimilation, and plant nitrogen status. *Plant Cell Rep.* 29, 1069–1077. doi: 10.1007/s00299-010-0898-5
- Imssande, J., and Touraine, B. (1994). N demand and the regulation of nitrate uptake. *Plant Physiol.* 105, 3–7.
- Ito, O., Okano, K., Kuroiwa, M., and Totsuka, T. (1985). Effects of NO<sub>2</sub> and O<sub>3</sub> alone or in combination on kidney bean plants (*Phaseolus vulgaris* L.): partitioning of assimilates and root activities. *J. Exp. Bot.* 36, 652–662. doi: 10.1093/jxb/36.4.652
- Jonassen, E. M., Sevin, D. C., and Lillo, C. (2009). The bZIP transcription factors HY5 and HYH are positive regulators of the main nitrate reductase gene in Arabidopsis leaves, NIA2, but negative regulators of the nitrate uptake gene NRT1.1. *J. Plant Physiol.* 166, 2071–2076. doi: 10.1016/j.jplph.2009.05.010
- Kaiser, W. M., and Forster, J. (1989). Low CO<sub>2</sub> prevents nitrate reduction in leaves. *Plant Physiol.* 91, 970–974. doi: 10.1104/pp.91.3.970
- Kondo, K., Yamada, K., Nakagawa, A., Takahashi, M., Morikawa, H., and Sakamoto, A. (2008). Molecular characterization of atmospheric NO<sub>2</sub>-responsive germin-like proteins in azalea leaves. *Biochem. Biophys. Res. Commun.* 377, 857–861. doi: 10.1016/j.bbrc.2008.10.060
- Kosegarten, H. U., Hoffmann, B., and Mengel, K. (1999). Apoplastic pH and Fe<sup>3+</sup> reduction in intact sunflower leaves. *Plant Physiol.* 121, 1069–1079. doi: 10.1104/pp.121.4.1069
- Lejay, L., Tillard, P., Lepetit, M., Olive, F. D., Filleur, S., Daniel-Vedele, F., et al. (1999). Molecular and functional regulation of two NO<sub>3</sub><sup>-</sup> uptake systems by N- and C-status of Arabidopsis plants. *Plant J.* 18, 509–519. doi: 10.1046/j.1365-313X.1999.00480.x
- Loqué, D., Tillard, P., Gojon, A., and Lepetit, M. (2003). Gene expression of the NO<sub>3</sub><sup>-</sup> transporter NRT1.1 and the nitrate reductase NIA1 is repressed in Arabidopsis roots by NO<sub>2</sub><sup>-</sup>, the product of NO<sub>3</sub><sup>-</sup> reduction. *Plant Physiol.* 132, 958–967. doi: 10.1104/pp.102.018523
- Lv, Q. D., Tang, R. J., Liu, H., Gao, X. S., Li, Y. Z., Zheng, H. Q., et al. (2009). Cloning and molecular analyses of the Arabidopsis thaliana chloride channel gene family. *Plant Sci.* 176, 650–661. doi: 10.1016/j.plantsci.2009.02.006
- Ma, C. Y., Xu, X., Hao, L., and Cao, J. (2007). Nitrogen dioxide-induced responses in Brassica campestris seedlings: the role of hydrogen peroxide in the modulation of antioxidative level and induced resistance. *Agric. Sci. China* 6, 1193–1200. doi: 10.1016/S1671-2927(07)60163-1
- Marschner, P. (2012). *Mineral Nutrition of Higher Plants*. San Diego, CA: Academic Press.
- Morikawa, H., Higaki, A., Nohno, M., Takahashi, M., Kamada, M., Nakata, M., et al. (1998). More than a 600-fold variation in nitrogen dioxide assimilation among 217 plant taxa. *Plant Cell Environ.* 21, 180–190. doi: 10.1046/j.1365-3040.1998.00255.x
- Mühlhling, K. H., and Lauchli, A. (2001). Influence of chemical form and concentration of nitrogen on apoplastic pH of leaves. *J. Plant Nutr.* 25, 399–411. doi: 10.1081/PLN-100104968
- Müller, C., Scheible, W. R., Stitt, M., and Krapp, A. (2001). Influence of malate and 2-oxoglutarate on the NIA transcript level and nitrate reductase activity in tobacco leaves. *Plant Cell Environ.* 24, 191–203. doi: 10.1111/j.1365-3040.2001.00664.x
- Murray, F., Wilson, S., and Samaraweera, S. (1994). NO<sub>2</sub> increases wheat grain yield even in the presence of SO<sub>2</sub>. *Agric. Ecosyst. Environ.* 48, 115–123. doi: 10.1016/0167-8809(94)90082-5
- Nussbaum, S., Ballmoos, P. V., Gfeller, H., Schlunegger, U. P., Fuhrer, J., Rhodes, D., et al. (1993). Incorporation of atmospheric <sup>15</sup>NO<sub>2</sub>-nitrogen into free amino

- acids by Norway spruce *Picea abies* (L.) Karst. *Oecologia* 94, 408–414. doi: 10.1007/BF00317117
- Okamoto, M., Vidmar, J. J., and Glass, A. D. (2003). Regulation of NRT1 and NRT2 gene families of *Arabidopsis thaliana*: responses to nitrate provision. *Plant Cell Physiol.* 44, 304–317. doi: 10.1093/pcp/pcg036
- Okano, K., Totsuka, T., Fukuzawa, T., and Tazaki, T. (1985). Growth responses of plants to various concentrations of nitrogen dioxide. *Environ. Pollut.* 38, 361–373. doi: 10.1016/0143-1471(85)90107-2
- Ono, F., Frommer, W. B., and von Wire'n, N. (2000). Coordinated diurnal regulation of low- and high-affinity nitrate transporters in tomato. *Plant Biol.* 2, 17–23. doi: 10.1055/s-2000-297
- Orebamjo, T. O., and Stewart, G. R. (1975). Ammonium inactivation of nitrate reductase in *Lemna minor* L. *Planta* 122, 37–44. doi: 10.1007/BF00385402
- Orsel, M., Eulenburg, K., Krapp, A., and Daniel-Vedele, F. (2004). Disruption of the nitrate transporter genes AtNRT2.1 and AtNRT2.2 restricts growth at low external nitrate concentration. *Planta* 219, 714–721. doi: 10.1007/s00425-004-1266-x
- Orsel, M., Krapp, A., and Daniel-Vedele, F. (2002). Analysis of the NRT2 nitrate transporter family in *Arabidopsis*: structure and gene expression. *Plant Physiol.* 129, 886–896. doi: 10.1104/pp.005280
- Padidam, M., Venkateswarlu, K., and Johri, M. M. (1991). Ammonium represses NADPH nitrate reductase in the moss *Funaria hygrometrica*. *Plant Sci.* 75, 185–194. doi: 10.1016/0168-9452(91)90233-X
- Popova, O. V., Dietz, K. J., and Gollmack, D. (2003). Salt-dependent expression of a nitrate transporter and two amino acid transporter genes in *Mesembryanthemum crystallinum*. *Plant Mol. Biol.* 52, 569–578. doi: 10.1023/A:1024802101057
- Qiao, Z., and Murray, F. (1997). Effects of atmospheric nitrogen dioxide on uptake and assimilation of ammonium in soybean plants. *J. Plant Nutr.* 20, 1183–1190. doi: 10.1080/01904169709365326
- Qiao, Z., and Murray, F. (1998). The effects of NO<sub>2</sub> on the uptake and assimilation of nitrate by soybean plants. *Environ. Exp. Bot.* 10, 33–40. doi: 10.1016/S0098-8472(97)00023-3
- Quaggiotti, S., Ruperti, B., Borsa, P., Destro, T., and Malagoli, M. (2003). Expression of a putative high-affinity NO<sub>3</sub><sup>-</sup> transporter and of an H<sup>+</sup>-ATPase in relation to whole plant nitrate transport physiology in two maize genotypes differently responsive to low nitrogen availability. *J. Exp. Bot.* 54, 1023–1031. doi: 10.1093/jxb/erg106
- Raven, J. A. (1988). Acquisition of nitrogen by the shoots of land plants: its occurrence and implication for acid-base regulation. *New Phytol.* 109, 1–20. doi: 10.1111/j.1469-8137.1988.tb00212.x
- Sabaratnam, S., and Gupat, G. (1988). Effects of nitrogen dioxide on biochemical and physiological characteristics of soybean. *Environ. Pollut.* 55, 149–158. doi: 10.1016/0269-7491(88)90125-X
- Sattelmacher, B. (2001). The apoplast and its significance for plant mineral nutrition. *New Phytol.* 149, 167–192. doi: 10.1046/j.1469-8137.2001.00034.x
- Savchenko, G., Wiese, C., Neimanis, S., Hedrich, R., and Heber, U. (2000). pH regulation in apoplastic and cytoplasmic cell compartments of leaves. *Planta* 211, 246–255. doi: 10.1007/s004250000280
- Scheible, W. R., Gonzalez-Fontes, A., Lauerer, M., Muller-Rober, B., Caboche, M., and Stitt, M. (1997). Nitrate acts as a signal to induce organic acid metabolism and repress starch metabolism in tobacco. *Plant Cell* 9, 783–798. doi: 10.1105/tpc.9.5.783
- Schiffgens-Gruber, A., and Lutz, C. (1992). Ultrastructure of mesophyll cell chloroplasts of spruce needles exposed to O<sub>3</sub>, SO<sub>2</sub> and NO<sub>2</sub> alone and in combination. *Environ. Exp. Bot.* 32, 243–254. doi: 10.1016/0098-8472(92)90007-O
- Schiltz, S., Munier-Jolain, N., Jeudy, C., Burstin, J., and Salon, C. (2005). Dynamics of exogenous nitrogen partitioning and nitrogen remobilization from vegetative organs in pea revealed by <sup>15</sup>N *in vivo* labeling throughout seed filling. *Plant Physiol.* 137, 1463–1473. doi: 10.1104/pp.104.056713
- Schmutz, P., Tarjan, D., Günthardt-Goerg, M. S., Matyssek, R., and Bucher, J. B. (1995). Nitrogen dioxide - a gaseous fertilizer of Poplar trees. *Phyton* 35, 219–232.
- Schumaker, K. S., and Sze, H. (1987). Decrease of pH gradients in tonoplast vesicles by NO<sub>3</sub><sup>-</sup> and Cl<sup>-</sup>: evidence for H<sup>+</sup>-coupled anion transport. *Plant Physiol.* 83, 490–496. doi: 10.1104/pp.83.3.490
- Shimazaki, K., Yu, S. W., Sakaki, T., and Tanaka, K. (1992). Differences between spinach and kidney bean plants in terms of sensitivity to fumigation with NO<sub>2</sub>. *Plant Cell Physiol.* 33, 267–273.
- Siegwolf, R. T. W., Matyssek, R., Saurer, M., Maurer, S., Günthardt-Goerg, M. S., Schmutz, P., et al. (2001). Stable isotope analysis reveals differential effects of soil nitrogen and nitrogen dioxide on the water use efficiency in hybrid poplar leaves. *New Phytol.* 149, 233–246. doi: 10.1046/j.1469-8137.2001.00032.x
- Sparks, J. P., Monson, R. K., Sparks, K. L., and Lerdau, M. (2001). Leaf uptake of nitrogen dioxide (NO<sub>2</sub>) in a tropical wet forest: implications for tropospheric chemistry. *Oecologia* 127, 214–221. doi: 10.1007/s004420000594
- Speer, M., and Kaiser, W. M. (1991). Ion relations of symplastic and apoplastic space in leaves from *Spinacia oleracea* L. and *Pisum sativum* L. under salinity. *Plant Physiol.* 97, 990–997. doi: 10.1104/pp.97.3.990
- Srivastava, H. S., Jolliffe, P. A., and Runeckles, V. C. (1974). Inhibition of gas exchange in bean leaves by NO<sub>2</sub>. *Can. J. Bot.* 53, 466–474. doi: 10.1139/b75-057
- Stiegler, J. C., Richardson, M. D., and Karcher, D. E. (2011). Foliar nitrogen uptake following urea application to putting green turfgrass species. *Crop Sci.* 51, 1253–1260. doi: 10.2135/cropsci2010.06.0377
- Stulen, I., Perez-Soba, M., De Kok, L. J., and Van Der Eerden, L. (1998). Impact of gaseous nitrogen deposition on plant functioning. *New Phytol.* 139, 61–70. doi: 10.1046/j.1469-8137.1998.00179.x
- Takagi, M., and Gyokusen, K. (2004). Light and atmospheric pollution affect photosynthesis of street trees in urban environments. *Urban For. Urban Green.* 2, 167–171. doi: 10.1078/1618-8667-00033
- Takahashi, M., Furuhashi, T., Ishikawa, N., Horiguchi, G., Sakamoto, A., Tsukaya, H., et al. (2014). Nitrogen dioxide regulates organ growth by controlling cell proliferation and enlargement in *Arabidopsis*. *New Phytol.* 201, 1304–1315. doi: 10.1111/nph.12609
- Takahashi, M., Sakamoto, A., Ezura, H., and Morikawa, H. (2011). Prolonged exposure to atmospheric nitrogen dioxide increases fruit yield of tomato plants. *Plant Biotechnol.* 8, 485–487. doi: 10.5511/plantbiotechnology.11.0819a
- Takeuchi, Y. C., Nihira, J., Kondo, N., and Tezuka, T. (1985). Change in Nitrate-reducing activity in squash seedlings with NO<sub>2</sub> fumigation. *Plant Cell Physiol.* 26, 1027–1035.
- Touraine, B., Grignon, N., and Grignon, C. (1988). Charge balance in NO<sub>3</sub><sup>-</sup> fed soybean: estimation of K<sup>+</sup> and carboxylate recirculation. *Plant Physiol.* 88, 605–612. doi: 10.1104/pp.88.3.605
- Touraine, B., Muller, B., and Grignon, C. (1992). Effect of phloem-translocated malate on NO<sub>3</sub><sup>-</sup> uptake by roots of intact soybean plants. *Plant Physiol.* 99, 1118–1123. doi: 10.1104/pp.99.3.1118
- Uscola, M., Villar-Salvador, P., Oliet, J., and Warren, C. (2014). Foliar absorption and root translocation of nitrogen from different chemical forms in seedlings of two Mediterranean trees. *Environ. Exp. Bot.* 104, 34–43. doi: 10.1016/j.envexpbot.2014.03.004
- Vidmar, J. J., Zhuo, D., Siddiqi, M. Y., Schjoerring, J. K., Touraine, B., and Glass, A. D. M. (2000). Regulation of high-affinity nitrate transporter genes and high-affinity nitrate influx by nitrogen pools in roots of barley. *Plant Physiol.* 123, 307–318. doi: 10.1104/pp.123.1.307
- Wang, R. C., Okamoto, M., Xing, X., and Crawford, N. M. (2003). Microarray analysis of the nitrate response in *Arabidopsis* roots and shoots reveals over 1000 rapidly responding genes and new linkages to glucose, trehalose-6-phosphate, iron, and sulfate metabolism. *Plant Physiol.* 132, 556–567. doi: 10.1104/pp.103.021253
- Wardlaw, I. F. (2005). Viewpoint: consideration of apoplastic water in plant organs: a reminder. *Funct. Plant Biol.* 32, 561–569. doi: 10.1071/FP04127
- Weber, P., Nussbaum, S., Fuhrer, J., Gfeller, H., Schlunegger, U. P., Brunold, C., et al. (1995). Uptake of atmospheric <sup>15</sup>NO<sub>2</sub> and its incorporation into free amino acids in wheat (*Triticum aestivum* L.). *Physiol. Plant* 94, 71–77. doi: 10.1111/j.1399-3054.1995.tb00786.x
- Wellburn, A. R. (1990). Why are atmospheric oxides of nitrogen usually phytotoxic and not alternative fertilizers? *New Phytol.* 115, 395–429. doi: 10.1111/j.1469-8137.1990.tb00467.x
- Wellburn, A. R., Higginson, C., Robinson, D., and Walmsley, C. (1981). Biochemical explanations of more than additive inhibitory effects of low atmospheric levels of sulphur dioxide plus nitrogen dioxide upon plants. *New Phytol.* 88, 223–237. doi: 10.1111/j.1469-8137.1981.tb01719.x



- Wilkinson, J. Q., and Crawford, N. M. (1993). Identification and characterization of a chlorate-resistant mutant of *Arabidopsis thaliana* with mutations in both nitrate reductase structural genes NIA1 and NIA2. *Mol. Gen. Genet.* 239, 289–297.
- Wippel, K., Wittek, A., Hedrich, R., and Sauer, N. (2010). Inverse pH regulation of plant and fungal sucrose transporters: a mechanism to regulate competition for sucrose at the host/pathogen interface? *PLoS ONE* 5:e12429. doi: 10.1371/journal.pone.0012429
- Yoneyama, T., and Sasakawa, H. (1979). Transformation of atmospheric NO<sub>2</sub> absorbed in spinach leaves. *Plant Cell Physiol.* 20, 263–266.
- Yoneyama, T., Sasakawa, H., Ishizuka, S., and Totsuka, T. (1979). Absorption of atmospheric NO<sub>2</sub> by plants and soils. *Soil Sci. Plant Nutr.* 25, 267–275. doi: 10.1080/00380768.1979.10433167
- Yu, S. W., Li, L., and Shimazaki, K. I. (1988). Response of spinach and kidney bean plants to nitrogen dioxide. *Environ. Pollut.* 55, 1–13. doi: 10.1016/0269-7491(88)90155-8

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 28 March 2014; accepted: 04 July 2014; published online: 30 July 2014.

Citation: Hu Y, Fernández V and Ma L (2014) Nitrate transporters in leaves and their potential roles in foliar uptake of nitrogen dioxide. *Front. Plant Sci.* 5:360. doi: 10.3389/fpls.2014.00360

This article was submitted to *Crop Science and Horticulture*, a section of the journal *Frontiers in Plant Science*.

Copyright © 2014 Hu, Fernández and Ma. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.