



Protein based molecular markers provide reliable means to understand prokaryotic phylogeny and support Darwinian mode of evolution

Vaibhav Bhandari, Hafiz S. Naushad and Radhey S. Gupta*

Department of Biochemistry and Biomedical Sciences, McMaster University, Hamilton, ON, Canada

Edited by:

Yousef Abu Kwaik, University of Louisville School of Medicine, USA

Reviewed by:

Srinand Sreevastan, University of Minnesota, USA
Andrey P. Anisimov, State Research Center for Applied Microbiology and Biotechnology, Russia

*Correspondence:

Radhey S. Gupta, Department of Biochemistry and Biomedical Sciences, McMaster University, 1200 Main Street West, Health Sciences Center, Hamilton, ON L8N 3Z5, Canada.
e-mail: gupta@mcmaster.ca

The analyses of genome sequences have led to the proposal that lateral gene transfers (LGTs) among prokaryotes are so widespread that they disguise the interrelationships among these organisms. This has led to questioning of whether the Darwinian model of evolution is applicable to prokaryotic organisms. In this review, we discuss the usefulness of taxon-specific molecular markers such as conserved signature indels (CSIs) and conserved signature proteins (CSPs) for understanding the evolutionary relationships among prokaryotes and to assess the influence of LGTs on prokaryotic evolution. The analyses of genomic sequences have identified large numbers of CSIs and CSPs that are unique properties of different groups of prokaryotes ranging from phylum to genus levels. The species distribution patterns of these molecular signatures strongly support a tree-like vertical inheritance of the genes containing these molecular signatures that is consistent with phylogenetic trees. Recent detailed studies in this regard on the Thermotogae and Archaea, which are reviewed here, have identified large numbers of CSIs and CSPs that are specific for the species from these two taxa and a number of their major clades. The genetic changes responsible for these CSIs (and CSPs) initially likely occurred in the common ancestors of these taxa and then vertically transferred to various descendants. Although some CSIs and CSPs in unrelated groups of prokaryotes were identified, their small numbers and random occurrence has no apparent influence on the consistent tree-like branching pattern emerging from other markers. These results provide evidence that although LGT is an important evolutionary force, it does not mask the tree-like branching pattern of prokaryotes or understanding of their evolutionary relationships. The identified CSIs and CSPs also provide novel and highly specific means for identification of different groups of microbes and for taxonomical and biochemical studies.

Keywords: conserved indels, signature proteins, phylogenetic trees, lateral gene transfers, Thermotogae, Archaea, Crenarchaeota, RpoB signatures

INTRODUCTION

The understanding of prokaryotic relationships is one of the most important goals of evolutionary sciences. These relationships have been difficult to understand due to the simplicity and antiquity of prokaryotic organisms and disagreements in viewpoints among evolutionary biologists regarding the importance of different factors when grouping prokaryotes. Although earlier studies in this regard were based on morphology or physiology (Cowan, 1965; Buchanan and Gibbons, 1974; Stanier et al., 1976), the field itself has evolved to account for new information brought about by technological or informational breakthroughs, viz. molecular data, DNA hybridization and 16S rRNA (Zuckerandl and Pauling, 1965; Woese and Fox, 1977; Woese, 1987). The most recent breakthrough involves rapid and easily available sequencing of entire genomic sequences (Fleischmann et al., 1995; Iguchi et al., 2009; NCBI genomic database, 2012). This has allowed determination of evolutionary relationships among different organisms based upon large numbers of different

gene/protein sequences using a variety of approaches (Gupta, 1998; Haggerty et al., 2009; Puigbo et al., 2009; Blair and Murphy, 2011).

The comparative genomic analyses have revealed that phylogenetic relationships deduced based upon different genes and protein sequences are not congruent and lateral gene transfer (LGT) among different taxa is indicated as the main factor responsible for this lack of concordance (Gogarten et al., 2002; Baptiste and Boucher, 2008; Dagan et al., 2008; Puigbo et al., 2009; Swithers et al., 2009; Andam and Gogarten, 2011). This has led to questioning of whether the Darwinian model of evolution involving vertical inheritance of genes from parents to progenies (Darwin, 1859) is applicable to the prokaryotes (Doolittle, 1999; Pennisi, 1999; Gogarten et al., 2002; Dagan and Martin, 2006; Doolittle and Baptiste, 2007; Dagan et al., 2008; Baptiste et al., 2009; Williams et al., 2011). Multiple mechanisms are known to contribute to the evolution of an organism's genomes including genes that are acquired vertically from the parent organism,

evolution of new genes by gene duplication and divergence, gain of new genes by means of LGTs, as well as gene losses in various lineages (Baptiste et al., 2009; Ragan and Beiko, 2009; Treangen and Rocha, 2011; Williams et al., 2011). LGT, in particular, is being increasingly thought to have an overbearing influence on prokaryotic genome composition. Although rRNAs, ribosomal proteins and other genes involved in the information transfer processes are considered less prone to LGTs due to their involvement in complex gene networks (Jain et al., 1999; Sorek et al., 2007), recent studies indicate that no single gene/protein is completely immune to this process (Yap et al., 1999; Doolittle and Baptiste, 2007; Dagan et al., 2008). Some recent studies have estimated that over time most genes ($81 \pm 15\%$) have undergone at least one LGT event (Doolittle, 1999; Dagan and Martin, 2007; Doolittle and Baptiste, 2007; Dagan et al., 2008). These studies in large part form the basis of the hypothesis that LGTs have led to abolishment of all signals that can be used for determination of prokaryotic evolutionary relationships and a call for uprooting the tree of life (Martin, 1999; Pennisi, 1999; Doolittle, 2000; Gogarten et al., 2002; Delsuc et al., 2005; Baptiste et al., 2009).

Although the importance of LGTs in genome evolution is widely accepted, there is considerable disagreement concerning the prevalence of LGTs and their impact on prokaryotic evolutionary relationships. While some authors have indicated that LGT is so profuse that its influence disguises the Darwinian mode of evolution involving vertical inheritance of genes (Gogarten et al., 2002; Baptiste et al., 2005b, 2009; Doolittle and Baptiste, 2007; Koonin, 2007), others have inferred that the incidences of LGTs are either very minimal or limited and those genes that are laterally transferred have little impact on prokaryotic phylogeny (Wolf et al., 2002; Kurland et al., 2003; Dutilh et al., 2004; Beiko et al., 2005; Kunin et al., 2005; Kurland, 2005; Galtier, 2007; Puigbo et al., 2009; Gao and Gupta, 2012a). However, there are no standardized methods to assess LGTs and the methods used to infer LGTs are varied and based upon large numbers of often poorly supported assumptions (Koski and Golding, 2001; Koski et al., 2001; Ragan, 2001; Beiko et al., 2005; Boto, 2010). Thus, the prevalence of LGTs differ greatly among different studies and often similar datasets have led to dissimilar conclusions (Koski et al., 2001; Ragan, 2001; Wang, 2001; Lerat et al., 2003; Susko et al., 2006; Zhaxybayeva et al., 2007; Marri and Golding, 2008; Roettger et al., 2009). Therefore, prior to concluding that in view of LGTs the Darwinian mode of evolution is not a suitable model for prokaryotes, reliability of the incidences of LGTs and their overall impact on the evolutionary relationships should be critically examined.

Despite the prevalence of LGTs, phylogenetic trees based upon 16S rRNA as well as numerous single genes as well multi-gene analyses strongly support the existence of large numbers of distinct phyla of bacteria (Ludwig and Klenk, 2005). Additionally, these trees also clearly delineate many discrete taxonomic clades within these phyla (Woese, 1987; Ludwig and Klenk, 2005; Ciccarelli et al., 2006; Wu et al., 2009; Gao and Gupta, 2012a). In a recent detailed study Puigbo et al. (2009) reported construction of phylogenetic trees for 6901 prokaryotic genes. Although there were significant topological differences among these trees,

a consistent phylogenetic signal was observed in most of these trees, indicating that the LGT events, which were of random nature, did not obscure the central trend resulting from the vertical transfer of genes. The fact that similar prokaryotic clades at different taxonomic levels (ranging from phyla to genera) are consistently identified in phylogenetic trees based upon different gene/protein sequences strongly indicates that the distinctness of the prokaryotic taxa and their evolutionary relationships are in large part discernible and they have not been obliterated by LGTs (Woese, 1987; Daubin et al., 2002; Kurland et al., 2003; Lerat et al., 2003; Beiko et al., 2005; Kurland, 2005; Ludwig and Klenk, 2005; Ciccarelli et al., 2006; Ragan and Beiko, 2009; Wu et al., 2009; Boto, 2010; Yarza et al., 2010; Gupta, 2010b; Gao and Gupta, 2012a). To account for the above observations and the occurrences of LGTs, it has been suggested that the prokaryotic evolution has both tree-like (at intermediate phylogenetic depths) and non-tree (or net-like) (at the base and tips) characteristics (Dagan et al., 2008; Puigbo et al., 2009, 2010; Swithers et al., 2009; Boto, 2010; Beiko, 2011; Dagan, 2011; Kloesges et al., 2011; Popa et al., 2011).

The availability of genome sequences is also enabling development of novel and independent sequence based approaches for determining the evolutionary relationships among organisms and to assess the impact of LGTs on these relationships. In this review, we provide a summary of our recent work in this area based upon two different types of molecular markers that we have used successfully for understanding the evolutionary relationships among prokaryotes. Based upon these markers it is now possible to identify different prokaryotic taxa ranging from phyla to genera in clear molecular terms and the evolutionary relationships among them can also be reliably deduced (Gupta and Griffiths, 2002; Gupta, 2009, 2010a; Gao and Gupta, 2012b). The relationships revealed by these new approaches strongly support a tree-like branching pattern among prokaryotes and the observed incidences of LGTs, which exhibit no specific pattern or statistical significance, apparently have no major impact on the derived relationships. It is contended that these molecular markers provide valuable means for developing a reliable phylogeny and taxonomy of the prokaryotic organisms.

USEFULNESS OF CONSERVED SIGNATURE INDELS (CSIs) AND CONSERVED SIGNATURE PROTEINS (CSPs) FOR UNDERSTANDING EVOLUTIONARY RELATIONSHIPS AMONG PROKARYOTES

Of the two kinds of molecular markers that we are using for studying prokaryotic evolution, the conserved signature indels (inserts or deletions), or CSIs, in protein sequences comprises an important category (Gupta, 1998, 2010a; Griffiths and Gupta, 2001). The CSIs that provide useful molecular markers for evolutionary studies are generally of the same lengths and they are flanked on both sides by conserved regions to ensure that the observed changes are not caused by alignment artifacts (Gupta, 1998; Gupta and Griffiths, 2002; Jordan and Goldman, 2012). When such CSIs are present in the same position in a given protein in a group of related species, their presence is most parsimoniously explained by postulating that the genetic change leading to the CSI occurred in a common ancestor of this group

and then this gene with the indel was vertically transmitted to its progeny (Rivera and Lake, 1992; Baldauf and Palmer, 1993; Gupta, 1998, 2000b; Rokas and Holland, 2000; Cutino-Jimenez et al., 2010). The CSIs that are uniquely shared by organisms of one taxa provide molecular tools for identifying the species from this taxa and consolidating the relationships among bacteria of that taxa by delimiting it in molecular terms (Gupta, 2004). Additionally, depending upon the presence or absence of a given CSI in the outgroup species, it can be determined whether the indel represents an insert or a deletion and based upon this a rooted relationship among the species of interest can be derived. Our earlier work in this regard has led to identification of large numbers of CSIs that are specific for different groups of microbes at various phylogenetic levels (**Table 1**; Gupta and Griffiths, 2006; Gupta, 2009; Gupta and Bhandari, 2011; Gupta and Shami, 2011; Gao and Gupta, 2012b).

The second kind of molecular markers that we have usefully employed in our systematic and evolutionary studies are whole proteins that are uniquely found in particular groups or subgroups of bacteria (Gupta, 2006; Gupta and Griffiths, 2006; Gupta and Mok, 2007; Gao and Gupta, 2012b). Comparative analyses of genomic sequences have indicated that many conserved proteins are uniquely present in all species from particular groups, at different phylogenetic depths (Daubin and Ochman, 2004; Lerat et al., 2005; Gupta, 2006; Gupta and Griffiths, 2006; Gupta and Mok, 2007; Dutilh et al., 2008; Gao and Gupta, 2012b). Because of their unique presence in species from particular phylogenetic clades of species, it is likely that the genes for these CSPs originated once in a common ancestor of these groups and then vertically acquired by all its descendants. Because of their taxa specificity these CSPs again provide valuable molecular markers for identifying different groups of species in molecular terms and for evolutionary studies (Gao and Gupta, 2007; Gupta and Mathews, 2010; Gupta, 2010b). However, when a CSP (or CSI) is confined to certain species/strains, then based upon this information alone, it is often difficult to determine whether these species form a clade in the phylogenetic sense or not. Hence, to understand the evolutionary significance of these signatures, such studies are generally performed in conjunction with phylogenetic analysis, which provides a reference point for evaluating the significance of various CSIs and CSPs (Gao and Gupta, 2007; Gupta and Mathews, 2010; Gupta, 2010b).

Molecular markers in the form of CSIs and CSPs have proven useful for examining or consolidating prokaryotic relationships at domain, phylum as well as intra-phylum levels. **Table 1** provides a summary of some bacterial and archaeal taxa for which CSIs and CSPs have been identified (Gupta, 2010a). Two recent detailed studies based upon CSIs and CSPs have focused upon understanding evolutionary relationships within the phylum Thermotogae and the domain Archaea (Gao and Gupta, 2007; Gupta and Bhandari, 2011; Gupta and Shami, 2011). To illustrate the usefulness of these molecular markers for elucidation of prokaryotic evolutionary relationships, and to assess the influence of LGTs on the derived inferences, results for these two taxonomic groups are reviewed here.

MOLECULAR MARKERS FOR THE THERMOTOGAE


The species of the phylum Thermotogae are a group of hyperthermophilic, anaerobic, gram-negative bacteria recognized by a distinctive toga-like sheath structure and their ability to grow at high temperatures (Huber et al., 1986). The approximately 90 species of this phylum are currently divided into nine Genera within a single family termed the Thermotogaceae (Euzéby, 2011; NCBI Taxonomy, 2012). The Thermotogae species, prospectively, are important tools for industrial and biotechnological applications due to the ecological niche they inhabit and the thermo-stable proteins that they harbor (Connors et al., 2006). With the publication of the genome for *T. maritima*, the first species from this phylum (Nelson et al., 1999), the Thermotogae were brought to the forefront of LGT debate. This was due to the fact that based upon Blast searches it was determined that for about 25% of the genes from *T. maritima* genome, the closest blast hits were from archaeal species rather than any bacteria, leading to the inference that Thermotogae species have incurred high degree of LGTs with the archaeal organisms (Nelson et al., 1999). Upon revisiting this issue, Zhaxybayeva et al. (2009) found that for only about 11% of the Thermotogae proteins Archaea were the closest hits, but that the Thermotogae proteins exhibited maximal similarity (42–48% of genes) to the Firmicutes. Based upon these observations, the Thermotogae species genomes were proposed to be a chimera composed of different bacterial and archaeal sources (Zhaxybayeva et al., 2009). However, these estimates for LGTs have been questioned in other studies which indicate that much less (6–7%) of the Thermotogae genome has been laterally transferred (Garcia-Vallve et al., 2000; Ochman et al., 2000). Further, in view of the fact that Thermotogae species branch in proximity of the Firmicutes phylum (Gupta, 2001; Griffiths and Gupta, 2004b), the observation that a preponderance of the top hits for the Thermotogae species are from Firmicutes is an expected result, and it does not indicate that these genes have been laterally transferred (Zhaxybayeva et al., 2009; Andam and Gogarten, 2011).

Apart from their unique protein toga, the species of the phylum Thermotogae are assigned to this group and divided into its different genera primarily on the basis of their branching in the 16S rRNA trees (Reysenbach, 2001; Huber and Hannig, 2006; Zhaxybayeva et al., 2009; Yarza et al., 2010). Until recently, no unique molecular or biochemical characteristics were known that could distinguish the species of this phylum from other bacteria. For identification of molecular markers that could possibly define this phylum and its sub-taxa, a genome wide analysis was performed on protein sequences from 12 Thermotogae spp. whose genomes were available (Gupta and Bhandari, 2011). The protein sequences from these 12 species as well as species representing other bacteria phyla were aligned and examined for the presence of CSIs that were uniquely present in Thermotogae species or those that were commonly shared with some other bacteria. The analysis identified numerous CSIs specific for all Thermotogae. An example of a CSI consisting of a 3 aa long insert in the ribosomal protein L7 that is exclusively present in all sequenced Thermotogae species, including two recently sequenced species, is shown in **Figure 1A**. The unique presence of this CSI of the same length, at the same position in

Table 1 | Overview of the CSIs and CSPs that have been identified for some major prokaryotic taxa.

Taxonomic group	Number of CSPs/CSIs	References
Archaea	<i>Archaeal Kingdom specific:</i> 16 CSPs <i>Subgroups:</i> Thaumarchaeota—6 CSIs/201 CSPs, Euryarchaeota—6 CSPs, Thermoacidophiles—77 CSPs, Halophiles—127 CSPs, Methanogens—31 CSPs, Thermococcus-Pyrococcus clade—141 CSPs	Gao and Gupta, 2007; Gupta and Shami, 2011
Crenarchaeota	<i>Phylum specific:</i> 6 CSIs, 13 CSPs <i>Subgroups:</i> Sulfolobales—3 CSIs/151 CSPs, Thermoproteales—5 CSIs/25 CSPs, Desulfurococcales—4CSPs, Sulfolobales-Desulfurococcales clade—2 CSIs/18 CSPs	Gupta and Shami, 2011
Thaumarchaeota	>200 CSPs	Gupta and Shami, 2011
Thermotogae	<i>Phylum specific:</i> 18 CSIs <i>Subgroups:</i> Thermotoga genus—13 CSIs, Thermosipho genus—7 CSIs, Thermosipho-Fervidobacterium clade—13 CSIs, Thermotoga-Thermosipho-Fervidobacterium clade—5 CSIs, Petrotoga-Kosmotoga clade—4 CSIs	Gupta and Bhandari, 2011
Cyanobacteria	<i>Phylum specific:</i> 39 CSPs/10 CSIs <i>Subgroups:</i> Cyanobacterial Clade A—14 CSPs/1 CSI, Other Cyanobacteria (outside clade A)—5 CSPs/4 CSIs, Cyanobacterial Clade C—60 CSPs, Nostocales—65 CSPs, Chroococcales—8 CSPs, <i>Synechococcus</i> —14 CSPs, <i>Prochlorococcus</i> —19 CSPs, Low B/A type <i>Prochlorococcus</i> —67 CSPs	Gupta, 2009; Gupta and Mathews, 2010
Chlamydiae	<i>Phylum specific:</i> 59 CSPs/8 CSIs <i>Subgroups:</i> Chlamydiaceae—79 CSPs, Chlamydochlamydia—20 CSPs, Chlamydia—20 CSPs	Gupta and Griffiths, 2006
Bacteroidetes, chlorobi and fibrobacteres	<i>Phylum specific:</i> 1 CSP/2 CSIs <i>Subgroup specific:</i> Bacteroidetes—27 CSPs/2 CSIs, Chlorobi—51 CSPs/2 CSIs, Bacteroidetes and Chlorobi clade—5 CSPs/3CSIs	Gupta, 2004
Actinobacteria	<i>Phylum specific:</i> 24 CSPs/4 CSIs <i>Subgroup specific:</i> CMN group—13 CSPs, <i>Mycobacterium</i> and <i>Nocardia</i> —14 CSIs, <i>Mycobacterium</i> —24 CSPs, <i>Micrococcineae</i> —24 CSPs, Corynebacteriales—4 CSPs/2 CSIs, Bifidobacteriales—14 CSPs/1 CSI	Gao and Gupta, 2005, 2012b; Gao et al., 2006
Deinococcus-thermus	<i>Phylum specific:</i> 65 CSPs/8 CSIs <i>Subgroup specific:</i> Deinococci—206 SPs	Griffiths and Gupta, 2004a, 2007a
Aquificae	<i>Phylum specific:</i> 10 CSPs/5 CSIs	Griffiths and Gupta, 2006b, 2004b
α-proteobacteria	<i>Class specific:</i> 6 CSPs/13 CSIs <i>Subgroups:</i> Rickettsiales—3 CSPs/2 CSIs, Rickettsiaceae—4 CSPs/5 CSIs, Anaplasmataceae—5 CSPs/2 CSIs, Rhodobacterales-Caulobacter-Rhizobiales clade—2 CSIs, Rhodobacterales-Caulobacter clade—1 CSI, Rhizobiales—6 CSPs/1CSI, Bradyrhizobiaceae—62 CSPs/2CSIs	Gupta and Mok, 2007
γ-proteobacteria	<i>Class specific:</i> 4 CSPs/1 CSI <i>Subgroups:</i> 20 CSPs, 2 CSIs for various subgroup combinations of subgroups	Gao et al., 2009
ε-proteobacteria	<i>Class specific:</i> 49 CSPs/4 CSIs <i>Subgroups:</i> <i>Wolinella-Helicobacter</i> clade—11 CSPs/2 CSIs, <i>Campylobacter</i> genus—18 CSPs/1 CSI	Gupta, 2006
Pasteurellales	<i>Order specific:</i> 44 CSIs <i>Subgroups:</i> Pasteurellales Clade I—13 CSIs, Pasteurellales Clade II—9 CSIs	Naushad and Gupta, 2012
Clostridia sensu stricto	<i>Genus specific:</i> 10 CSPs/3 CSIs	Gupta and Gao, 2009

The table provides general information regarding the number of CSIs and CSPs identified for many taxonomic groups on which genomic studies have been conducted. Further details can be obtained from the corresponding studies.



this universally distributed protein, in different species from the phylum Thermotogae indicates that the genetic change leading to this CSI occurred once in the common ancestor of the Thermotogae species. In addition to this CSI, this study also identified 17 other CSIs in other important proteins such as DNA recombination protein RecA, DNA polymerase I and tryptophanyl-tRNA synthetase that are also specific for the species from the phylum Thermotogae (Gupta and Bhandari, 2011).

In addition to the large numbers of CSIs that were uniquely present in all Thermotogae species, this study also identified many CSIs that were specific for different sub-groups within the phylum Thermotogae (Gupta and Bhandari, 2011). These included 13 CSIs that were specific for the species of the genus

Thermotoga and seven others that distinguished species of the genus *Thermosiphonia* from all others. However, it was observed that the species *Thermotoga lettingae* shared only 1 of 13 CSIs that were otherwise commonly present in other species of this genus. This suggests that *T. lettingae*, which is distantly related to all other *Thermotoga* species, should be assigned to a separate genus. Besides these CSIs that were specific for the species of these two genera, 13 CSIs supported a specific relationships among species of the *Fervidobacterium* and *Thermosiphonia* genera; 5 CSIs were shared by species from the genus *Thermotoga* and those from the *Fervidobacterium-Thermosiphonia* clade; and 4 CSIs supported a grouping of the *Petrotoga* and *Kosmotoga* genera along with the species *Thermotogales bacterium MesG1.Ag.4.2* (Figure 1C, left panel; Gupta and Bhandari, 2011). Importantly, all of the

relationships indicated by various CSIs were also independently observed in a phylogenetic tree for the Thermotogae species based upon concatenated sequences for 12 conserved proteins (**Figure 1B**).


The CSIs identified in the above study independently and strongly supported different nodes observed in the phylogenetic tree for Thermotogae species all the way from phylum to genus level. If the hypothesis that LGT events have abolished the ability to discern prokaryotic relationships was correct, then it should have been difficult to identify discrete molecular markers supporting distant relationships among these species. At the very least, the Thermotogae species would have shown relationships with species of other prokaryotic groups such as Firmicutes or Archaea as frequently as they did with one another. In this study, in addition to the CSIs that were specific for the Thermotogae species (**Figure 1C**, left panel), several CSIs were also identified that the Thermotogae shared with species from other prokaryotic or eukaryotic organisms (**Figure 1C**, right panel). However, such CSIs, suggesting possible LGT between Thermotogae and other taxa, were far outweighed by CSIs supporting the monophyletic, tree-like relationships among the species of the phylum (left panel) (Gupta and Bhandari, 2011). Assuming that all the CSIs that the Thermotogae shared with other groups are due to LGT, less than 20% (16 of 85) of all Thermotogae genes containing these CSIs have incurred LGTs (Gupta and Bhandari, 2011). Moreover, these presumed LGT events are of random nature and in no case do the Thermotogae species share more than a total of 3 CSIs with any particular phyla of species. Additionally, in most of these cases only a few species from these other taxa contained the indels that were present in most or all Thermotogae species (Gupta and Bhandari, 2011). Thus, these other CSIs, although they are present in a few isolated species from other taxa, are also largely specific for the Thermotogae species and they do not affect the ability of other CSIs to clearly discriminate Thermotogae species from all other bacteria or to deduce the evolutionary relationships amongst species from this phylum.

The shared presence of similar CSI in unrelated taxa can result from two different possibilities, either the gene with the CSI was laterally transferred among the two groups or that independent CSIs owing to two separate genetic events are responsible for these CSIs. After identification of such CSIs, tree-making approaches can be used to test if the presence of the indel in the two groups is due to LGT. Previously, in our work, a number of CSIs in the GlyA and MurA proteins that were commonly shared by the Chlamydiae and a subgroup of Actinobacteria were shown to be due to lateral transfer of genes from Actinobacteria to a common ancestor of the Chlamydiae (Griffiths and Gupta, 2006a). Recently, the shared presence of several CSIs in the bacteriochlorophyll biosynthesis proteins by unrelated phyla of photosynthetic prokaryotes has also been shown to be due to LGTs (Raymond et al., 2002; Gupta, 2012). However, in many other instances phylogenetic analyses have not supported LGT as the possible reason for the presence of a related CSI in unrelated taxa. In these cases, similar CSIs have originated independently in these lineages due to their presumed similar functions in these particular taxa.

MOLECULAR MARKERS FOR THE ARCHAEA AND ITS SUB-GROUPS

Archaea are widely recognized as the third domain of life. They generally inhabit extreme environments such as those of extreme temperature, pH or salinity, where little to no other life exists (Woese et al., 1990). However, recent studies indicate that archaeal species are widespread in the environment and they play a major role in the carbon and nitrogen cycles (Pace, 1997; Herndl et al., 2005; Leininger et al., 2006). Some archaeal species have been found to be commensal organisms residing in human colons (Oxley et al., 2010). The Archaea are generally divided into two main phyla, the Crenarchaeota and Euryarchaeota, based on 16S rRNA data and other phylogenetic data (Woese et al., 1990; Gribaldo and Brochier-Armanet, 2006). The Crenarchaeotes are described as thermophiles with sulfur-reducing capabilities while the Euryarchaeotes are metabolically and morphologically quite diverse (Gribaldo and Brochier-Armanet, 2006; Gupta and Shami, 2011). The mesophilic Crenarchaeota have been recently placed into a separate phylum called the Thaumarchaeota (Brochier-Armanet et al., 2008; Gupta and Shami, 2011).

Despite the importance of Archaea in different environments and in understanding of the evolutionary history of life on earth (Woese et al., 1990; Gupta, 2000a), until recently, very few molecular characteristics were known that are uniquely shared by all Archaea. Additionally, as the higher taxonomic groups within Archaea are described primarily based upon 16S rRNA trees, the characteristics that are unique to different phyla, classes, orders and families of the Archaea have scarcely been elucidated (Boone et al., 2001). The utilization of archaeal genomes for discovery of CSPs as well as CSIs has provided significant information in the form of molecular markers that are distinctive characteristics of Archaea and its taxonomic sub-groups. In 2007, a comprehensive analysis was performed on available archaeal genomes to search for CSPs that were unique to either all Archaea or many of its sub-groups (Gao and Gupta, 2007). Over 1400 such proteins distinctive of Archaea or its main taxa were discovered (**Figure 2**). In the analysis, sixteen proteins specific to all or most Archaea were identified that were not present in any bacterial or eukaryotic organism. Numerous proteins whose homologs were limited to the Crenarchaeota, Euryarchaeota and other sub-groups such as the Thermococci, Thermoplasmata, and Halobacteriales were also detected (**Figure 2**). Significantly, this study also identified 31 proteins that were commonly shared by all methanogenic bacteria (Gao and Gupta, 2007). In the 16S rRNA and other phylogenetic trees, the methanogenic Archaea do not form a monophyletic lineage, but instead are split into a number of distinct clusters separated by non-methanogenic Archaea (Burggraf et al., 1991; Brochier et al., 2004; Bapteste et al., 2005a; Gao and Gupta, 2007). Because most of the proteins that are commonly shared by various methanogens are generally involved in functions related to methanogenesis and their genes are clustered into a few large operons in genomes (Harms et al., 1995; Tersteegen and Hedderich, 1999; Grabarse et al., 2001; Gao and Gupta, 2007), it is likely that the genes for these proteins have been laterally acquired by different Archaea. This could provide a plausible explanation for the observed discrepancy in the branching of methanogenic Archaea in phylogenetic trees and



their unique sharing of genes for these proteins (Gao and Gupta, 2007).

A recent analysis has further added to the catalogue of molecular signatures for the archaeal organisms (Gupta and Shami, 2011). The focus of this study was on identifying CSIs and CSPs that were specific for the Crenarchaeota and Thaumarchaeota phyla (Gupta and Shami, 2011). Six CSIs and 13 CSPs specific for all species of the phylum Crenarchaeota were identified along with numerous markers for its different orders: the Sulfolobales (151 CSPs, 3 CSIs), Thermoproteales (25 CSPs, 5 CSIs) and the Desulfurococcales (4 CSPs). The study also described the markers (18 CSPs and 2 CSIs) indicative of a close relationship among the Sulfolobales and the Desulfurococcales. The discriminative ability of CSPs is highlighted by the results of blast searches on some CSPs that are specific for the Crenarchaeota or its main groups (Sulfolobales, Thermoproteales, Desulfurococcales and Acidilobales) that are shown in **Table 2**. In these cases, BLASTP searches were carried out on these proteins and the results for all species for whom the observed *E*-values were significant are shown. From the results presented in **Table 2**, it is evident that the first 2 CSPs are specific for the Crenarchaeota phylum, the next two are uniquely found in various species belonging to the orders Desulfurococcales, Acidilobales and Sulfolobales, whereas the last 5 CSPs are distinctive characteristics of species belonging to either

the Desulfurococcales (and Acidilobales), the Sulfolobales, or the Thermoproteales orders.

In this study, more than 200 CSPs for various members of the newly defined Thaumarchaeota phylum were also identified (Gupta and Shami, 2011). The Thaumarchaeota are composed of several organisms previously included in the Crenarchaeota (Brochier-Armanet et al., 2008). The two phyla appear as sister groups in phylogenetic analysis and they also share 3 CSIs and 10 CSPs with each other (Gupta and Shami, 2011). Nevertheless, the two groups can be phylogenetically differentiated and numerous markers have been identified for each group that helps to define them molecularly as individual taxa (Gupta and Shami, 2011). A summary diagram depicting the various molecular markers specific for the archaeal species is shown in **Figure 2**. It should be noted that CSIs were only identified for the Thaumarchaeota and the Crenarchaeota and no detailed analysis to identify CSIs has thus far been carried out on the Euryarchaeota.

The two studies noted above have identified numerous CSIs and CSPs for the Archaea, its main phyla (Euryarchaeota, Crenarchaeota, Thaumarchaeota) and a number of its sub-phylum level taxa (Sulfolobales, Thermococcales, Halobacteriales, etc.; Gao and Gupta, 2007; Gupta and Shami, 2011). Except for the methanogens, the distribution patterns of the identified CSIs and CSPs are also strongly supported by the phylogenetic

Table 2 | A series of proteins specific for the Crenarchaeota and its sub-groups.

Protein accession #	NP_147640	NP_147284	BAA81469	NP_147588	YP_001041009	YP_254810	YP_254922	NP_559041	NP_559897
Protein length	262 aa	143 aa	98 aa	228 aa	127 aa	228 aa	270 aa	626 aa	113 aa
Desulfurococcales									
<i>Aeropyrum pernix</i>	0.0	9e-98	5e-64	7e-161	7e-22	-	-	-	-
<i>Hyperthermus butylicus</i>	3e-46	9e-43	1e-20	1e-23	3e-25	-	-	-	-
<i>Ignicoccus hospitalis</i>	3e-41	-	5e-27	4e-19	3e-25	-	-	-	-
<i>Desulfurococcus kamchatkensis</i>	7e-46	1e-21	2e-20	5e-17	7e-32	-	-	-	-
<i>Staphylothermus marinus</i>	4e-56	1e-25	3e-21	3e-21	2e-85	-	-	-	-
Acidilobales									
<i>Acidilobus saccharovorans</i>	9e-56	4e-36	4e-21	1e-46	1e-19	-	-	-	-
Sulfolobales									
<i>Sulfolobus tokodaii</i>	4e-40	2e-29	3e-20	7e-26	-	1e-77	1e-80	-	-
<i>Sulfolobus islandicus</i>	4e-42	6e-30	1e-25	1e-15	-	7e-50	8e-65	-	-
<i>Sulfolobus acidocaldarius</i>	7e-34	3e-23	4e-22	4e-24	-	2e-162	0.0	-	-
<i>Sulfolobus solfataricus</i>	1e-41	7e-30	5e-26	8e-15	-	5e-50	8e-64	-	-
<i>Metallosphaera sedula</i>	3e-31	3e-33	3e-20	1e-22	-	4e-39	8e-60	-	-
Thermoproteales									
<i>Pyrobaculum aerophilum</i>	9e-18	3e-11	-	-	-	-	-	0.0	2e-73
<i>Pyrobaculum islandicum</i>	3e-18	3e-11	-	-	-	-	-	0.0	6e-54
<i>Pyrobaculum arsenaticum</i>	1e-18	1e-10	-	-	-	-	-	0.0	2e-63
<i>Pyrobaculum caldifontis</i>	6e-22	7e-11	-	-	-	-	-	0.0	1e-60
<i>Thermofilum pendens</i>	1e-35	5e-30	-	-	-	-	-	1e-42	3e-10
<i>Caldivirga maquilingensis</i>	1e-17	4e-8	-	-	-	-	-	1e-87	2e-22
<i>Thermoproteus neutrophilus</i>	2e-19	7e-11	-	-	-	-	-	0.0	5e-61
<i>Thermoproteus tenax</i>	3e-15	6e-10	-	-	-	-	-	0.0	4e-46
Top non-Crenarchaeota hit									
<i>Brucella melitensis</i>	(2e-1)	<i>Desulfobacterium autotrophicum</i>	<i>Aromatoleum aromaticum</i>	<i>Serpula lacrymans</i>	<i>Clonorchis sinensis</i>	<i>Granulicatella elegans</i>	<i>Encephali-tozoon cuniculi</i>	<i>Burkholderia cenocepacia</i>	<i>Sordaria macrospora</i>
		(8e-1)	(4e-1)	(7e-1)	(3e-1)	(6e-1)	(7e-1)	(9e-1)	(1e-1)

BlastP searches were carried out on proteins specific for the Crenarchaeota or its sub-groups and the results for representative species from different sub-groups of the Crenarchaeota are shown with the observed E-values. E-values greater than 1e-3 are considered insignificant hits with lack of homology to the query protein sequence.

The dashes (-) indicate that the homolog for the protein query was not detected in the BlastP searches.

Top non-Crenarchaeota hits indicate detection of species outside the Crenarchaeota that were observed to have the lowest E-value scores.

shared by all of the sequenced species belonging to the phyla Proteobacteria (different subclasses), Aquificae, Chlamydiae, Verrucomicrobiae, Bacteroidetes-Chlorobi, and Planctomycetes (Griffiths and Gupta, 2007b). This insert is present in all of the >1500 sequences that are available from species from these phyla. On the other hand, this CSI is not found in any of the >1500 sequences available from various species belonging to the phyla Firmicutes, Actinobacteria, Chloroflexi, Cyanobacteria, Deinococcus-Thermus, Synergistetes, Spirochaetes, etc. This insert is also not found in the archaeal RpoB homologs, thus providing evidence that this indel is an insert in the groups of species where it is found (Griffiths and Gupta, 2004b). Based upon its highly specific species distribution pattern, which argues strongly against the lateral transfer of this gene amongst various phyla, the genetic change responsible for this CSI most likely occurred in a common ancestor of the group of species that contain this CSI, after the divergence of other bacterial phyla that lack this indel as indicated in **Figure 3A** (right panel). A number of other main-line CSIs, which based upon their species distribution patterns have occurred at other important branch points in prokaryotic evolution, have been described in our earlier works (Griffiths and Gupta, 2001, 2004b; Gupta and Griffiths, 2002). Based upon these CSIs, it is possible to determine the branching order of most of the bacterial phyla (Gupta, 1998, 2001, 2003; Griffiths and Gupta, 2004b; see also www.bacterialphylogeny.info).

Within the highly conserved RpoB protein, in addition to the large CSI that is commonly shared by a number of bacterial phyla, several other CSIs have been identified that are specific for different groups/phyla of bacteria. The taxon specificities of these CSIs and their positions within in the RpoB polypeptide are shown in **Figure 3B**. These CSIs include a 4 aa deletion that is commonly and uniquely shared by a number of different orders of the γ -proteobacteria (399/399 species), a 3 aa insert that is specifically present in all of the Chlamydiae-Verrucomicrobiae species (47/47), another 3 aa insert that is a distinctive property of the Clade C cyanobacteria (50/50; Gupta, 2009), a 25 aa insert in various species from the order Rhodospirillales (103/103) and a 6 aa insert in all species from the genus *Thermotoga* except *T. lettingae* (Gupta and Griffiths, 2006; Gupta and Mok, 2007; Griffiths and Gupta, 2007b; Gao et al., 2009; Gupta and Bhandari, 2011). It is highly significant that within a single gene/protein multiple highly specific CSIs are present, each of which is specific for a different group of bacteria and help distinguish these groups from all other bacteria. These CSIs are not present in any species outside of the indicated taxa. The presence of these different taxon-specific characteristics in a single gene/protein strongly indicates that the genetic changes responsible for these CSIs occurred in the gene for this key protein at different stages in the evolution of bacterial domain and that no LGT of the gene for the RpoB protein has occurred among these taxa. Similar to the RpoB protein, multiple CSIs that are specific for different groups of prokaryotes have also been identified in many other important genes/proteins. These observations indicate that strong and consistent phylogenetic signals that are very likely not affected to any significant extent by the LGTs are still present in many conserved and universally distributed genes/proteins and these can be used to trace the evolutionary relationships among prokaryotes.

It is important to point out that virtually all of the higher taxonomic clades (above the Genus rank) within prokaryotes are currently identified solely on the basis of their branching in the 16S rRNA trees. Because the phylogenetic trees are a continuum, based upon them it has proven difficult to clearly define or delimit the boundaries of different taxonomic groups. Additionally, for virtually all of the higher prokaryotic taxa, no molecular, biochemical or physiological characteristics are known that are unique to them. Hence, a very important aspect of microbiology that needs to be understood is that in what respects do species from different main groups of bacteria differ from each other and what, if any, unique molecular, biochemical, structural or physiological characteristics are commonly shared by species from different groups? In this context, the large numbers of CSIs and CSPs for different taxonomic clades of bacteria that are being discovered by comparative genomic analyses provide novel and valuable tools for taxonomic, diagnostic, and biochemical studies (Gupta and Bhandari, 2011; Gao and Gupta, 2012b). In view of the specificities of the discovered CSIs and CSPs for different groups of prokaryotes and their retention by all species from these groups of prokaryotes, it is highly likely that these CSIs and CSPs are involved in functions that are essential for prokaryotes (Galperin and Koonin, 2004; Fang et al., 2005; Singh and Gupta, 2009; Schoeffler et al., 2010). Indeed, recent work on several CSIs have shown that they are essential for the group of organisms where they are found and the deletion or substantial changes in them led to failure of cell growth (Singh and Gupta, 2009; Schoeffler et al., 2010). Hence, further studies on understanding the cellular functions of the different taxa-specific CSIs and CSPs could lead to identification of novel biochemical and other functional characteristics that are specific for these groups of organisms.

It should also be noted that the identified CSIs and CSPs generally constitute robust molecular characteristics that exhibit high degree of predictive ability. Many of these CSIs and CSPs were discovered when the sequence information was available for very few prokaryotic species. However, despite the large increase in the number of sequenced genomes, most of these CSIs and CSPs are still specific for the originally indicated groups of prokaryotes (Gupta, 2009, 2011; Gao and Gupta, 2012b). Additionally, for several Chlamydiae-, Aquificae-, Deinococcus-Thermus- and Actinobacteria- specific degenerate primers based on conserved flanking sequences have been designed and they have been used to amplify the sequence regions predicted to contain the CSIs from large numbers of organisms for whom no sequences were available (Griffiths and Gupta, 2004a,b; Gao and Gupta, 2005; Griffiths et al., 2005). In these studies, in almost all cases the expected inserts or deletions were found to be present in previously un-sequenced organisms from the indicated groups, thus providing evidence that these CSIs and CSPs provide powerful new tools for identification of both known as well as novel species from different groups of prokaryotes.

CONCLUSIONS

There is considerable debate at present concerning the impact of LGTs on understanding prokaryotic phylogeny. While there

is little dispute that LGT plays an important role in microbial evolution, the extreme view taken by some that LGTs are so rampant within the prokaryotes that it totally masks the evolutionary signal from vertical transfer of genes (Doolittle, 2000; Gogarten et al., 2002; Doolittle and Bapteste, 2007; Dagan et al., 2008; Bapteste et al., 2009) is not supported by available evidence. As reviewed here, in phylogenetic trees based upon most gene/protein sequences all of the major groups within prokaryotes (from phylum down to genus level) are generally clearly identified, thus indicating that a strong phylogenetic signal emanating from vertical transfer of genes is maintained throughout prokaryotic evolution (Gupta, 1998, 2000b; Dutilh et al., 2004; Ludwig and Klenk, 2005; Ciccarelli et al., 2006; Puigbo et al., 2009). Most of the differences seen amongst these trees are either at the tips (i.e., species/strains levels) or at the base, i.e., relationships among the higher taxonomic clades such as phyla, class, etc. A recent study indicates that the incidence of LGTs shows linear correlation with the genome sequence and the GC content similarities of the donor and recipient organisms (Kloesges et al., 2011). Hence, while many of the observed inconsistencies between different gene trees at the species/strain levels could be due to LGTs (Puigbo et al., 2009; Kloesges et al., 2011), the differences in branching pattern at the higher taxonomic levels are perhaps in large parts due to loss of the phylogenetic signal and the lack of resolving power of the tree-based phylogenetic approaches (Gupta, 1998; Ludwig and Klenk, 2005; Puigbo et al., 2009).

In this review we have discussed the usefulness of CSIs and CSPs, as novel and important class of molecular markers for understanding the evolutionary relationships among prokaryotes. We have presented compelling evidence that based upon the species distribution patterns of these molecular signatures different prokaryotic taxa from phylum down to the genus levels can be clearly identified. Additionally, based upon these markers it is also possible to reliably deduce the evolutionary relationships amongst different prokaryotic taxa, both within a phylum and among different phyla. The evolutionary relationships deduced based upon these molecular markers generally exhibit high degree of congruency with those indicated by 16S rRNA trees or other gene/protein sequences. The analyses based upon these markers have also been able to clarify some relationships that are not resolved in phylogenetic trees. The species distribution patterns of these markers thus provide strong evidence that different clades of bacteria have evolved in a tree-like manner and that the prokaryotic organisms are not an exception to the Darwinian model of evolution. The relatively small numbers of these CSIs where the indel is also present in some unrelated species, which could be due to LGTs, show no specific pattern or relationship, thus they have minimal or no impact on the strong and consistent tree-like branching pattern that is evident from all other identified CSIs. However, it should be acknowledged that all of the work using CSIs and CSPs on understanding the evolutionary relationships among prokaryotes has thus far been carried out at genus level or higher taxa. Hence, it remains to be seen whether this approach will prove equally useful in clarifying the evolutionary relationships at the

species or strain levels or not, where the evolutionary flux and the incidences of LGTs are deemed to be the highest (Daubin et al., 2003; Lerat et al., 2003; Dagan et al., 2008; Puigbo et al., 2009; Kloesges et al., 2011).

The molecular markers such as those described here in addition to their usefulness for understanding prokaryotic phylogeny also provide valuable means to address/clarify a number of important aspects of microbiology. (1) Based upon these markers different prokaryotic taxa can now be identified in clear molecular terms rather than only as phylogenetic entities. (2) Based upon them the boundaries of different taxonomic clades can also be more clearly defined. (3) Due to their high degree of specificity and predictive ability, they provide important diagnostic tools for identifying both known and unknown species belonging to these groups of bacteria. (4) The shared presence of these CSIs by unrelated groups of bacteria provides potential means for identifying novel cases of LGTs. (5) Functional studies on these molecular markers should help in the discovery of novel biochemical or physiological properties that are distinctive characteristics of different groups of prokaryotes.

Lastly, it should be acknowledged that the number of genes which harbor rare genetic changes such as these CSIs is generally small in comparison to the total number of genes that are present in any genome. However, the genes containing these CSIs are involved in different essential functions and they are often among the most conserved proteins found in various organisms. Although, the criticism could be levied that the inferences based upon small numbers of genes/proteins containing these CSIs are not representative of the entire genomes (Dagan and Martin, 2006; Bapteste and Boucher, 2008), it should be emphasized that in a number of studies such as those discussed here, the reported CSIs or CSPs represent analyses of the entire genomes. Based upon these CSIs and/or CSPs, no other significant or consistent relationships or patterns among these organisms, other than those indicated here, can be derived from consideration of all of the gene/protein sequences in these genomes using these approaches. In this context it is also helpful to remember that molecular sequences like all other fossils change and disintegrate over long evolutionary periods of time and they lose their information content at different rates. Hence, a well-preserved fossil is generally considered to be far more informative than hundreds or even thousands of disintegrated fossils. Following this analogy, it is expected that not all genes/proteins will prove equally useful for understanding the evolutionary history of prokaryotes, which spans > 3.5 billion years. Thus, the best we can hope for is to find significant numbers of conserved genes/proteins, which contain consistent and reliable signals such as those described in the present work, whose inferences are generally consistent with all/most other available information.

ACKNOWLEDGMENTS

This work was supported by a research grant from the Natural Science and Engineering Research Council of Canada. Hafiz S. Naushad was partly supported by a scholarship from the Islamia University of Bahawalpur.

REFERENCES

- Andam, C. P., and Gogarten, J. P. (2011). Biased gene transfer in microbial evolution. *Nat. Rev. Microbiol.* 9, 543–555.
- Baldauf, S. L., and Palmer, J. D. (1993). Animals and fungi are each other's closest relatives: congruent evidence from multiple proteins. *Proc. Natl. Acad. Sci. U.S.A.* 90, 11558–11562.
- Bapteste, E., and Boucher, Y. (2008). Lateral gene transfer challenges principles of microbial systematics. *Trends Microbiol.* 16, 200–207.
- Bapteste, E., Brochier, C., and Boucher, Y. (2005a). Higher-level classification of the Archaea: evolution of methanogenesis and methanogens. *Archaea* 1, 353–363.
- Bapteste, E., Susko, E., Leigh, J., MacLeod, D., Charlebois, R. L., and Doolittle, W. F. (2005b). Do orthologous gene phylogenies really support tree-thinking? *BMC Evol. Biol.* 5, 33.
- Bapteste, E., O'Malley, M. A., Beiko, R. G., Ereshefsky, M., Gogarten, J. P., Franklin-Hall, L., Lapointe, F. J., Dupre, J., Dagan, T., Boucher, Y., and Martin, W. (2009). Prokaryotic evolution and the tree of life are two different things. *Biol. Dir.* 4, 34.
- Beiko, R. G. (2011). Telling the whole story in a 10,000-genome world. *Biol. Dir.* 6, 34.
- Beiko, R. G., Harlow, T. J., and Ragan, M. A. (2005). Highways of gene sharing in prokaryotes. *Proc. Natl. Acad. Sci. U.S.A.* 102, 14332–14337.
- Blair, C., and Murphy, R. W. (2011). Recent trends in molecular phylogenetic analysis: where to next? *J. Hered.* 102, 130–138.
- Boone, D. R., Castenholz, R. W., and Garrity, G. M. (2001). *Bergey's Manual of Systematic Bacteriology*. New York, NY: Springer, 1–721.
- Boto, L. (2010). Horizontal gene transfer in evolution: facts and challenges. *Proc. Biol. Sci.* 277, 819–827.
- Brochier, C., Forterre, P., and Gribaldo, S. (2004). Archaeal phylogeny based on proteins of the transcription and translation machineries: tackling the Methanopyrus kandleri paradox. *Genome Biol.* 5, R17.
- Brochier-Armanet, C., Bousseau, B., Gribaldo, S., and Forterre, P. (2008). Mesophilic Crenarchaeota: proposal for a third Archaeal phylum, the Thaumarchaeota. *Nat. Rev. Microbiol.* 6, 245–252.
- Buchanan, R. E., and Gibbons, N. E. (1974). *Bergey's Manual of Determinative Bacteriology*. Baltimore, MD: Williams and Wilkins.
- Burggraf, S., Stetter, K. O., Rouviere, P., and Woese, C. R. (1991). Methanopyrus kandleri: an Archaeal methanogen unrelated to all other known methanogens. *Syst. Appl. Microbiol.* 14, 346–351.
- Ciccarelli, F. D., Doerks, T., von Mering, C., Creevey, C. J., Snel, B., and Bork, P. (2006). Toward automatic reconstruction of a highly resolved tree of life. *Science* 311, 1283–1287.
- Connors, S. B., Mongodin, E. F., Johnson, M. R., Montero, C. I., Nelson, K. E., and Kelly, R. M. (2006). Microbial biochemistry, physiology, and biotechnology of hyperthermophilic Thermotoga species. *FEMS Microbiol. Rev.* 30, 872–905.
- Cowan, S. T. (1965). Principles and practice of bacterial taxonomy—a forward look. *J. Gen. Microbiol.* 39, 143–153.
- Cutino-Jimenez, A. M., Martins-Pinheiro, M., Lima, W. C., Martin-Tornet, A., Morales, O. G., and Menck, C. F. (2010). Evolutionary placement of Xanthomonadales based on conserved protein signature sequences. *Mol. Phylogenet. Evol.* 54, 524–534.
- Dagan, T. (2011). Phylogenomic networks. *Trends Microbiol.* 19, 483–491.
- Dagan, T., Artzy-Randrup, Y., and Martin, W. (2008). Modular networks and cumulative impact of lateral transfer in prokaryote genome evolution. *Proc. Natl. Acad. Sci. U.S.A.* 105, 10039–10044.
- Dagan, T., and Martin, W. (2006). The tree of one percent. *Genome Biol.* 7, 118.
- Dagan, T., and Martin, W. (2007). Ancestral genome sizes specify the minimum rate of lateral gene transfer during prokaryote evolution. *Proc. Natl. Acad. Sci. U.S.A.* 104, 870–875.
- Darwin, C. (1859). *The Origin of Species by Means of Natural Selection or the Preservation of Favoured Races in the Struggle for Life*. London: John Murray.
- Daubin, V., Gouy, M., and Perriere, G. (2002). A phylogenomic approach to bacterial phylogeny: evidence of a core of genes sharing a common history. *Genome Res.* 12, 1080–1090.
- Daubin, V., and Ochman, H. (2004). Bacterial genomes as new gene homes: the genealogy of ORFans in *E. coli*. *Genome Res.* 14, 1036–1042.
- Daubin, V., Moran, N. A., and Ochman, H. (2003). Phylogenetics and the cohesion of bacterial genomes. *Science* 301, 829–832.
- Delsuc, F., Brinkmann, H., and Philippe, H. (2005). Phylogenomics and the reconstruction of the tree of life. *Nat. Rev. Genet.* 6, 361–375.
- Doolittle, W. F. (1999). Phylogenetic classification and the universal tree. *Science* 284, 2124–2129.
- Doolittle, W. F. (2000). Uprooting the tree of life. *Sci. Am.* 282, 90–95.
- Doolittle, W. F., and Bapteste, E. (2007). Pattern pluralism and the Tree of Life hypothesis. *Proc. Natl. Acad. Sci. U.S.A.* 104, 2043–2049.
- Dutilh, B. E., Huynen, M. A., Bruno, W. J., and Snel, B. (2004). The consistent phylogenetic signal in genome trees revealed by reducing the impact of noise. *J. Mol. Evol.* 58, 527–539.
- Dutilh, B. E., Snel, B., Ettema, T. J., and Huynen, M. A. (2008). Signature genes as a phylogenomic tool. *Mol. Biol. Evol.* 25, 1659–1667.
- Euzeby, J. P. (2011). List of prokaryotic names with standing in nomenclature. <http://www.bacterio.cict.fr/classifphylo.html>. (Ref Type: Generic).
- Fang, G., Rocha, E., and Danchin, A. (2005). How essential are nonessential genes? *Mol. Biol. Evol.* 22, 2147–2156.
- Fleischmann, R. D., Adams, M. D., White, O., Clayton, R. A., Kirkness, E. F., Kerlavage, A. R., Bult, C. J., Tomb, J. F., Dougherty, B. A., and Merrick, J. M. (1995). Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science* 269, 496–512.
- Galperin, M. Y., and Koonin, E. V. (2004). 'Conserved hypothetical' proteins: prioritization of targets for experimental study. *Nucleic Acids Res.* 32, 5452–5463.
- Galtier, N. (2007). A model of horizontal gene transfer and the bacterial phylogeny problem. *Syst. Biol.* 56, 633–642.
- Gao, B., and Gupta, R. S. (2005). Conserved indels in protein sequences that are characteristic of the phylum Actinobacteria. *Int. J. Syst. Evol. Microbiol.* 55, 2401–2412.
- Gao, B., and Gupta, R. S. (2007). Phylogenomic analysis of proteins that are distinctive of Archaea and its main subgroups and the origin of methanogenesis. *BMC Genomics* 8, 86.
- Gao, B., and Gupta, R. S. (2012a). Microbial systematics in the post-genomics era. *Antonie Van Leeuwenhoek* 101, 45–54.
- Gao, B., and Gupta, R. S. (2012b). Phylogenetic framework and molecular signatures for the main clades of the phylum actinobacteria. *Microbiol. Mol. Biol. Rev.* 76, 66–112.
- Gao, B., Mohan, R., and Gupta, R. S. (2009). Phylogenomics and protein signatures elucidating the evolutionary relationships among the Gammaproteobacteria. *Int. J. Syst. Evol. Microbiol.* 59, 234–247.
- Gao, B., Paramanathan, R., and Gupta, R. S. (2006). Signature proteins that are distinctive characteristics of Actinobacteria and their subgroups. *Antonie Van Leeuwenhoek* 90, 69–91.
- Garcia-Vallve, S., Romeu, A., and Palau, J. (2000). Horizontal gene transfer in bacterial and archaeal complete genomes. *Genome Res.* 10, 1719–1725.
- Gogarten, J. P., Doolittle, W. F., and Lawrence, J. G. (2002). Prokaryotic evolution in light of gene transfer. *Mol. Biol. Evol.* 19, 2226–2238.
- Grabarse, W., Mahlert, E., Duin, E. C., Goubeaud, M., Shima, S., Thauer, R. K., Lamzin, V., and Ermler, U. (2001). On the mechanism of biological methane formation: structural evidence for conformational changes in methyl-coenzyme M reductase upon substrate binding. *J. Mol. Biol.* 309, 315–330.
- Gribaldo, S., and Brochier-Armanet, C. (2006). The origin and evolution of Archaea: a state of the art. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 361, 1007–1022.
- Griffiths, E., and Gupta, R. S. (2001). The use of signature sequences in different proteins to determine the relative branching order of bacterial divisions: evidence that Fibrobacter diverged at a similar time to Chlamydia and the Cytophaga-Flavobacterium-Bacteroides division. *Microbiology* 147, 2611–2622.
- Griffiths, E., and Gupta, R. S. (2004a). Distinctive protein signatures provide molecular markers and evidence for the monophyletic nature of the deinococcus-thermus phylum. *J. Bacteriol.* 186, 3097–3107.
- Griffiths, E., and Gupta, R. S. (2004b). Signature sequences in diverse proteins provide evidence for the late divergence of the Order Aquificales. *Int. Microbiol.* 7, 41–52.
- Griffiths, E., and Gupta, R. S. (2006a). Lateral transfers of serine hydroxymethyltransferase (glyA) and UDP-N-acetylglucosamine enolpyruvyl transferase (murA) genes from free-living Actinobacteria to the parasitic chlamydiae. *J. Mol. Evol.* 63, 283–296.
- Griffiths, E., and Gupta, R. S. (2006b). Molecular signatures in protein sequences that are characteristic of the phylum Aquificae. *Int. J. Syst. Evol. Microbiol.* 56, 99–107.
- Griffiths, E., and Gupta, R. S. (2007a). Identification of signature proteins that are distinctive of the

- Deinococcus-Thermus phylum. *Int. Microbiol.* 10, 201–208.
- Griffiths, E., and Gupta, R. S. (2007b). Phylogeny and shared conserved inserts in proteins provide evidence that Verrucomicrobia are the closest known free-living relatives of chlamydiae. *Microbiology* 153, 2648–2654.
- Griffiths, E., Petrich, A. K., and Gupta, R. S. (2005). Conserved indels in essential proteins that are distinctive characteristics of Chlamydiales and provide novel means for their identification. *Microbiology* 151, 2647–2657.
- Gupta, R. S. (1998). Protein phylogenies and signature sequences: a reappraisal of evolutionary relationships among archaeobacteria, eubacteria, and eukaryotes. *Microbiol. Mol. Biol. Rev.* 62, 1435–1491.
- Gupta, R. S. (2000a). The natural evolutionary relationships among prokaryotes. *Crit. Rev. Microbiol.* 26, 111–131.
- Gupta, R. S. (2000b). The phylogeny of proteobacteria: relationships to other eubacterial phyla and eukaryotes. *FEMS Microbiol. Rev.* 24, 367–402.
- Gupta, R. S. (2001). The branching order and phylogenetic placement of species from completed bacterial genomes, based on conserved indels found in various proteins. *Int. Microbiol.* 4, 187–202.
- Gupta, R. S. (2003). Evolutionary relationships among photosynthetic bacteria. *Photosynth. Res.* 76, 173–183.
- Gupta, R. S. (2004). The phylogeny and signature sequences characteristics of Fibrobacteres, Chlorobi, and Bacteroidetes. *Crit. Rev. Microbiol.* 30, 123–143.
- Gupta, R. S. (2006). Molecular signatures (unique proteins and conserved indels) that are specific for the epsilon proteobacteria (Campylobacteriales). *BMC Genomics* 7, 167.
- Gupta, R. S. (2009). Protein signatures (molecular synapomorphies) that are distinctive characteristics of the major cyanobacterial clades. *Int. J. Syst. Evol. Microbiol.* 59, 2510–2526.
- Gupta, R. S. (2010a). “Applications of conserved indels for understanding microbial phylogeny,” in *Molecular Phylogeny of Microorganisms*, eds A. Oren and R. T. Papke (Norfolk, UK: Caister Academic Press), 135–150.
- Gupta, R. S. (2010b). Molecular signatures for the main phyla of photosynthetic bacteria and their subgroups. *Photosynth. Res.* 104, 357–372.
- Gupta, R. S. (2011). Origin of diderm (Gram-negative) bacteria: antibiotic selection pressure rather than endosymbiosis likely led to the evolution of bacterial cells with two membranes. *Antonie Van Leeuwenhoek* 100, 171–182.
- Gupta, R. S. (2012). Origin and spread of photosynthesis based upon conserved sequence Features in key bacteriochlorophyll biosynthesis proteins. *Mol. Biol. Evol.* PMID: 22628531. [Epub ahead of print].
- Gupta, R. S., and Bhandari, V. (2011). Phylogeny and molecular signatures for the phylum Thermotogae and its subgroups. *Antonie Van Leeuwenhoek* 100, 1–34.
- Gupta, R. S., and Gao, B. (2009). Phylogenomic analyses of clostridia and identification of novel protein signatures that are specific to the genus *Clostridium* sensu stricto (cluster I). *Int. J. Syst. Evol. Microbiol.* 59, 285–294.
- Gupta, R. S., and Griffiths, E. (2002). Critical issues in bacterial phylogeny. *Theor. Popul. Biol.* 61, 423–434.
- Gupta, R. S., and Griffiths, E. (2006). Chlamydiae-specific proteins and indels: novel tools for studies. *Trends Microbiol.* 14, 527–535.
- Gupta, R. S., and Mathews, D. W. (2010). Signature proteins for the major clades of Cyanobacteria. *BMC Evol. Biol.* 10, 24.
- Gupta, R. S., and Mok, A. (2007). Phylogenomics and signature proteins for the alpha proteobacteria and its main groups. *BMC Microbiol.* 7, 106.
- Gupta, R. S., and Shami, A. (2011). Molecular signatures for the Crenarchaeota and the Thaumarchaeota. *Antonie Van Leeuwenhoek* 99, 133–157.
- Haggerty, L. S., Martin, F. J., Fitzpatrick, D. A., and McInerney, J. O. (2009). Gene and genome trees conflict at many levels. *Philos. Trans. R. Soc. Lond. B Biol Sci.* 364, 2209–2219.
- Harms, U., Weiss, D. S., Gartner, P., Linder, D., and Thauer, R. K. (1995). The energy conserving N5-methyltetrahydromethanopterin: coenzyme M methyltransferase complex from *Methanobacterium thermoautotrophicum* is composed of eight different subunits. *Eur. J. Biochem.* 228, 640–648.
- Herndl, G. J., Reinthaler, T., Teira, E., van Aken, H., Veth, C., Pernthaler, A., and Pernthaler, J. (2005). Contribution of Archaea to total prokaryotic production in the deep Atlantic Ocean. *Appl. Environ. Microbiol.* 71, 2303–2309.
- Huber, R., and Hannig, M. (2006). “Thermotogales,” in *The Prokaryotes*, eds M. Dworkin, S. Falkow, E. Rosenberg, K. H. Schleifer, and E. Stackebrandt (New York, NY: Springer), 899–922.
- Huber, R., Langworthy, T. A., Konig, H., Thomm, M., Woese, C. R., Sleytr, U. B., and Stetter, K. O. (1986). *Thermotoga maritima* sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90 °C*. *Arch. Microbiol.* 144, 324–333.
- Iguchi, A., Thomson, N. R., Ogura, Y., Saunders, D., Ooka, T., Henderson, I. R., Harris, D., Asadulghani, M., Kurokawa, K., Dean, P., Kenny, B., Quail, M. A., Thurston, S., Dougan, G., Hayashi, T., Parkhill, J., and Frankel, G. (2009). Complete genome sequence and comparative genome analysis of enteropathogenic *Escherichia coli* O127, H6 strain E2348/69. *J. Bacteriol.* 191, 347–354.
- Jain, R., Rivera, M. C., and Lake, J. A. (1999). Horizontal gene transfer among genomes: the complexity hypothesis. *Proc. Natl. Acad. Sci. U.S.A.* 96, 3801–3806.
- Jordan, G., and Goldman, N. (2012). The effects of alignment error and alignment filtering on the sitewise detection of positive selection. *Mol. Biol. Evol.* 29, 1125–1139.
- Kloesges, T., Popa, O., Martin, W., and Dagan, T. (2011). Networks of gene sharing among 329 proteobacterial genomes reveal differences in lateral gene transfer frequency at different phylogenetic depths. *Mol. Biol. Evol.* 28, 1057–1074.
- Koonin, E. V. (2007). The Biological Big Bang model for the major transitions in evolution. *Biol. Dir.* 2, 21.
- Koski, L. B., and Golding, G. B. (2001). The closest BLAST hit is often not the nearest neighbor. *J. Mol. Evol.* 52, 540–542.
- Koski, L. B., Morton, R. A., and Golding, G. B. (2001). Codon bias and base composition are poor indicators of horizontally transferred genes. *Mol. Biol. Evol.* 18, 404–412.
- Kunin, V., Goldovsky, L., Darzentas, N., and Ouzounis, C. A. (2005). The net of life: reconstructing the microbial phylogenetic network. *Genome Res.* 15, 954–959.
- Kurland, C. G. (2005). What tangled web: barriers to rampant horizontal gene transfer. *Bioessays* 27, 741–747.
- Kurland, C. G., Canback, B., and Berg, O. G. (2003). Horizontal gene transfer: a critical view. *Proc. Natl. Acad. Sci. U.S.A.* 100, 9658–9662.
- Leininger, S., Ulrich, T., Schloter, M., Schwark, L., Qi, J., Nicol, G. W., Prosser, J. I., Schuster, S. C., and Schleper, C. (2006). Archaea predominate among ammonia-oxidizing prokaryotes in soils. *Nature* 442, 806–809.
- Lerat, E., Daubin, V., and Moran, N. A. (2003). From gene trees to organismal phylogeny in prokaryotes: the case of the gamma-Proteobacteria. *PLoS Biol.* 1:E19. doi: 10.1371/journal.pbio.0000019
- Lerat, E., Daubin, V., Ochman, H., and Moran, N. A. (2005). Evolutionary origins of genomic repertoires in bacteria. *PLoS Biol.* 3:e130. doi: 10.1371/journal.pbio.0030130
- Ludwig, W., and Klenk, H.-P. (2005). “Overview: a phylogenetic backbone and taxonomic framework for prokaryotic systematics,” in *Bergey’s Manual of Systematic Bacteriology*, eds D. J. Brenner, N. R. Krieg, J. T. Staley, and G. M. Garrity (Berlin: Springer-Verlag), 49–65.
- Marri, P. R., and Golding, G. B. (2008). Gene amelioration demonstrated: the journey of nascent genes in bacteria. *Genome* 51, 164–168.
- Martin, W. (1999). Mosaic bacterial chromosomes: a challenge en route to a tree of genomes. *Bioessays* 21, 99–104.
- Naushad, H. S., and Gupta, R. S. (2012). Molecular signatures (conserved indels) in protein sequences that are specific for the order Pasteurellales and distinguish two of its main clades. *Antonie Van Leeuwenhoek* 101, 105–124.
- NCBI genomic database. (2012). <http://www.ncbi.nlm.nih.gov/genomes/lproks.cgi>. (Ref Type: Electronic Citation).
- NCBI Taxonomy. (2012). <http://www.ncbi.nlm.nih.gov/taxonomy>. (Ref Type: Electronic Citation).
- Nelson, K. E., Clayton, R. A., Gill, S. R., Gwinn, M. L., Dodson, R. J., Haft, D. H., Hickey, E. K., Peterson, J. D., Nelson, W. C., Ketchum, K. A., McDonald, L., Utterback, T. R., Malek, J. A., Linher, K. D., Garrett, M. M., Stewart, A. M., Cotton, M. D., Pratt, M. S., Phillips, C. A., Richardson, D., Heidelberg, J., Sutton, G. G., Fleischmann, R. D., Eisen, J. A., White, O., Salzberg, S. L., Smith, H. O., Venter, J. C., and Fraser, C. M. (1999). Evidence for lateral gene transfer between Archaea and bacteria from genome sequence of *Thermotoga maritima*. *Nature* 399, 323–329.
- Ochman, H., Lawrence, J. G., and Groisman, E. A. (2000). Lateral gene transfer and the nature of bacterial innovation. *Nature* 405, 299–304.

- Oxley, A. P., Lanfranco, M. P., Wurdemann, D., Ott, S., Schreiber, S., McGenity, T. J., Timmis, K. N., and Nogales, B. (2010). Halophilic Archaea in the human intestinal mucosa. *Environ. Microbiol.* 12, 2398–2410.
- Pace, N. R. (1997). A molecular view of microbial diversity and the biosphere. *Science* 276, 734–740.
- Pennisi, E. (1999). Is it time to uproot the tree of life? *Science* 284, 1305–1307.
- Popa, O., Hazkani-Covo, E., Landan, G., Martin, W., and Dagan, T. (2011). Directed networks reveal genomic barriers and DNA repair bypasses to lateral gene transfer among prokaryotes. *Genome Res.* 21, 599–609.
- Puigbo, P., Wolf, Y. I., and Koonin, E. V. (2009). Search for a ‘Tree of Life’ in the thicket of the phylogenetic forest. *J. Biol.* 8, 59.
- Puigbo, P., Wolf, Y. I., and Koonin, E. V. (2010). The tree and net components of prokaryote evolution. *Genome Biol. Evol.* 2, 745–756.
- Ragan, M. A. (2001). On surrogate methods for detecting lateral gene transfer. *FEMS Microbiol. Lett.* 201, 187–191.
- Ragan, M. A., and Beiko, R. G. (2009). Lateral genetic transfer: open issues. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 364, 2241–2251.
- Raymond, J., Zhaxybayeva, O., Gogarten, J. P., Gerdes, S. Y., and Blankenship, R. E. (2002). Whole-genome analysis of photosynthetic prokaryotes. *Science* 298, 1616–1620.
- Reysenbach, A.-L. (2001). “Phylum BII. Thermotogae phy. nov.,” in *Bergey’s Manual of Systematic Bacteriology* eds G. M. Garrity, D. R. Boone, and R. W. Castenholz (Berlin: Springer), 369–387.
- Rivera, M. C., and Lake, J. A. (1992). Evidence that eukaryotes and eocyte prokaryotes are immediate relatives. *Science* 257, 74–76.
- Roettger, M., Martin, W., and Dagan, T. (2009). A machine-learning approach reveals that alignment properties alone can accurately predict inference of lateral gene transfer from discordant phylogenies. *Mol. Biol. Evol.* 26, 1931–1939.
- Rokas, A., and Holland, P. W. (2000). Rare genomic changes as a tool for phylogenetics. *Trends Ecol. Evol.* 15, 454–459.
- Schoeffler, A. J., May, A. P., and Berger, J. M. (2010). A domain insertion in *Escherichia coli* GyrB adopts a novel fold that plays a critical role in gyrase function. *Nucleic Acids Res.* 38, 7830–7844.
- Singh, B., and Gupta, R. S. (2009). Conserved inserts in the Hsp60 (GroEL) and Hsp70 (DnaK) proteins are essential for cellular growth. *Mol. Genet. Genomics* 281, 361–373.
- Sorek, R., Zhu, Y., Creevey, C. J., Francino, M. P., Bork, P., and Rubin, E. M. (2007). Genome-wide experimental determination of barriers to horizontal gene transfer. *Science* 318, 1449–1452.
- Stanier, R. Y., Adelberg, E. A., and Ingraham, J. L. (1976). *The Microbial World*. Englewood Cliffs, NJ: Prentice-Hall Inc., 1–871.
- Susko, E., Leigh, J., Doolittle, W. F., and Baptiste, E. (2006). Visualizing and assessing phylogenetic congruence of core gene sets: a case study of the gamma-proteobacteria. *Mol. Biol. Evol.* 23, 1019–1030.
- Swithers, K. S., Gogarten, J. P., and Fournier, G. P. (2009). Trees in the web of life. *J. Biol.* 8, 54.
- Tersteegen, A., and Hedderich, R. (1999). Methanobacterium thermoautotrophicum encodes two multisubunit membrane-bound [NiFe] hydrogenases. Transcription of the operons and sequence analysis of the deduced proteins. *Eur. J. Biochem.* 264, 930–943.
- Treangen, T. J., and Rocha, E. P. (2011). Horizontal transfer, not duplication, drives the expansion of protein families in prokaryotes. *PLoS Genet.* 7:e1001284. doi: 10.1371/journal.pgen.1001284
- Wang, B. (2001). Limitations of compositional approach to identifying horizontally transferred genes. *J. Mol. Evol.* 53, 244–250.
- Williams, D., Fournier, G. P., Lapierre, P., Swithers, K. S., Green, A. G., Andam, C. P., and Gogarten, J. P. (2011). A rooted net of life. *Biol. Dir.* 6, 45.
- Woese, C. R. (1987). Bacterial evolution. *Microbiol. Rev.* 51, 221–271.
- Woese, C. R., and Fox, G. E. (1977). Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proc. Natl. Acad. Sci. U.S.A.* 74, 5088–5090.
- Woese, C. R., Kandler, O., and Wheelis, M. L. (1990). Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc. Natl. Acad. Sci. U.S.A.* 87, 4576–4579.
- Wolf, Y. I., Rogozin, I. B., Grishin, N. V., and Koonin, E. V. (2002). Genome trees and the Tree of Life. *Trends Genet.* 18, 472–479.
- Wu, D., Hugenholtz, P., Mavromatis, K., Pukall, R., Dalin, E., Ivanova, N. N., Kunin, V., Goodwin, L., Wu, M., Tindall, B. J., Hooper, S. D., Pati, A., Lykidis, A., Spring, S., Anderson, I. J., D’Haeseleer, P., Zemla, A., Singer, M., Lapidus, A., Nolan, M., Copeland, A., Han, C., Chen, F., Cheng, J. F., Lucas, S., Kerfeld, C., Lang, E., Gronow, S., Chain, P., Bruce, D., Rubin, E. M., Kyrpides, N. C., Klenk, H. P., and Eisen, J. A. (2009). A phylogeny-driven genomic encyclopaedia of Bacteria and Archaea. *Nature* 462, 1056–1060.
- Yap, W. H., Zhang, Z., and Wang, Y. (1999). Distinct types of rRNA operons exist in the genome of the actinomycete *Thermomonospora chromogena* and evidence for horizontal transfer of an entire rRNA operon. *J. Bacteriol.* 181, 5201–5209.
- Yarza, P., Ludwig, W., Euzéby, J., Amann, R., Schleifer, K. H., Glockner, F. O., and Rossello-Mora, R. (2010). Update of the All-Species Living Tree Project based on 16S and 23S rRNA sequence analyses. *Syst. Appl. Microbiol.* 33, 291–299.
- Zhaxybayeva, O., Nesbo, C. L., and Doolittle, W. F. (2007). Systematic overestimation of gene gain through false diagnosis of gene absence. *Genome Biol.* 8, 402.
- Zhaxybayeva, O., Swithers, K. S., Lapierre, P., Fournier, G. P., Bickhart, D. M., DeBoy, R. T., Nelson, K. E., Nesbo, C. L., Doolittle, W. F., Gogarten, J. P., and Noll, K. M. (2009). On the chimeric nature, thermophilic origin, and phylogenetic placement of the Thermotogales. *Proc. Natl. Acad. Sci. U.S.A.* 106, 5865–5870.
- Zuckerklund, E., and Pauling, L. (1965). Molecules as documents of evolutionary history. *J. Theor. Biol.* 8, 357–366.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 02 April 2012; accepted: 27 June 2012; published online: 26 July 2012.

Citation: Bhandari V, Naushad HS and Gupta RS (2012) Protein based molecular markers provide reliable means to understand prokaryotic phylogeny and support Darwinian mode of evolution. *Front. Cell. Inf. Microbiol.* 2:98. doi: 10.3389/fcimb.2012.00098

Copyright © 2012 Bhandari, Naushad and Gupta. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in other forums, provided the original authors and source are credited and subject to any copyright notices concerning any third-party graphics etc.