



Nitrate reductase ^{15}N discrimination in *Arabidopsis thaliana*, *Zea mays*, *Aspergillus niger*, *Picea angusta*, and *Escherichia coli*

Eli Carlisle^{1†}, Chris Yarnes¹, Michael D. Toney² and Arnold J. Bloom^{1*}

¹ Department of Plant Sciences, University of California, Davis, CA, USA

² Department of Chemistry, University of California, Davis, CA, USA

Edited by:

Cristina Cruz, Faculty of Sciences of the University of Lisbon, Portugal

Reviewed by:

Adrian Hegeman, University of Minnesota-Twin Cities, USA
Uener Kolukisaoglu, University of Tuebingen, Germany

*Correspondence:

Arnold J. Bloom, Department of Plant Sciences, University of California, One Shields Avenue, Davis, CA 95776, USA
e-mail: ajbloom@ucdavis.edu

† Present address:

Eli Carlisle, Department of Land, Air, and Water Resources, University of California, Davis, USA

Stable ^{15}N isotopes have been used to examine movement of nitrogen (N) through various pools of the global N cycle. A central reaction in the cycle involves the reduction of nitrate (NO_3^-) to nitrite (NO_2^-) catalyzed by nitrate reductase (NR). Discrimination against ^{15}N by NR is a major determinant of isotopic differences among N pools. Here, we measured *in vitro* ^{15}N discrimination by several NRs purified from plants, fungi, and a bacterium to determine the intrinsic ^{15}N discrimination by the enzyme and to evaluate the validity of measurements made using ^{15}N -enriched NO_3^- . Observed NR isotope discrimination ranged from 22 to 32‰ (kinetic isotope effects of 1.022–1.032) among the different isozymes at natural abundance ^{15}N (0.37%). As the fractional ^{15}N content of substrate NO_3^- increased from natural abundance, the product ^{15}N fraction deviated significantly from that expected based on substrate enrichment and ^{15}N discrimination measured at natural abundance. Additionally, isotopic discrimination by denitrifying bacteria used to reduce NO_3^- and NO_2^- in some protocols became a greater source of error as ^{15}N enrichment increased. We briefly discuss potential causes of the experimental artifacts with enriched ^{15}N and recommend against the use of highly enriched ^{15}N tracers to study N discrimination in plants or soils.

Keywords: nitrate reductase, ^{15}N discrimination, natural abundance, enriched ^{15}N , plant, fungi, bacteria, kinetic isotope effect

INTRODUCTION

Nitrogen (N) is a key limiting resource in many ecosystems, and so N biogeochemical cycles—particularly the processes of organismal N uptake and assimilation—are of great importance (Epstein and Bloom, 2005). The main N source for most plants is soil inorganic N in the forms of nitrate (NO_3^-) and ammonium (NH_4^+) ions dissolved in the soil solution. A primary pathway through which inorganic N is converted into organic N is NO_3^- assimilation via the biochemical pathway involving the sequential reactions catalyzed by the enzymes nitrate reductase (NR), nitrite reductase (NiR), glutamine synthetase (GS), and glutamate synthase (GOGAT). Reduction of NO_3^- to NO_2^- via NR is the rate limiting step in this pathway (Robinson et al., 1998; Evans, 2001), and thus the behavior of NR is central to biogeochemical plant-soil interactions as well as to internal plant N cycling (Tcherkez and Farquhar, 2006).

Eukaryotic NRs are members of the sulfite oxidase family of molybdoenzymes. They are found in three forms that use different reductants: an NADH-specific form typically found in higher plants and algae, an NAD(P)H-bispecific form found in higher plants, algae, and fungi, and an NADPH-specific form found only in fungi. All three forms are similar in overall composition (Campbell, 1999; Fischer et al., 2005) and require multimerization for full function, typically as either homodimers or

homotetramers (Campbell, 1999; Barbier and Campbell, 2005). Each subunit contains one each of three cofactors: a flavin adenine dinucleotide (FAD), a heme-Fe, and a Mo-molybdopterin (Mo-MPT) cofactor (Campbell, 1999).

Prokaryotic NRs are unrelated to the eukaryotic NRs. They are members of the DMSO (dimethylsulfoxide) reductase family of molybdoenzymes and have an assimilatory (Nas) and two dissimilatory forms. The dissimilatory forms include a membrane-bound respiratory form (Nar) and a periplasmic form (Nap) (Moreno-Vivián et al., 1999). All prokaryotic forms, like all eukaryotic forms, have Mo as a key component of the cofactor that reduces NO_3^- . In prokaryotic NRs, however, the Mo is bound to a bis-molybdopterin guanine dinucleotide (MGD) rather than the MPT found in eukaryotes (Berks et al., 1995). The MGD molecule differs from the MPT molecule by the linkage of a guanine monophosphate (GMP) to the phosphate group of MPT (Temple and Rajagopalan, 2000). All prokaryotic NRs are heterotrimeric: they are composed of a Mo-MGD cofactor, a iron-sulfur (FeS) center, and either (a) FAD, (b) cytochrome b prosthetic, or (c) cytochrome c group in the Nas, Nap, or Nar forms, respectively (Moreno-Vivián et al., 1999). The prokaryotic NRs use a variety of electron donors, including ferredoxin, flavodoxin, and NADH for the Nas enzyme, quinones for the Nap enzymes, and quinols for the Nar enzymes.

Stable N isotopes (¹⁵N and ¹⁴N) are a standard method for examining N cycling and transformations on many scales, ranging from global biogeochemical N cycles (Shearer and Kohl, 1986; Schlesinger, 1997; Amundson et al., 2003) to mechanistic enzyme studies (Cleland, 2005). Two approaches are common, natural abundance and ¹⁵N-enriched. In the first approach, one monitors how the natural abundance ¹⁵N/¹⁴N ratio of various compounds in a pathway differs from a reference, typically that of atmospheric N₂ gas (0.37% ¹⁵N). An advantage of the natural abundance approach is that no additional N need be added that might perturb the experimental system. For example, inorganic N (e.g., NO₃⁻), if added above background levels, can inhibit N₂-fixation or alter N transformations and assimilation (Sprenst, 1979).

In the second approach, one employs a ¹⁵N-enriched reactant as a tracer and monitors δ¹⁵N differences through various N pools (Hauck and Bremner, 1976; Androssoff et al., 1995). Slight differences in substrate ¹⁵N content and variability introduced through mixing and fractionation of different N pools in the soil and the plant are overwhelmed by the tracer signal. This approach is not typically used to examine enzymatic isotope fractionation. Regardless of the experimental approach, a key assumption is that ¹⁵N behaves identically to ¹⁴N in biological systems except for the differences due to mass (Hauck and Bremner, 1976).

When environmental or substrate concentrations of NO₃⁻ are limiting, little to no discrimination against the ¹⁵N isotope is observed during NO₃⁻ assimilation (Kolb and Evans, 2003). Discrimination against the ¹⁵N isotope has been demonstrated to occur only when the NO₃⁻ concentration exceeds plant needs or uptake ability. The key to understanding these observations is that isotopic discrimination is only detected to the extent that both isotopic species are equilibrated between the plant and the environment during the course of the measurement (Kolb and Evans, 2003). In such cases, plant NR has typically been cited as discriminating against ¹⁵N by 15‰ when the enzyme is supplied adequate amounts of NO₃⁻ substrate containing N isotopes at natural abundance (i.e., product NO₂⁻ is depleted in ¹⁵N relative to substrate NO₃⁻ by 15 parts per 1000; Mariotti et al., 1982; Ledgard et al., 1985; Evans et al., 1996; Tcherkez and Farquhar, 2006). Nonetheless, reported values for NR ¹⁵N discrimination vary from 5 to 30‰ (Schmidt and Medina, 1991; Granger et al., 2004; Needoba et al., 2004; Tcherkez and Farquhar, 2006).

Previously, we obtained significantly different results for the rate of NO₃⁻ assimilation in plant shoots when using ¹⁴N or ¹⁵N as a tracer (Bloom et al., 2010). *Arabidopsis*, grown under ambient CO₂ concentrations with natural abundance NO₃⁻ nutrition and pulsed with 25% enriched ¹⁵NO₃⁻, demonstrated 20–30% less NO₃⁻ assimilation than *Arabidopsis* grown on 99% enriched ¹⁵NO₃⁻ and pulsed with natural abundance NO₃⁻. In wheat, these experimental protocols resulted in similar trends (Bloom et al., 2010). These differences in assimilation rates may have been confounded by artifacts of the isotope measurements by the isotope ratio mass spectrometer (IRMS).

In this study, we examined how NR discrimination varies among different organisms, and determined whether ¹⁵N enrichment of substrate NO₃⁻ affects the measurement of NR discrimination, which may have caused the unusual results in our previous work (Bloom et al., 2010). We conducted *in vitro*

experiments on five purified NRs from plants, fungi, and a bacterium to address two hypotheses: (1) natural abundance ¹⁵N discrimination is similar among NRs; (2) substrate NO₃⁻ ¹⁵N-enrichment affects instrumental analysis of samples and the estimation of ¹⁵N discrimination.

MATERIALS AND METHODS

We measured ¹⁵N-isotope discrimination by commercial preparations of five different nitrate reductases. Four of these enzymes are assimilatory nitrate reductases including that from *Arabidopsis thaliana* (AtNaR2 NADH:nitrate reductase E.C. 1.7.1.1), *Aspergillus niger* (NAD(P)H:nitrate reductase E.C. 1.7.1.2), *Pichia angusta* (YNaR1 NAD(P)H:nitrate reductase E.C. 1.7.1.2), and *Zea mays* (maize; NADH:nitrate reductase E.C. 1.7.1.1); these were assessed using similar protocols. The fifth is a prokaryotic NR (*Escherichia coli*, cytochrome:nitrate reductase E.C. 1.9.6.1), which is a dissimilatory reductase (Nar form); this enzyme required a different experimental protocol. The maize, *Aspergillus*, and *E. coli* enzymes are isolated from the native organisms, while the *Arabidopsis* and yeast enzymes are recombinant enzymes expressed in the yeast *Pichia pastoris*. The *Aspergillus* and *E. coli* enzymes were purchased from Sigma, and the rest were purchased from NECi (nitrate.com; also see Barbier et al., 2004). We were unable to detect any nitrate or nitrite in the enzyme preparations (Miranda et al., 2001).

ASSIMILATORY NITRATE REDUCTASES

One unit of enzyme activity is defined as the amount of NR that is capable of catalyzing the conversion of 1 μmol nitrate to nitrite per minute at optimal pH and temperature. Assimilatory nitrate reductase analyses were carried out by reconstituting the freeze-dried *Arabidopsis*, yeast, and maize enzymes in 25 mM potassium phosphate, 0.1 mM Na₂EDTA, pH 7.5, at 22–25°C, to a final concentration of 1 unit/mL. For the *Aspergillus* enzyme, a 100 mM potassium phosphate buffer, 0.1 mM Na₂EDTA, pH 7.5 was used. The NR K_m for NO₃⁻ in *Arabidopsis* and maize ranges from 15 to 40 μM (Su et al., 1997; Campbell, 1999), 199 μM for *Aspergillus* (Gilliam et al., 1993), and 470 μM for the dissimilatory NR in *E. coli* (Adams and Mortenson, 1982). There is no information for the specific yeast NR used in our study, but NRs from other genera of yeast have a K_m ranging from 110 (Hipkin et al., 1986) to 740 μM (Morozkina et al., 2005).

The enzyme reaction solution contained 1 unit (1 mL) of reconstituted NR, 165 mL of additional phosphate buffer (pH 7.5), and 70 μmol of NADH or NADPH for the plant and fungal enzymes, respectively. Additionally, the *Aspergillus* solution contained 4 μmol of FAD. After mixing, 10 mL of solution was added to each of 16 sterile 15 mL centrifuge tubes. One mL of KNO₃ solution (0.2 M KNO₃) was then added to each centrifuge tube, and the tubes were placed into a water bath at 30°C (25°C for *Aspergillus*) for 30 min. The use of 0.2 mmol of KNO₃ in the reaction tubes was calculated to provide a surplus of NO₃⁻ for the enzymatic reaction, while minimizing the possibility that residual NO₃⁻ might saturate the mass spectrometer detectors. Therefore, the enzyme reactions were not necessarily operating under NO₃⁻ saturating conditions. The KNO₃ was enriched with different levels of ¹⁵N from natural abundance to >99 atom % ¹⁵N to examine

how substrate ^{15}N enrichment influences the estimation of NR ^{15}N discrimination using an IRMS. At the end of the incubation, samples were placed into 20 mL scintillation vials. The reaction was halted by adding 0.25 mL of 1 M HCl and swirling the vials (decreasing pH to below 4). After 1 min, 40 μL of 10 M KOH were added to the solution, raising its pH to 10–12 to prevent oxidation of NO_2^- to NO_3^- . Concentrations of NO_2^- and NO_3^- were determined using the Griess reaction (Miranda et al., 2001).

DISSIMILATORY NITRATE REDUCTASE

The reaction catalyzed by dissimilatory *E. coli* NR requires reducing conditions. The enzyme was reconstituted in 2 mL of cold degassed deionized water and added to 55 mL of degassed buffer (0.125 M MOPS, pH 7.0) containing 30 mg benzyl viologen (BV; 1,1'-Dibenzyl-4,4'-bipyridinium dichloride, Sigma) as an electron donor. Ten milliliter of buffer/enzyme solution were added to 15 mL centrifuge tubes containing 0.2 mmoles of KNO_3 of various ^{15}N enrichments and 0.25 mL of a sodium dithionite/sodium bicarbonate solution (500 mg of each added to 50 mL degassed DI H_2O) to maintain reducing conditions. If the blue color of the reduced BV dye started to fade, additional dithionite solution was added to the vial. The vials were incubated in a water bath at 30°C for 90 min. To stop the reaction, vials were shaken vigorously to oxidize the BV. At the end of the incubation, 75 μL of 10 N NaOH was added to each tube to minimize conversion of NO_2^- to NO_3^- .

PREPARATION OF BACTERIAL CULTURES FOR HEADSPACE N_2O ANALYSIS

For all forms of the NR enzyme, we analyzed the ^{15}N isotope composition of the headspace N_2O produced through bacterial denitrification of NO_3^- and NO_2^- . Bacterial cultures and solutions were prepared as in Böhlke et al. (2007). The bacterial cultures used in this study were *Stenotrophomonas nitritireducens* (ATCC# BAA-12) and *Pseudomonas chlororaphis* (ATCC# 43928), both of which were obtained from the American Type Culture Collection (www.atcc.org). Both species lack the enzymatic ability to reduce N_2O to N_2 . *S. nitritireducens* is limited to reducing NO_2^- , while *P. chlororaphis* is capable of reducing both NO_3^- and NO_2^- to N_2O . This specificity allowed a sequential reduction of our experimental samples to estimate the ^{15}N concentrations of the NO_3^- and NO_2^- independently.

The denitrifying bacteria were streaked on tryptic soy agar (TSA) plates composed of TSA (40 g/L) and tryptic soy broth (TSB; 10 g/L) for *S. nitritireducens*, and TSA (40 g/L) and KNO_3 (1 g/L) for *P. chlororaphis*. The TSA plates for the *S. nitritireducens* cultures were prepared with 10 mM Na_2WO_4 to minimize any potential NO_3^- reduction (Nielsen et al., 2004). Both cultures were grown aerobically at 22–24°C. After 5–7 days, bacterial colonies were streaked onto a second plate. If re-plated more than 2 or 3 times, higher background levels of N_2O were observed for both bacteria. Therefore, all N_2O analyses were made using colonies from the second plate; additional analyses required starting the process over from the original frozen aliquot.

Individual *S. nitritireducens* colonies were transferred 4–5 days after streaking to sterile 50 mL centrifuge tubes with 35 mL of a growth medium containing TSB solution (15.0 g/L), $(\text{NH}_4)_2\text{SO}_4$ (0.25 g/L), and KH_2PO_4 (2.45 g/L). Tubes were capped leaving

an air headspace and agitated on a reciprocal shaker for 1–2 days at 22°C. The tubes were opened daily to allow for aerobic growth. The contents of the centrifuge tubes were combined and an antifoaming agent was added, 2–3 drops of Antifoam B (Sigma) per 105 mL solution. Four milliliter of the combined solution were pipetted into each 20 mL vial, which was capped with a septum and crimped, and then placed on a manifold that purged the vials with N_2 gas for 4 h to remove any residual oxygen, CO_2 , and N_2O . The contents of these vials were used for ^{15}N analysis of NO_2^- , as well as for providing substrate for the NO_3^- reducing *P. chlororaphis* cultures.

Pseudomonas chlororaphis cultures were used to inoculate a starter tube containing 1.5 mL of TSB solution (30 g/L), KNO_3 (1 g/L), and $(\text{NH}_4)_2\text{SO}_4$ (0.066 g/L), which was placed on a reciprocal shaker for 1 day. TSB solution (400 mL) was added to a 500 mL media bottle, which was inoculated with the contents of the starter tube and sealed tightly. The media bottle was placed on the reciprocal shaker, and the bacteria were allowed to grow for 6–10 days under anaerobic conditions. At the end of the 6–10 days, the solution was centrifuged (10 min at 20°C and 6500 rpm). The supernatant was decanted into a large beaker and the pellets remaining in the centrifuge tubes were resuspended in 60 mL of the supernatant solution. An antifoam agent—one drop of Antifoam B (Sigma) per 60 mL—was added and the concentrated bacterial solution was mixed thoroughly. Two milliliter of this solution were pipetted into each 20 mL headspace vial, capped with a septum and crimped, and then purged for 4 h on the manifold system with N_2 gas. The manifold system was carefully sterilized between sample sets.

BACTERIAL NO_2^- AND NO_3^- REDUCTION TO N_2O

Solutions from the enzyme reaction were added to the purged *S. nitritireducens* vials to determine the ^{15}N enrichment of N_2O derived from NO_2^- . The amount of solution added to the vial was determined by the concentration of the NO_2^- in the solution. A target value of 0.025 μmol of NO_2^- were added to each vial. This corresponded to volumes ranging from 0.25 to 1.5 mL depending on the enzyme. The vials were then placed on a reciprocal shaker for 24 h. At the end of 24 h, a sample was removed from the vial using a gas-tight syringe and injected into a vial containing *P. chlororaphis*. Again, a volume of solution equivalent to 0.025 μmol of NO_3^- was added to the vials (volumes ranged from 25–200 μL). Amounts of N added to all samples were matched to ensure that the samples remained in the linear range of the detectors in the IRMS. The *P. chlororaphis* vials were placed on the reciprocal shaker.

Once the subsample had been removed from the vials containing *S. nitritireducens*, and after 24 h incubation for the *P. chlororaphis* vials, 0.2 mL of 10 M NaOH were added to both sets of vials killing the bacteria and removing any residual headspace CO_2 . All vials were then taken to the UC Davis Stable Isotope Facility where they were analyzed for $[\text{N}_2\text{O}]$ and ^{15}N (ThermoFinnigan GasBench + PreCon trace gas concentration system interfaced to a ThermoScientific Delta V Plus IRMS, Bremen, Germany). To control for potential contamination, blanks (including bacterial solution alone and with DDI water), natural abundance NO_2^- and NO_3^- samples, and appropriately ^{15}N -enriched NO_2^- and

NO₃⁻ standards, were included in each sample run. The blanks also allowed us to correct for any N₂O derived from residual NO₃⁻ remaining in the *P. chlororaphis* solution and instrumental leaks. Samples of the enzyme reaction solution were also tested for inorganic N contamination and found not to contain detectable amounts. No remaining inorganic N was found in the spent nutrient solution after the bacterial incubations. Thus, conversion of the NO₃⁻ and NO₂⁻ to N₂O was complete and quantitative.

ANALYSIS OF ¹⁵N

We defined ⁴⁵R and ⁴⁶R as the ratios of measured ion peak areas [⁴⁵R = (m/z 45)/(m/z 44), and ⁴⁶R = (m/z 46)/(m/z 44)]. ¹⁵R or the ratio of ¹⁵N/¹⁴N of the N₂O headspace was determined as:

$$^{15}R = \frac{^{45}R - ^{17}R}{2} \quad (1)$$

where ¹⁷R (¹⁷O/¹⁶O) is fixed at the isotopic oxygen signature of ambient N₂O (¹⁷R = 0.00038855; Kaiser et al., 2003). ¹⁵R may equivalently be determined using ⁴⁶R as described in Stevens and Laughlin (1998):

$$^{15}R = -^{17}R + \sqrt{[(^{17}R)^2 + ^{46}R - ^{18}R]} \quad (2)$$

where ¹⁸R (¹⁸O/¹⁶O) is fixed at the isotopic oxygen signature of N₂O (δ¹⁸O = 44.62 and ¹⁸R = 0.0020947; Kaiser et al., 2003).

Isotope analyses are often expressed in terms of either δ¹⁵N or atom % ¹⁵N. δ¹⁵N is expressed as the per mil difference between the ¹⁵N/¹⁴N ratio of a sample relative to the primary reference material for nitrogen, atmospheric N₂ ("Air," 0‰):

$$\delta^{15}N(\text{‰}) = \left(\frac{\frac{^{15}N_{\text{sample}}}{^{14}N_{\text{sample}}}}{\frac{^{15}N_{\text{standard}}}{^{14}N_{\text{standard}}}} - 1 \right) * 1000 \quad (3)$$

For all analyses, initial measurements are made relative to a working reference gas of the same species (i.e., N₂O, N₂), co-measured during analysis. Subsequently, all measurements were corrected using working standards previously calibrated to the following standard reference materials IAEA-N1, IAEA-N2, IAEA-N3, USGS-40, and USGS-41.

There are, however, no internationally accepted reference standards for N₂O at high ¹⁵N enrichments and, because of this, we compared the ¹⁵N/¹⁴N ratio of the sample N₂O to the ¹⁵N/¹⁴N ratio of the substrate NO₃⁻ as determined by the quantitative combustion of substrate NO₃⁻ to N₂ (Stickrod and Marshall, 2000; See Supplemental Table 1) via a PDZ Europa ANCA-GSL Elemental Analyzer interfaced to a PDZ Europa 20-20 IRMS (Sercon Ltd., Cheshire, UK) according to:

$$\delta^{15}N(\text{‰}) = \left(\frac{\frac{^{15}N_{\text{sample}}}{^{14}N_{\text{sample}}}}{\frac{^{15}N_{\text{substrate}}}{^{14}N_{\text{substrate}}}} - 1 \right) * 1000 \quad (4)$$

This approach minimizes the impact of absolute errors in calibration because only differences between the substrate and product are considered, and the δ notation is very sensitive to large absolute differences between the sample and reference ratios (Fry, 2006). Also, in order to examine the performance of the bacterial reduction method at high levels of ¹⁵N enrichment, we calculated the atom % ¹⁵N from the atomic ratios, ¹⁵R:

$$\text{Atom \% } ^{15}N = \frac{(100 * ^{15}R)}{(1 + ^{15}R)} \quad (5)$$

Alternatively, the atom % ¹⁵N could be determined from the molecular ratios (e.g., ⁴⁵R and ⁴⁶R) as outlined in Stevens and Laughlin (1998).

The kinetic isotope effect (KIE) for a chemical reaction is the relative reactivity of molecules labeled with different isotopes (Farquhar et al., 1989). Data for competitive KIE experiments is generally taken at different values of fractional reaction of the initial substrate and fitted to Equation (6a), where *f* is fractional conversion of the initial substrate, *R*₀ is the isotopic ratio of the initial substrate (equal to the isotopic ratio of the product at 100% conversion), and *R*_{*f*} is the isotopic ratio of the product at fractional reaction *f*. Here, because all reactions were halted at very low conversions of the initial NO₃⁻ (0.5% maximal conversion), the KIE is well approximated by the molar ¹⁵N/¹⁴N ratio of the initial nitrate substrate to the nitrite product (Equation (6b); Bigeleisen and Wolfsberg, 1958):

$$KIE = \frac{\ln(1 - f)}{\ln(1 - f \times (R_f/R_0))} \quad (6a)$$

$$KIE = \frac{R_0}{R_f} \quad (6b)$$

Therefore, the KIE is related to the measured isotopic discrimination (Δ) by

$$\Delta = KIE - 1 \quad (7)$$

Alternatively, it can be derived by the following equation (Evans, 2001):

$$\Delta = 1000 \left[\frac{\delta^{15}N_0 - \delta^{15}N_f}{1 + \frac{\delta^{15}N_f}{1000}} \right] \quad (8)$$

where δ¹⁵N₀ and δ¹⁵N_{*f*} are the ¹⁵N content of the initial substrate and the product at fractional reaction *f*, respectively. Enzyme discrimination data will be presented using the Δ notation for the remainder of this paper.

The bacterial reduction method was compared with a chemical reduction method to examine potential bacterial discrimination against ¹⁵N during the reduction of NO₂⁻ to N₂O. For the chemical reduction, we used sulfamic acid to reduce NO₂⁻ to N₂O (Granger and Sigman, 2009). As a result of this reaction, the solution NO₂⁻ supplies one of the N atoms in the N₂O molecule,

while sulfamic acid supplies the other (Granger and Sigman, 2009). No remaining NO₃⁻ was observed in the samples after the reduction.

STATISTICAL ANALYSIS

The experiments were formulated as a randomized complete block design. Blocks were represented by the enzyme batch to control for inter-batch variation in enzyme activity and purity. Depending on the particular enzyme, a single batch of enzyme was used to produce from one to three repetitions of each ¹⁵NO₃ concentration to NO₂⁻. A linear mixed effects model (PROC MIXED, SAS 9.1, Cary, NC, USA) was used to analyze the data.

RESULTS

NR discrimination at natural abundance ranged (mean ± SE) from 22.5 ± 0.8‰ for *Arabidopsis* to 31.6 ± 2.3‰ for *E. coli* (Table 1). Interestingly, the two recombinant NRs (*Arabidopsis* and yeast) had the lowest discrimination among the enzymes. The average discrimination for the non-recombinant assimilatory eukaryotic NRs was 29.4 ± 0.4‰, significantly greater ($p < 0.0001$) than the average of 22.8 ± 0.5‰ for the recombinant NRs. The isotope discriminations of the maize and *E. coli* enzymes were significantly greater than that of the enzymes from *Arabidopsis* ($p = 0.0171$ and $p = 0.0052$, respectively) and yeast ($p = 0.0254$ and $p = 0.0052$, respectively). The KIEs had similar values among all species (Table 2) at natural abundance ¹⁵N.

There was no significant difference between the values obtained via the chemical and bacterial reduction methods (Figure 1; $p = 0.5633$), although the data from the chemical reduction method tended to be closer to the expected ¹⁵N enrichment at high ¹⁵N enrichments.

Observed KIEs (R_0/R_f) increased with increasing substrate ¹⁵N enrichment for all enzymes ($p < 0.001$; Figure 2, Table 2), particularly above 50 atom % ¹⁵N. The *Arabidopsis* enzyme showed the greatest deviation from expected, but all enzymes produced significantly less ¹⁵N-NO₂⁻ than expected at ¹⁵N enrichments above 50 atom % ($p < 0.001$).

DISCUSSION

NATURAL ABUNDANCE

The results show that ¹⁵N discrimination in the NR-catalyzed reaction is similar across eukaryotic NRs, particularly among non-recombinant NRs. All NRs had ¹⁵N discrimination falling between 22 and 32‰ (corresponding to KIEs of 1.023 and 1.033, respectively; Table 2). This is higher than the 15‰ typically cited in the literature (e.g., Robinson, 2001; Werner and Schmidt, 2002). The relatively high ¹⁵N discrimination observed here

may partially derive from differences in how the measurements were made. Other work has generally focused on *in vivo* discrimination (e.g., Mariotti et al., 1982) or *in vitro* experiments using isolated organelles or cytosolic fractions (e.g., Ledgard et al., 1985). *In vitro* experiments with purified enzymes typically produce greater isotopic discrimination than experiments

Table 2 | Kinetic isotope effects (KIEs) for the five tested NRs.

¹⁵ N enrichment	<i>Arabidopsis</i>	Yeast	<i>Z. mays</i>	<i>A. niger</i>	<i>E. coli</i>
0.37%	1.023 (0.001)	1.023 (0.002)	1.031 (0.001)	1.028 (0.003)	1.032 (0.003)
0.5%	1.029 (0.002)				
1%	1.034 (0.001)				
10%	1.052 (0.001)	1.051 (0.001)	1.056 (0.001)		
25%	1.053 (0.002)	1.053 (0.002)	1.058 (0.001)		
50%	1.075 (0.002)	1.068 (0.001)	1.071 (0.002)	1.078 (0.004)	1.078 (0.002)
75%	1.278 (0.018)	1.179 (0.010)	1.186 (0.020)		
99%	4.483 (0.317)	3.470 (0.536)	2.723 (0.026)	3.957 (0.450)	3.482 (0.345)

Standard errors are presented in parentheses. With highly ¹⁵N enriched substrate pools, it is more appropriate to assess isotope discrimination as the KIE. Increasing ¹⁵N enrichment results in larger deviation from natural abundance KIEs and greater variation in the measurements.





FIGURE 1 | Observed ¹⁵N concentration in N₂O across a range of substrate ¹⁵N enrichments. The N₂O was produced from reduction of NO₂⁻ to N₂O via *S. nitritireducens* cultures (bacterially derived) or via sulfamic acid (chemically derived). Also plotted are the expected ¹⁵N concentrations. The bacterial and chemical data were calculated using Equation (5), and both approaches used the same NO₂⁻ substrate. The expected plot assumes a constant ¹⁵N discrimination of -29.44‰. Error bars represent the standard errors of the means ($n = 5$).

Table 1 | Observed ¹⁵N natural abundance isotope discrimination (Δ in ‰, $n = 6-10$) for the five tested NRs.

¹⁵ N enrichment	<i>Arabidopsis</i>	Yeast	<i>Z. mays</i>	<i>A. niger</i>	<i>E. coli</i>
0.37%	22.5 (0.8)	23.0 (0.5)	30.9 (1.5)	28.1 (0.4)	31.6 (2.3)

Standard errors are presented in parentheses.



with cellular organelles, multiple enzymes, or whole organisms because additional factors diminish ^{15}N discrimination as the catalytic system becomes more complex (Werner and Schmidt, 2002). We performed *in vitro* assays using purified NRs and electron donors, providing unlimited substrate, and operating under optimal conditions to prevent subsequent conversion of the product NO_2^- . Olleros-Izard (1983; cited in Schmidt and Medina, 1991) used a similar approach with purified corn NR and observed a discrimination of 30%, which agrees with our values for non-recombinant enzymes. Karsh et al. (2012) using purified enzymes and the bacterial reduction method found that the assimilatory NR from *A. niger* discriminated against $^{15}\text{NO}_3^-$ by $26.6 \pm 0.2\%$ (Karsh et al., 2012), again a value similar to ours.

The four assimilatory eukaryotic NRs (*Arabidopsis*, yeast, maize, and *A. niger*) are structurally similar, although the two recombinant enzymes (*Arabidopsis* and yeast) lack a portion of the complete native enzyme. For example, the recombinant *Arabidopsis* NR, which is natively homotetrameric, is missing in three of its four subunits the Mo-MPT cofactor, and thus the NO_3^- reducing site (Skipper et al., 2001). Such differences in structure may account for the lower ^{15}N discrimination observed in the recombinant NRs than in the maize and *A. niger* enzymes (Table 1). Some of the observed difference between the *Arabidopsis* and maize NRs may be due to differences between monocot and eudicot NRs; however, there is no published evidence for this possibility. Differences in glycosylation between native enzymes and recombinant enzymes expressed in *P. pastoris* also might affect enzymatic activity and possibly isotope discrimination (e.g., Henriksson et al., 2003).

Despite the structural disparity (Moreno-Vivián et al., 1999) between the eukaryotic and prokaryotic NRs, the *E. coli* NR did not grossly differ in NR ^{15}N discrimination from the other NRs tested.

The electron donor did not strongly affect ^{15}N discrimination. The plant-derived NRs (*Arabidopsis* and maize) used NADH as an electron donor, while the fungal NRs used NADPH. The similar level of discrimination that we found despite the difference donors supports the hypothesis that substantial structural differences among the eukaryotic enzymes are largely localized to the FAD binding domain (Campbell, 1999; Karsh et al., 2012). The dissimilatory *E. coli* NR used BV as an artificial electron donor, which provides electrons to the Mo-MGD cofactor directly, in contrast to the biological donor quinol, which provides electrons through the FeS or cytochrome c cofactors. No studies have measured the effects of the biological electron donor on ^{15}N discrimination in nitrate reduction in prokaryotes, and it is not clear whether the use of a different electron donor would affect ^{15}N discrimination. Nonetheless, use of an artificial electron donor (e.g., methyl viologen) rather than NADH had little to no influence on eukaryotic NR ^{15}N discrimination (Karsh et al., 2012).

ESTIMATION OF NR DISCRIMINATION USING ENRICHED ^{15}N

The systematic and large deviation from expected values of the KIEs at high ^{15}N enrichment (particularly > 50 atom %; Figure 2) cannot derive from substrate dependent changes in the rate limiting nature of N-O bond cleavage because the observed KIEs are greater than their theoretical limits (Tcherkez and Farquhar, 2006). Our measurements at high ^{15}N enrichment were affected by artifacts. Similar artifacts may have affected the ^{15}N pulse labeling experiment in our previous work (Bloom et al., 2010). The following discusses some of these potential artifacts.

First, although the sequential bacterial reduction method to assess ^{15}N discrimination at natural abundance ^{15}N worked very well, we observed increasingly large standard errors and anomalously low ^{15}N enrichment in the N_2O derived from NO_3^- reduction by *P. chlororaphis* as ^{15}N NO_3^- enrichment increased (Supplemental Table 2). At enrichments of 99 atom %, the N_2O derived from the NO_2^- was substantially more enriched in ^{15}N than the N_2O derived from NO_3^- (97.6 vs. 58.3 atom % ^{15}N ; Supplemental Table 2). If we used a different NO_3^- reducing bacterium, *Pseudomonas aureofaciens* (Sigman et al., 2001), the observed differences were even larger (97.6 vs. 7.9 atom %; Supplemental Table 2). By contrast, *S. nitritireducens* produced N_2O from NO_2^- with ^{15}N enrichment comparable to that produced via chemical reduction (Figure 1). These results suggest that the sequential bacterial reduction method should be limited to ^{15}N enrichments of less than 10 atom %. The reason for the variation among the bacterial strains is not clear, although ages of the bacterial strains or possible bacterial contamination may play a role. NO_3^- concentrations in the spent bacterial reduction solutions contained no measurable NO_3^- after reduction, so the reaction was apparently quantitative (data not shown).

Second, instrument performance and the mathematical models currently available for the calculation of ^{15}N in N_2O at high levels of ^{15}N enrichment are likely responsible for some of the observed variation. Using an IRMS rather than using an elemental analyzer coupled with a mass spectrometer contributed to the observed error, particularly at ^{15}N enrichments greater than 50%, however, the use of the IRMS was necessary to measure NR discrimination at natural abundance ^{15}N concentrations. High

enrichments of ¹⁵N lead to problems with IRMS collector saturation, and we could not measure the 44 peak necessary for the calculation of the mass ratios.

The results from highly ¹⁵N enriched substrates presented here illustrate a more general methodological issue regarding stable N isotopes: one should avoid the use of highly ¹⁵N enriched substrates in discrimination studies, particularly if small experimental changes in ¹⁵N concentration are expected. High ¹⁵N enrichments exacerbate methodological difficulties in sample preparation, measurement, and instrument calibration. This ultimately increases the final experimental error and unnecessarily confounds experimental outcomes.

Indeed, the problems in the use of highly enriched ¹⁵N encountered here may affect measurements at larger scales. For example, it is common in agricultural field studies to use enriched ¹⁵N as a tracer for the cycling of N among different soil pools. Experiments conducted using enriched ¹⁵N tracers consistently estimate lower N cycling rates than those conducted using natural abundance or non-isotopic methods (e.g., Hauck and Bremner, 1976; Androsoff et al., 1995; Ladha et al., 2005). Other factors beyond the measurement issues described above certainly play a role in the observed differences among these approaches, but artifacts linked to the use of high ¹⁵N enrichments such as those observed here may contribute to these differences.

In conclusion, despite the use of different (i.e., natural and artificial) electron donors, NR ¹⁵N discrimination at natural abundance levels of ¹⁵N varied between 22 to 32‰ among four assimilatory eukaryotic NRs as well as a prokaryotic dissimilatory NR. These discriminations are similar to those observed or reported in other recent studies (Tcherkez and Farquhar, 2006; Karsh et al., 2012). Difficulties encountered using heavily ¹⁵N-enriched samples demonstrate the undesirability of protocols requiring such labeling for the measurement of enzyme isotope discrimination.

ACKNOWLEDGMENTS

This work was supported by NSF-IOS-08-18435, NSF-IOS-13-58675, and the National Research Initiative Competitive Grant 2008-35100-04459 from the USDA National Institute of Food and Agriculture. We would like to thank Hsien Easlon for his comments and critical review and Joy Mathews for her assistance with the instrumentation. Finally we would like to thank two anonymous reviewers for their critical reviews of the manuscript. No conflict of interest, financial or otherwise, exists for any of the authors.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://www.frontiersin.org/Journal/10.3389/fpls.2014.00317/abstract>

REFERENCES

- Adams, M. W. W., and Mortenson, L. E. (1982). The effect of cyanide and ferricyanide on the activity of the dissimilatory nitrate reductase of *Escherichia coli*. *J. Biol. Chem.* 257, 1791–1799.
- Amundson, R., Austin, A. T., Schuur, E. A. G., Yoo, K., Matzek, V., Kendall, C., et al. (2003). Global patterns of the isotopic composition of soil and plant nitrogen. *Glob. Biogeochem. Cycles* 17, 1031. doi: 10.1029/2002GB001903
- Androsoff, G. L., van Kessel, C., and Pennock, D. J. (1995). Landscape-scale estimates of dinitrogen fixation by *Pisum sativum* by nitrogen-15 natural abundance and enriched isotope dilution. *Biol. Fertil. Soils* 20, 33–40. doi: 10.1007/BF00307838
- Barbier, G. G., and Campbell, W. H. (2005). Viscosity effects on eukaryotic nitrate reductase activity. *J. Biol. Chem.* 280, 26049–26054. doi: 10.1074/jbc.M409694200
- Barbier, G. G., Joshi, J. C., Campbell, E. R., and Campbell, W. H. (2004). Purification and biochemical characterization of simplified eukaryotic nitrate reductase expressed in *Pichia pastoris*. *Protein Expr. Purif.* 37, 61–71. doi: 10.1016/j.pep.2004.05.021
- Berks, B. C., Ferguson, S. J., Moir, J. W. B., and Richardson, D. J. (1995). Enzymes and associated electron transport systems that catalyse the respiratory reduction of nitrogen oxides and oxyanions. *Biochim. Biophys. Acta* 1232, 97–173. doi: 10.1016/0005-2728(95)00092-5
- Bigeleisen, J., and Wolfsberg, M. (1958). “Theoretical and experimental aspects of isotope effects in chemical kinetics,” in *Advances in Chemical Physics*, Vol. 1, eds I. Prigogine and P. Debye (Hoboken, NJ: John Wiley and Sons, Inc), 15–76.
- Bloom, A. J., Burger, M., Asensio, J. S. R., and Cousins, A. B. (2010). Carbon dioxide enrichment inhibits nitrate assimilation in wheat and *Arabidopsis*. *Science* 328, 899–903. doi: 10.1126/science.1186440
- Böhlke, J. K., Smith, R. L., and Hannon, J. E. (2007). Isotopic analysis of N and O in nitrite and nitrate by sequential selective bacterial reduction to N₂O. *Anal. Chem.* 79, 5888–5895. doi: 10.1021/ac070176k
- Campbell, W. H. (1999). Nitrate reductase structure, function and regulation: bridging the gap between biochemistry and physiology. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 50, 277–303. doi: 10.1146/annurev.arplant.50.1.277
- Cleland, W. W. (2005). The use of isotope effects to determine enzyme mechanisms. *Arch. Biochem. Biophys.* 433, 2–12. doi: 10.1016/j.abb.2004.08.027
- Epstein, E., and Bloom, A. J. (2005). *Mineral Nutrition of Plants: Principles and Perspectives, 2nd Edn.* Sunderland, MA: Sinauer Associates.
- Evans, R. D. (2001). Physiological mechanisms influencing plant nitrogen isotope composition. *Trends Plant Sci.* 6, 121–126. doi: 10.1016/S1360-1385(01)01889-1
- Evans, R. D., Bloom, A. J., Sukrapanna, S. S., and Ehrlinger, J. R. (1996). Nitrogen isotope composition of tomato (*Lycopersicon esculentum* Mill. cv. T-5) grown under ammonium or nitrate nutrition. *Plant Cell Environ.* 19, 1317–1323. doi: 10.1111/j.1365-3040.1996.tb00010.x
- Farquhar, G. D., Ehleringer, J. R., and Hubick, K. T. (1989). Carbon isotope discrimination and photosynthesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 40, 503–537. doi: 10.1146/annurev.pp.40.060189.002443
- Fischer, K., Barbier, G. G., Hecht, H. J., Mendel, R. R., Campbell, W. H., and Schwarz, G. (2005). Structural basis of eukaryotic nitrate reduction: crystal structure of the nitrate reductase active site. *Plant Cell* 17, 1167–1179. doi: 10.1105/tpc.104.029694
- Fry, B. (2006). *Stable Isotope Ecology*. New York, NY: Springer.
- Gilliam, M. B., Sherman, M. P., Griscavage, J. M., and Ignarro, L. J. (1993). A spectrophotometric assay for nitrate using NADPH oxidation by *Aspergillus* nitrate reductase. *Anal. Biochem.* 212, 359–365. doi: 10.1006/abio.1993.1341
- Granger, J., and Sigman, D. (2009). Removal of nitrite with sulfamic acid for nitrate N and O isotope analysis with the denitrifier method. *Rapid Commun. Mass Spectrom.* 23, 3753–3762. doi: 10.1002/rcm.4307
- Granger, J., Sigman, D. M., Needoba, J. A., and Harrison, P. J. (2004). Coupled nitrogen and oxygen isotope fractionation of nitrate during assimilation by cultures of marine phytoplankton. *Limnol. Oceanogr.* 49, 1763–1773. doi: 10.4319/lo.2004.49.5.1763
- Hauck, R. D., and Bremner, J. M. (1976). Use of tracers for soil and fertilizer nitrogen research. *Adv. Agron.* 28, 219–266. doi: 10.1016/S0065-2113(08)60556-8
- Henriksson, H., Denman, S. E., Campuzano, I. D. G., Ademark, P., Master, E. R., Teeri, T. T., et al. (2003). N-linked glycosylation of native and recombinant cauliflower xyloglucan endotransglycosylase 16A. *Biochem. J.* 375, 61–73. doi: 10.1042/BJ20030485
- Hipkin, C. R., Ali, A. H., and Cannons, A. (1986). Structure and properties of assimilatory nitrate reductase from the yeast *Candida nitratophila*. *J. Gen. Microbiol.* 132, 1997–2003. doi: 10.1099/00221287-132-7-1997
- Kaiser, J., Rockmann, T., and Brenninkmeijer, C. A. M. (2003). Complete and accurate mass spectrometric isotope analysis of tropospheric nitrous oxide. *J. Geophys. Res.* 108, 1–17. doi: 10.1029/2003JD003613

- Karsh, K. L., Granger, J. L., Kritec, K., and Sigman, D. M. (2012). Eukaryotic assimilatory nitrate reductase fractionates N and O isotopes with a ratio near unity. *Environ. Sci. Technol.* 46, 5727–5735. doi: 10.1021/es204593q
- Kolb, K. J., and Evans, R. D. (2003). Influence of nitrogen source and concentration on nitrogen isotopic discrimination in two barley genotypes (*Hordeum vulgare* L.). *Plant Cell Environ.* 26, 1431–1440. doi: 10.1046/j.1365-3040.2003.01066.x
- Ladha, J. K., Pathak, H., Krupnik, T. J., Six, J., and van Kessel, C. (2005). Efficiency of fertilizer nitrogen in cereal production: retrospects and prospects. *Adv. Agron.* 87, 85–156. doi: 10.1016/S0065-2113(05)87003-8
- Ledgard, S. E., Woo, K. C., and Bergersen, F. J. (1985). Isotopic fractionation during reduction of nitrate and nitrite by extracts of spinach leaves. *Aust. J. Plant Physiol.* 12, 631–640. doi: 10.1071/PP9850631
- Mariotti, A., Mariotti, F., Champigny, M.-L., Amarger, N., and Moysé, A. (1982). Nitrogen isotope fractionation associated with nitrate reductase activity and uptake of NO₃⁻ by Pearl Millet. *Plant Physiol.* 69, 880–884. doi: 10.1104/pp.69.4.880
- Miranda, K. M., Espey, M. G., and Wink, D. A. (2001). A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide* 5, 62–71. doi: 10.1006/niox.2000.0319
- Moreno-Vivián, C., Cabello, P., Martínez-Luque, M., Blasco, R., and Castillo, F. (1999). Prokaryotic nitrate reduction: molecular properties and functional distinction among bacterial nitrate reductases. *J. Bacteriol.* 181, 6573–6584.
- Morozkina, E. V., Nosikov, A. N., Zvyagilskaya, R. A., and L'Vov, N. P. (2005). Isoation purification, and characterization of nitrate reductase from a salt-tolerant *Rhodotorula glutinis* yeast strain grown in the presence of tungsten. *Biochemistry (Mosc.)* 70, 809–814. doi: 10.1007/s10541-005-0188-y
- Needoba, J. A., Sigman, D. M., and Harrison, P. J. (2004). The mechanism of isotope fractionation during algal nitrate assimilation as illuminated by the ¹⁵N/¹⁴N of intracellular nitrate. *J. Phycol.* 40, 517–522. doi: 10.1111/j.1529-8817.2004.03172.x
- Nielsen, M., Larsen, L. H., Jetten, M. S. M., and Revsbech, N. P. (2004). Bacterium-based NO₂⁻ biosensor for environmental applications. *Appl. Environ. Microbiol.* 70, 6551–6558. doi: 10.1128/AEM.70.11.6551-6558.2004
- Olleros-Izard, T. (1983). *Kinetische Isotopeneffekte der Arginase und Nitratreduktase Reaktion: Ein Beitrag zur Aufklärung der Entsprechenden Reaktionsmechanismen*. Ph.D. thesis, Technischen Universität München, Freising-Weihenstephan.
- Robinson, D. (2001). ⁸¹⁵N as an integrator of the nitrogen cycle. *Trends Ecol. Evol.* 16, 153–162. doi: 10.1016/S0169-5347(00)02098-X
- Robinson, D., Handley, L. L., and Scrimgeour, C. M. (1998). A theory for ¹⁵N/¹⁴N fractionation in nitrate-grown vascular plants. *Planta* 205, 397–406. doi: 10.1007/s004250050336
- Schlesinger, W. H. (1997). *Biogeochemistry: An Analysis of Global Change*. San Diego, CA: Academic Press.
- Schmidt, H. L., and Medina, R. (1991). Possibilities and scope of the double isotope effect method in the elucidation of mechanisms of enzyme catalyzed reactions. *Isotopenpraxis* 27, 1–4. doi: 10.1080/10256019108622450
- Shearer, G., and Kohl, D. H. (1986). N₂-fixation in field settings: estimations based on natural ¹⁵N abundance. *Aust. J. Plant Physiol.* 13, 699–756.
- Sigman, D. M., Casciotti, K. L., Andreani, M., Barford, C. M., Galanter, M., and Bohlke, J. K. (2001). A bacterial method for the nitrogen isotopic analysis of nitrate in seawater and freshwater. *Anal. Chem.* 73, 4145–4153. doi: 10.1021/ac010088e
- Skipper, L., Campbell, W. H., Mertens, J. A., and Lowe, D. J. (2001). Pre-steady state kinetic analysis of recombinant Arabidopsis NADH:nitrate reductase: rate-limiting processes in catalysis. *J. Biol. Chem.* 276, 26995–27002. doi: 10.1074/jbc.M100356200
- Sprent, J. (1979). *The Biology of Nitrogen Fixation*. New York, NY: McGraw Hill.
- Stevens, R. J., and Laughlin, R. J. (1998). Measurement of nitrous oxide and di-nitrogen emissions from agricultural soils. *Nutr. Cycling Agroecosyst.* 52, 131–139. doi: 10.1023/A:1009715807023
- Stickrod, R. D. 2nd., and Marshall, J. D. (2000). On-line nitrate-^δ¹⁵N extracted from groundwater determined by continuous-flow elemental analyzer/isotope ratio mass spectrometry. *Rapid Commun. Mass Spectrom.* 14, 1266–1268. doi: 10.1002/1097-0231(20000730)14:14<1266::AID-RCM21>3.0.CO;2-R
- Su, W., Mertens, J. A., Kanamaru, K., Campbell, W. H., and Crawford, N. M. (1997). Analysis of wild-type and mutant plant nitrate reductase expressed in the methylotrophic yeast *Pichia pastoris*. *Plant Physiol.* 115, 1135–1143. doi: 10.1104/pp.115.3.1135
- Tcherkez, G., and Farquhar, G. D. (2006). Isotopic fractionation by plant nitrate reductase, twenty years later. *Funct. Plant Biol.* 33, 531–537. doi: 10.1071/FP05284
- Temple, C. A., and Rajagopalan, K. V. (2000). Mechanism of assembly of the bis(molybdopterin guanine dinucleotide) molybdenum cofactor in *Rhodobacter sphaeroides* dimethyl sulfoxide reductase. *J. Biol. Chem.* 275, 40202–40210. doi: 10.1074/jbc.M007407200
- Werner, R. A., and Schmidt, H. L. (2002). The *in vivo* nitrogen isotope discrimination among organic plant compounds. *Phytochemistry* 61, 465–484. doi: 10.1016/S0031-9422(02)00204-2

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 29 January 2014; accepted: 15 June 2014; published online: 02 July 2014.
Citation: Carlisle E, Yarnes C, Toney MD and Bloom AJ (2014) Nitrate reductase ¹⁵N discrimination in *Arabidopsis thaliana*, *Zea mays*, *Aspergillus niger*, *Picea angusta*, and *Escherichia coli*. *Front. Plant Sci.* 5:317. doi: 10.3389/fpls.2014.00317
This article was submitted to *Plant Physiology*, a section of the journal *Frontiers in Plant Science*.

Copyright © 2014 Carlisle, Yarnes, Toney and Bloom. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.