



# $\omega$ 3 Polyunsaturated Fatty Acids as Immunomodulators in Colorectal Cancer: New Potential Role in Adjuvant Therapies

Stefania Miccadei<sup>1</sup>, Roberta Masella<sup>2</sup>, Anna Maria Mileo<sup>1</sup> and Sandra Gessani<sup>3\*</sup>

<sup>1</sup> Unit of Tumor Immunology and Immunotherapy, Department of Research, Advanced Diagnostic and Technological Innovation, Regina Elena National Cancer Institute, Rome, Italy, <sup>2</sup> Department of Veterinary Public Health and Food Safety, Istituto Superiore di Sanità, Rome, Italy, <sup>3</sup> Department of Hematology, Oncology and Molecular Medicine, Istituto Superiore di Sanità, Rome, Italy

## OPEN ACCESS

### Edited by:

José Mordoh,  
Fundación Instituto Leloir, Argentina

### Reviewed by:

Christine E. Loscher,  
Dublin City University, Ireland  
Chryssostomos Chatgililoglu,  
National Research Council (CNR), Italy

### \*Correspondence:

Sandra Gessani  
sandra.gessani@iss.it

### Specialty section:

This article was submitted to  
Immunotherapies and Vaccines,  
a section of the journal  
Frontiers in Immunology

**Received:** 27 June 2016

**Accepted:** 21 October 2016

**Published:** 15 November 2016

### Citation:

Miccadei S, Masella R, Mileo AM and Gessani S (2016)  $\omega$ 3 Polyunsaturated Fatty Acids as Immunomodulators in Colorectal Cancer: New Potential Role in Adjuvant Therapies. *Front. Immunol.* 7:486. doi: 10.3389/fimmu.2016.00486

Diet composition may affect the onset and progression of chronic degenerative diseases, including cancer, whose pathogenesis relies on inflammatory processes. Growing evidence indicates that diet and its components critically contribute to human health, affecting the immune system, secretion of adipokines, and metabolic pathways. Colorectal cancer (CRC) is one of the leading causes of death worldwide. Antineoplastic drugs are widely used for CRC treatment, but drug resistance and/or off-target toxicity limit their efficacy. Dietary  $\omega$ 3 polyunsaturated fatty acids (PUFA) have been gaining great interest in recent years as possible anti-inflammatory and anticancer agents, especially in areas such as the large bowel, where the pro-inflammatory context promotes virtually all steps of colon carcinogenesis. Growing epidemiological, experimental, and clinical evidence suggests that  $\omega$ 3 PUFA may play a role in several stages of CRC management exhibiting antineoplastic activity against human CRC cells, improving the efficacy of radiation and chemotherapy, ameliorating cancer-associated secondary complications, and preventing CRC recurrence. These effects are most likely related to the immunomodulatory activities of  $\omega$ 3 PUFA that are able to influence several aspects of the inflammatory process ranging from inflammasome activation, leukocyte recruitment, production of immune mediators to differentiation, and activation of immune cells. In this review, we will focus on the potential use of  $\omega$ 3 PUFA as adjuvant agents together with chemo/radiotherapy, highlighting the immunomodulatory effects most likely responsible for their beneficial effects in different stages of CRC management.

**Keywords:**  $\omega$ 3 polyunsaturated fatty acids, immunomodulation, colorectal cancer, adjuvant therapy, inflammation

## INTRODUCTION

It has become increasingly clear that dietary habits may affect the risk/progression of chronic diseases with a pathogenic inflammatory component, such as colorectal cancer (CRC) (1). CRC is the third most common cancer and the fourth leading cause of cancer death worldwide and represents a clinical model of cancer showing a close relationship with inflammation and environmental factors (2). Inflammation is nowadays considered an hallmark of cancer (3), and exacerbated inflammatory processes has been reported to favor the insurgence of several types of neoplastic

diseases, their progression, as well as a worse prognosis for the patient with cancer (World Cancer Report, 2014, IARC). Several types of inflammation differing by cause, mechanism, outcome, and intensity can promote cancer development and progression. Interestingly, rising CRC incidence trends are mainly regarded as a part of the change in lifestyle, including calorically excessive high-fat/low-fiber diet, lack of physical activity, and obesity (4). Epidemiological and clinical investigations have consistently evidenced a significant relationship between obesity-driven inflammation and most steps of CRC development, including initiation, promotion, progression, and metastasis (1). Furthermore, cancer therapy as well as surgery for cancer resection can trigger an inflammatory response by causing trauma, necrosis, and tissue injury that stimulate tumor re-emergence and resistance to therapy (5). In spite of the progress made in recent years in cancer treatment, the clinical relevance of the conventional therapeutic modalities alone in solid tumors, including CRC, still pose major questions due to the high costs of drugs and unsatisfactory long-term clinical results (6). Thus, due to the impact of inflammation as a risk factor for CRC development as well as in the course of the disease, new concepts are emerging on the benefits that may derive from the combination of conventional CRC treatments with intervention strategies targeted at inhibiting/attenuating chronic inflammation. Notably, dietary components influence inflammatory processes including intestinal inflammation, and dietary habits have been suggested to play a major role in CRC risk (7). Considerable attention has therefore been directed toward the ability of nutritional agents to target key molecular pathways involved in cancer-related inflammation. It has only recently gained acceptance that fatty acids (FAs) are major determinants in inflammation and may influence the risk of related pathologies. Given their precursor status to signaling lipid mediators, FAs act as ligands for key immune receptors, and their activation can initiate and perpetuate an innate immune response (8).

## DIETARY POLYUNSATURATED FATTY ACIDS

$\omega$ 3 and  $\omega$ 6 FAs are polyunsaturated fatty acids (PUFA) whose definition derives from counting the carbon atoms after the last double bond in the end of the fatty acid chain (9). The type of response they induce strongly depends on their biochemical properties, mainly related to number and position of double bonds (10). In this regard,  $\omega$ 3 and  $\omega$ 6 PUFA exhibit different behaviors with respect to the inflammatory process, exerting anti-inflammatory or pro-inflammatory activity, respectively. Owing to the proposed competitive role of  $\omega$ 3 and  $\omega$ 6 PUFA, their dietary composition has been suggested to be a biologically plausible target for CRC management.  $\omega$ 3 and  $\omega$ 6 FAs represent naturally occurring substances required for biological process with key roles in phospholipid membrane structure and function, cellular signaling, and lipid metabolism (11, 12). In humans, the  $\omega$ 3  $\alpha$ -linolenic acid (ALA, 18:3 $\omega$ 3) as well as the  $\omega$ 6 linoleic acid (LA, 18:2 $\omega$ 6), precursors of several PUFA, are essential FA that cannot be synthesized by humans and must therefore be assumed from dietary sources including fish,

flaxseeds, walnuts, and algae (9). Several kind of food, rich in  $\omega$ 3 PUFA, are typical of the so-called Mediterranean diet (MD). Several epidemiological studies have evaluated the effects of this dietary pattern as protective against several diseases associated with chronic low-grade inflammation such as cancer, diabetes, obesity, atherosclerosis, metabolic syndrome, and cognition disorders (13). Adherence to traditional MD was found to be associated with markedly and significantly reduced incidence of cardiovascular diseases and overall cancer (14, 15). Due to the poor endogenous conversion of ALA to its longer and more highly desaturated FAs, there is considerable debate as to whether the longer versions such as eicosapentaenoic acid (EPA, 20:5 $\omega$ 3) and docosahexaenoic acid (DHA, 22:6 $\omega$ 3) may be considered semi-essential (16). These PUFA are key components of cell membrane, lipoproteins, and adipose tissue (AT) whose composition closely reflects the dietary intake (17). For several years,  $\omega$ 3 PUFA have been studied extensively for the prevention of cardiovascular diseases (18, 19). More recently, their capacity to exert anticancer activity has been recognized. A plethora of studies pointed to  $\omega$ 3 PUFA and their oxidative metabolites as potential anti-inflammatory and anticancer agents, especially in areas such as the large bowel, where the influence of orally introduced substances is high and tumors show deranged PUFA patterns. In particular, modified  $\omega$ 3 and  $\omega$ 6 PUFA profiles have been found in serum and cancer tissue of CRC patients, suggesting possible alteration of PUFA metabolism that may play a role in inflammation-driven colorectal carcinogenesis (20, 21). Overall, these studies suggested that the metabolism of PUFA plays a role in inflammation-driven colorectal carcinogenesis and is likely influenced by the tumor size. In this regard, it is of relevance that  $\omega$ 3/ $\omega$ 6 PUFA ratio of the industrialized diets is imbalanced, and this condition results in a well-known pro-inflammatory effect (22). Noteworthy, a decreased  $\omega$ 3/ $\omega$ 6 PUFA ratio of visceral AT correlates with AT inflammatory status in CRC patients suggesting that qualitative, more than quantitative, PUFA changes in AT may influence tissue dysfunctions potentially linked to inflammatory conditions (23). Likewise, the sum of some PUFA [i.e., gamma linolenic acid (GLA), Di-homo-gamma linolenic acid (DGLA), and arachidonic acid (AA)] in both visceral and subcutaneous AT was found statistically higher in cancer patients with respect to healthy controls (24). Overall, these observations support the concept that PUFA derangement occurs not only in tumor tissue but also at microenvironment and systemic levels most likely contributing to tumor-associated inflammation.

There is now sufficient evidence from prospective studies indicating that fish consumption or fish oil  $\omega$ 3 PUFA intake is inversely related to CRC incidence in the general population (25). However, the results achieved in a recent meta-analysis carried out to evaluate the effects of  $\omega$ 3 PUFA on inflammatory markers in CRC patients highlighted that the beneficial effects of  $\omega$ 3 PUFA rely on supplementation protocols (e.g., dose, duration, and route of administration), the concomitant anticancer regimen adopted, as well as on the confounding effects due to other nutrients that might be present in the supplement (26). In this regard, it should be taken into account that the observational studies carried out to evaluate the potential anticancer effects of

$\omega$ 3 PUFA might have been limited by the insufficient homogeneity of the observations. An explanation for such heterogeneity relies likely on the inherent difficulties associated with epidemiology, including the confounding and dietary pattern context, measurement error, level of intake, and genetic polymorphisms (26, 27).

Despite the majority of studies has proven that  $\omega$ 3 PUFA administration is not toxic or carcinogenic, it is hard to imagine that these compounds could be exploited alone in the treatment of cancer. However, growing epidemiological, experimental, and clinical evidence showing the beneficial effects of  $\omega$ 3 PUFA in several stages of CRC management represent a solid rationale for their exploitation in therapeutic combinatory strategies (28). The antineoplastic effects of  $\omega$ 3 PUFA have been studied for many years, and multiple biological mechanisms have been suggested to explain their inhibitory effects on cancer cell growth (29, 30). However, in recent years, growing attention has been given to the antineoplastic activity of  $\omega$ 3 PUFA related to the action of these bioactive molecules on inflammatory process, due to the pathogenic role of inflammation in the development of many kinds of tumors.  $\omega$ 3 PUFA, particularly EPA and DHA, have emerged as nutrients capable of modulating both metabolic and immune processes. As summarized in **Table 1**, these PUFA are able to modulate a number of aspects of the inflammatory process ranging from inflammasome

activation, leukocyte recruitment, and production of immune mediators, to differentiation and activation of immune cells.  $\omega$ 3 PUFA effects are mediated by different mechanisms including both extracellular (e.g., GPR120) (31) and intracellular (e.g., PPAR $\gamma$ ) receptors that control inflammatory cell signaling and gene expression (32). In spite of many efforts, it remains to be fully defined how these compounds succeed in influencing such a variety of molecular pathways and cell/tissue functions with pleiotropic beneficial effects, and how these mechanisms are interconnected. Some of the  $\omega$ 3 PUFA-mediated events appear to rely, at least in part, on changes in FA composition of membrane phospholipids, relevant for the production of lipid mediators, and formation of lipid rafts in response to inflammatory stimuli (33).

## PROTECTIVE ANTI-INFLAMMATORY EFFECTS OF $\omega$ 3 PUFA IN COLORECTAL CANCER

Considerable debate still exists on the mechanisms underlying the protective effects exerted by the  $\omega$ 3 PUFA on the establishment/progression of inflammation and cancer (29). Furthermore, the precise mechanism that operates at specific body district, such as the intestine, remains to be defined. Several mechanisms have

**TABLE 1 | Immunomodulatory effects of  $\omega$ 3 PUFA in monocyte/macrophages and T lymphocytes.**

$\omega$ 3 PUFA exposure	Experimental/animal model	Observed effect	Suggested mechanism	Reference
DHA <sup>a</sup> , EPA <sup>a</sup>	MU and HU monocytic cell lines; MU and HU primary monocyte/M $\phi$ , C57BL/6 mice	↓ inflammasome activation	GPR120- $\beta$ arrestin-2-mediated NF- $\kappa$ B inhibition Enhanced autophagy Impaired TLR2/TLR1 dimerization	(34–36)
Isocaloric HFD-containing 27% menhaden fish oil (16% EPA, 9% DHA); isocaloric $\omega$ 3 PUFA-enriched diet (3% menhaden fish-oil + 7% safflower oil)	C57BL/6 mice	↓ leukocyte chemotaxis	GPR120, $\beta$ arrestin-2 NF- $\kappa$ B and STAT3 inhibition	(37, 38)
DHA <sup>a</sup> , EPA <sup>a</sup>	MU monocytic cell lines and primary M $\phi$	↑ efferocytic activity of M $\phi$	PPAR $\gamma$ and AKT activation	(39, 40)
DHA <sup>a</sup> , EPA <sup>a</sup> , isocaloric HFD-containing 27% menhaden fish oil (16% EPA, 9% DHA); HFD + DHA <sup>a</sup>	MU monocytic cell lines and primary M $\phi$ C57BL/6 mice	↑ M2 M $\phi$ polarization	GPR120, $\beta$ arrestin-2	(37, 40–42)
DHA <sup>a</sup> , EPA <sup>a</sup> , $\omega$ 3 PUFA-enriched diet <sup>b</sup> , DHA-enriched diet (omegavie DHA90 TG)	BALB/c and C57BL/6 mice MU spleen and bone marrow DC	↓ T cell activation ↑ Th2 lymphocyte polarization ↓ Th17 lymphocyte polarization ↑ expression and generation of T <sub>reg</sub> ↓ T <sub>reg</sub> suppressive and migratory function ↓ DC activation ↓ DC-mediated T cell activation ↑ DC-mediated T <sub>reg</sub> expansion	Interference with STAT3 signaling pathway <i>via</i> enhancement of PPAR $\gamma$ -induced SOCS3 M2 M $\phi$ mediated	(43–51)

The table shows a summary of the main publications on the immunomodulatory effects of  $\omega$ 3 PUFA in monocyte/macrophages and T lymphocytes, including both *in vitro* studies and *in vivo* animal models.

MU, murine; HU, human; M $\phi$ , macrophage; DC, dendritic cell; HFD, high fat diet; GPR120, G protein-coupled receptor 120.

<sup>a</sup>Chemically synthesized.

<sup>b</sup> $\omega$ 3 PUFA source: perilla seed oil, menhaden fish oil, and EPAX oil (EPAX-7010) containing ~85%  $\omega$ 3 PUFA (70% EPA, 12% DHA).

been proposed to explain the capacity of  $\omega$ 3 PUFA to interfere with important steps of CRC carcinogenesis including alterations of the cellular redox state and modulation of membrane dynamics, and surface receptor function which regulate cell proliferation and apoptosis. Due to the close association of CRC with inflammation, the capacity of  $\omega$ 3 PUFA to modulate some aspects of inflammation and immune response is becoming increasingly important. In this regard, it is of interest that  $\omega$ 3 PUFA inhibit agonist-induced activation of pattern recognition receptors, thus dampening inflammation (52). However, only few studies focused on the role of PUFA-mediated immunomodulatory effects in the therapy of CRC.

Due to the abundance of double bonds, EPA and DHA are susceptible to oxidation and can undergo spontaneous non-enzymatic peroxidation or enzymatic oxidation giving rise to a variety of bioactive lipids that induce inflammation, tumorigenesis, and thrombosis, as well as to mediators with antitumorogenic, pro-resolution properties (53, 54). One of the main mechanisms by which  $\omega$ 3 PUFA exert their anti-inflammatory activity involves enzymatic pathways that are hyper-activated in CRC (55), such as cyclooxygenase (COX) and lipoxygenase (LOX), and leads to prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) production, a potent pro-inflammatory and procarcinogenic agent. EPA and DHA have been shown to inhibit COX-2 expression in CRC cells (56), and this effect has been attributed to  $\omega$ 3 PUFA capacity to displace  $\omega$ 6 PUFA, in particular AA, from membranes. This metabolic competition is due to structural similarity of EPA and AA, which compete with each other for the same enzymes, COX, and LOX, a very important event in tumor development (57). Furthermore, supplementation with  $\omega$ 3 PUFA significantly decreases AA-derived eicosanoids, while increasing EPA-derived eicosanoids, shifting the balance toward a reduced level of inflammation (58). Interestingly, aspirin, a non-steroidal anti-inflammatory drug, possesses both COX and LOX inhibitory action, and enhances the production of the anti-inflammatory lipid mediator lipoxin A<sub>4</sub>. Likewise, anti-inflammatory compounds are formed from EPA and DHA by the action of aspirin termed as resolvins (from EPA and DHA) and protectins and maresins from DHA (59, 60). Of note, regular aspirin use substantially reduces the risk of CRC (61). Pro-inflammatory cytokines and tumor-infiltrating myeloid and immune cells play critical roles in almost every developmental stages of inflammation-related cancers, from initiation, promotion, and progression to malignant metastasis (1). A recently published meta-analysis of the  $\omega$ 3 PUFA effects on inflammatory markers highlighted a decreased level of some inflammatory mediators (cytokines or acute phase proteins) in CRC patients who received  $\omega$ 3 PUFA supplementation (26). Likewise, a recent *in vitro* study described a new anti-inflammatory and anticancer properties of DHA. Fluckiger and colleagues reported that inhibition of colon cancer growth and activation of apoptosis by DHA involve autocrine production of TNF $\alpha$  via microRNA (miR)-21 (62).

Another mechanism involved in anti-inflammatory and potentially antineoplastic effects of  $\omega$ 3 PUFA regards the possibility that these compounds may modulate epigenetically (through DNA methylation, acetylation modifications, and

miR gene regulation) the expression of crucial genes involved in cellular processes associated with colorectal carcinogenesis (63–65). More recently, the epigenetic regulation of gene expression and the polarization of macrophages toward pro-resolving M2 phenotype have emerged as novel mechanisms explaining the antineoplastic effects of  $\omega$ 3 PUFA at the colon level, although additional studies are needed for a deeper comprehension of this issue (30). Finally, the capacity of  $\omega$ 3 PUFA to modulate inflammatory pathways and to generate lipid mediators critical for the resolution of inflammation (e.g., resolvins, protectins, and maresins) has gained attention as a main mechanism involved in the beneficial effects of  $\omega$ 3 PUFA against CRC (66).

## $\omega$ 3 PUFA AS POTENTIAL ADJUVANTS IN THE THERAPY OF CRC

Fatty acids may influence carcinogenesis through various mechanisms. EPA and DHA are incorporated into cellular membranes and through the production of lipid mediators and may exert anticancer properties by affecting gene expression or activating signal transduction molecules involved in the control of cell proliferation, differentiation, apoptosis, and metastasis. These properties of  $\omega$ 3 PUFA suggest that they will have important therapeutic potential in cancer management.

Patients undergoing surgery are at risk of developing complications in the postoperative period partly caused by changes in the immune response following surgery (67). Thus, initially, a hyper-inflammatory response followed by a phase of relative immune incompetence occurs in relation to major surgery (68). Patients who undergo surgery for CRC have a 30% risk of infectious complications, and anastomotic leakages are seen in as many as 15% of patients (69).

Initial studies indicated that administration of PUFA-enriched diets leads to increased incorporation of EPA and DHA not only in liver and gut mucosa tissue but also in tumor tissue in patients with solid tumors of the upper gastrointestinal tract (70). This observation suggested that preoperative administration of oral PUFA-enriched diets could have an impact on the postoperative inflammatory response after major abdominal surgery. Subsequently, in a prospective randomized, double-blind, single-center, placebo-controlled study, it was demonstrated a higher total marine  $\omega$ 3 PUFA content in the colonic mucosa, but not in the muscular layer, after 7 days of oral EPA + DHA supplementation to patients admitted for elective CRC surgery (71). Furthermore, providing  $\omega$ 3 PUFA daily for 7 days before surgery resulted in a significant decrease in the formation of the pro-inflammatory leukotriene B<sub>4</sub> (LTB<sub>4</sub>) from neutrophils with a simultaneous increased production of LTB<sub>5</sub>. However, the clinical consequences of these changes are still unknown as no correlations between values of LTB<sub>4</sub> or LTB<sub>4</sub>/LTB<sub>5</sub>, and postoperative complication rates were found (72). Likewise, the higher incorporation of EPA, DHA, and docosapentaenoic acid in granulocytes of the  $\omega$ 3 PUFA-enriched supplement group of CRC patients was not associated with improved postoperative outcomes (69). Thus, the effects of  $\omega$ 3 PUFA on preoperative complications need to



be investigated in larger trials and for a longer period (months) of  $\omega$ 3 PUFA intake to clearly establish whether LTB formation from activated neutrophils is/is not an important determinant of surgical complications.

Several studies highlighted the beneficial effect of  $\omega$ 3 PUFA as chemopreventive and chemotherapeutic agents in the treatment of several chronic pathologies including cancer providing evidence for a potential use of these compounds to enhance chemo/radiotherapy efficacy and reduce the risk of tumor recurrence (73). Furthermore, several studies demonstrated that EPA and DHA, having immunomodulatory capacity, act to reduce inflammation, even when associated with chemo/radiotherapy immune suppression in lung (74, 75), and in other cancers (73).

In particular, Xue and colleagues evaluated whether  $\omega$ 3 PUFA supplementation could affect the efficacy of a cyclical combination treatment with Irinotecan (CPT-11)/5-fluorouracil (5-FU) in a rat CRC model. Administration of a diet enriched with  $\omega$ 3 PUFA prior to initiating chemotherapy, inhibited tumor growth, while, during chemotherapy,  $\omega$ 3 PUFA-enriched diet enhanced tumor chemosensitivity and reduced body weight loss with respect to control diet fed rats (76). Other studies addressed the issue of whether supplementation with  $\omega$ 3 PUFA to CRC patients undergoing chemotherapy could induce changes in inflammation markers, such as the ratio of C reactive protein (CRP) to albumin, considered a relevant clinical/inflammatory

marker. In this respect, some clinical trials demonstrated that EPA and DHA supplementation during chemotherapeutic treatments improved CRP values, CRP/albumin status, and positively modulated the nutritional status of CRC patients (77, 78). Likewise, evaluation of clinical outcomes during and after chemotherapy in CRC patients supplemented with fish oil containing EPA and DHA during the first 9 weeks of treatment unraveled that the time to tumor progression was significantly longer in the supplemented with respect to the control group (79). Interestingly, a recent phase II double-blind randomized, placebo-controlled trial of EPA administration to advanced CRC patients undergoing liver resection surgery demonstrated that EPA is incorporated into secondary CRC tissue and has systemic anti-inflammatory activity. In particular, a clear-cut decrease of the urinary metabolite PGE-M, reflecting systemic PGE<sub>2</sub> levels, as well as a reversible reduction in NF- $\kappa$ B binding in peripheral blood leukocytes was observed in EPA-treated patients with respect to the placebo group (80). Overall, these observations that supplementation may represent an innovative combination therapy strategy contributing to delay tumor progression and likely enhancing the antineoplastic activity of chemotherapeutic drugs. The design and main hallmarks of clinical trials performed on CRC patients supplemented with  $\omega$ 3 PUFA, described above, are summarized in **Table 2**.

In the last years, many studies have highlighted the presence of cancer stem cells (CSC) in most tumors, including CRC.

**TABLE 2 | Clinical effects of  $\omega$ 3 PUFA in CRC patients.**

$\omega$ 3 PUFA daily treatment	Surgery	n Subjects	Timing/duration	Chemotherapy	Observed effect	Outcome measures	Reference
1 g DHA + 2 g EPA orally	Yes	148	7 days pre-surgery	No	Potential beneficial effect on local immune function	$\omega$ 3 PUFA content in the colonic mucosa and muscular layer	(71)
1 g DHA + 2 g EPA orally	Yes	148	7 days pre-surgery	No	Anti-inflammatory effects 5-HEPE $\uparrow$ LTB4/%-HETE $\downarrow$	Inflammatory markers	(72)
1 g DHA + 2 g EPA orally	Yes	148	7 days pre- and 7 days post-surgery	No	No significant difference between group in infectious or non-infectious post-operative complications	Levels of $\omega$ 3 PUFA into granulocytes	(69)
0.6 g EPA/DHA (fish oil) orally	No	23	63 days	5-FU + irinotecan + folinic acid	Positive modulation of nutritional status. CRP/albumin $\downarrow$	Evaluation of nutritional status and inflammatory markers	(77)
0.6 g EPA/DHA (fish oil) orally	No	11	63 days	CAPE + OXA + 5-FU + leucovorin	Improved CRP values, CRP/albumin status, plasma fatty acid profile, and potentially prevented weight loss during treatment	Evaluation of inflammatory markers and nutritional status	(78)
0.6 g EPA/DHA (fish oil) orally	No	30	63 days	Standard chemotherapy	Delayed tumor progression time by enhancing the antineoplastic action of chemotherapy	Evaluation of tumor progression time and CEA values after chemotherapy	(79)
2 g EPA-FFA <sup>a</sup> orally	Yes	88	Median 30 days pre-operative	No	Preoperative treatment may have prolonged benefit on post-operative overall and disease-free survival	Ki67, CD31, and PGE2 levels. Survival statistical analysis (overall and disease-free survival)	(80)

<sup>a</sup>ALFA (SLA Pharma AG, Liestel, Switzerland).

5-HEPE, 5-hydroxyeicosatetraenoic acid; LTB4, leukotriene B4; 5-HETE, 5-hydroxyeicosatetraenoic acid; 5-FU, 5-fluorouracil; CAPE, capecitabine (Xeloda); OXA, oxaloplatin; CRP, C reactive protein; CEA, carcinoembryonic antigen; PGE2, prostaglandin E.2.

These cells are considered responsible for tumor relapse and resistance to conventional therapies (81). Thus, novel anticancer strategies have been designed to selectively target CSC, and to this purpose, natural compounds, including  $\omega$ 3 PUFA, might have a relevant role. In colon cancer stem-like cells (CSLC), De Carlo et al. showed that EPA has a pre-differentiating effect and this finding could, at least in part, explain the increased cellular sensitivity to 5-FU (82). In keeping with these observations, Yang and colleagues demonstrated that EPA and DHA, in a single or combined treatment, enhanced the chemotherapeutic sensitivity effect of 5-FU and mitomycin C by inducing apoptotic cell death of CSLC (83).

## CONCLUSION

Although the use of  $\omega$ 3 PUFA as potential chemopreventive and chemotherapeutic agents has been frequently proposed in cancer treatment, the evidence achieved points to these compounds as important agents for combined strategies against cancer. In particular, despite some contradictory results, several

experimental and clinical data strongly support a role of  $\omega$ 3 PUFA as adjuvants able to counteract the occurrence of inflammatory processes during CRC treatment as well as to increase chemotherapy efficacy or to decrease its toxicity. However, none of the studies has taken into consideration the initial status of the patients prior to PUFA supplementation or dietary advice, a relevant issue to better exploit the preventive/therapeutic/adjuvant potential of these compounds. Further and deepened studies are thus needed to characterize the real  $\omega$ 3 PUFA effectiveness and mechanisms of actions.

## AUTHOR CONTRIBUTIONS

All authors listed have made substantial, direct, and intellectual contribution to the work and approved it for publication.

## FUNDING

This study was supported by a grant of the Italian Association for Cancer Research (AIRC) project (IG 2013 No. 14185) to SG.

## REFERENCES

- Pietrzyk L, Torres A, Maciejewski R, Torres K. Obesity and obese-related chronic low-grade inflammation in promotion of colorectal cancer development. *Asian Pac J Cancer Prev* (2015) 16(10):4161–8. doi:10.7314/APJCP.2015.16.10.4161
- Favoriti P, Carbone G, Greco M, Pirozzi F, Pirozzi RE, Corcione F. Worldwide burden of colorectal cancer: a review. *Updates Surg* (2016) 68(1):7–11. doi:10.1007/s13304-016-0359-y
- Colotta F, Allavena P, Sica A, Garlanda C, Mantovani A. Cancer-related inflammation, the seventh hallmark of cancer: links to genetic instability. *Carcinogenesis* (2009) 30(7):1073–81. doi:10.1093/carcin/bgp127
- Morrison DS, Parr CL, Lam TH, Ueshima H, Kim HC, Jee SH, et al. Behavioural and metabolic risk factors for mortality from colon and rectum cancer: analysis of data from the Asia-Pacific Cohort Studies Collaboration. *Asian Pac J Cancer Prev* (2013) 14(2):1083–7. doi:10.7314/APJCP.2013.14.2.1083
- Grivennikov SI, Greten FR, Karin M. Immunity, inflammation, and cancer. *Cell* (2010) 140(6):883–99. doi:10.1016/j.cell.2010.01.025
- Ramakrishnan R, Gabrilovich DI. Novel mechanism of synergistic effects of conventional chemotherapy and immune therapy of cancer. *Cancer Immunol Immunother* (2013) 62(3):405–10. doi:10.1007/s00262-012-1390-6
- Turati F, Edefonti V, Bravi F, Ferraroni M, Talamini R, Giacosa A, et al. Adherence to the European food safety authority's dietary recommendations and colorectal cancer risk. *Eur J Clin Nutr* (2012) 66(4):517–22. doi:10.1038/ejcn.2011.217
- Wiktorowska-Owczarek A, Berezinska M, Nowak JZ. PUFAs: structures, metabolism and functions. *Adv Clin Exp Med* (2015) 24(6):931–41. doi:10.17219/acem/31243
- Ferreri C, Chatgililoglu C. *Membrane Lipidomics for Personalized Health*. Chichester: Wiley & Sons Ltd. (2015).
- Das UN. Essential fatty acids – a review. *Curr Pharm Biotechnol* (2006) 7(6):467–82. doi:10.2174/138920106779116856
- Chapkin RS, McMurray DN, Davidson LA, Patil BS, Fan YY, Lupton JR. Bioactive dietary long-chain fatty acids: emerging mechanisms of action. *Br J Nutr* (2008) 100(6):1152–7. doi:10.1017/S0007114508992576
- Akhtar Khan N. Polyunsaturated fatty acids in the modulation of T-cell signalling. *Prostaglandins Leukot Essent Fatty Acids* (2010) 82(4–6):179–87. doi:10.1016/j.plefa.2010.02.023
- Casas R, Sacanella E, Estruch R. The immune protective effect of the Mediterranean diet against chronic low-grade inflammatory diseases. *Endocr*
- Metab Immune Disord Drug Targets* (2014) 14(4):245–54. doi:10.2174/1871530314666140922153350
- Benetou V, Trichopoulos A, Orfanos P, Naska A, Lagiou P, Boffetta P, et al. Conformity to traditional Mediterranean diet and cancer incidence: the Greek EPIC cohort. *Br J Cancer* (2008) 99(1):191–5. doi:10.1038/sj.bjc.6604418
- Mena MP, Sacanella E, Vazquez-Agell M, Morales M, Fito M, Escoda R, et al. Inhibition of circulating immune cell activation: a molecular antiinflammatory effect of the Mediterranean diet. *Am J Clin Nutr* (2009) 89(1):248–56. doi:10.3945/ajcn.2008.26094
- Qi K, Hall M, Deckelbaum RJ. Long-chain polyunsaturated fatty acid accretion in brain. *Curr Opin Clin Nutr Metab Care* (2002) 5(2):133–8. doi:10.1097/00075197-200203000-00003
- Calder PC, Yaqoob P. Omega-3 polyunsaturated fatty acids and human health outcomes. *Biofactors* (2009) 35(3):266–72. doi:10.1002/biof.42
- Geelen A, Schouten JM, Kamphuis C, Stam BE, Burema J, Renkema JM, et al. Fish consumption, n-3 fatty acids, and colorectal cancer: a meta-analysis of prospective cohort studies. *Am J Epidemiol* (2007) 166(10):1116–25. doi:10.1093/aje/kwm197
- Mozaffarian D, Wu JH. Omega-3 fatty acids and cardiovascular disease: effects on risk factors, molecular pathways, and clinical events. *J Am Coll Cardiol* (2011) 58(20):2047–67. doi:10.1016/j.jacc.2011.06.063
- Wang S, Xie J, Li H, Yang K. Differences of polyunsaturated fatty acid in patients with colorectal cancer and healthy people. *J Cancer Res Ther* (2015) 11(2):459–63. doi:10.4103/0973-1482.147702
- Yang K, Li H, Dong J, Dong Y, Wang CZ. Expression profile of polyunsaturated fatty acids in colorectal cancer. *World J Gastroenterol* (2015) 21(8):2405–12. doi:10.3748/wjg.v21.i8.2405
- Simopoulos AP. The importance of the omega-6/omega-3 fatty acid ratio in cardiovascular disease and other chronic diseases. *Exp Biol Med (Maywood)* (2008) 233(6):674–88. doi:10.3181/0711-MR-311
- D'Archivio M, Scaccocchio B, Giammarioli S, Fiani ML, Vari R, Santangelo C, et al. omega-3-PUFAs exert anti-inflammatory activity in visceral adipocytes from colorectal cancer patients. *PLoS One* (2013) 8(10):e77432. doi:10.1371/journal.pone.0077432
- Giuliani A, Ferrara F, Scimo M, Angelico F, Olivieri L, Basso L. Adipose tissue fatty acid composition and colon cancer: a case-control study. *Eur J Nutr* (2014) 53(4):1029–37. doi:10.1007/s00394-013-0605-8
- Hall MN, Chavarro JE, Lee IM, Willett WC, Ma J. A 22-year prospective study of fish, n-3 fatty acid intake, and colorectal cancer risk in men. *Cancer Epidemiol Biomarkers Prev* (2008) 17(5):1136–43. doi:10.1158/1055-9965.EPI-07-2803

26. Mocellin MC, Camargo CQ, Nunes EA, Fiates GM, Trindade EB. A systematic review and meta-analysis of the n-3 polyunsaturated fatty acids effects on inflammatory markers in colorectal cancer. *Clin Nutr* (2016) 35(2):359–69. doi:10.1016/j.clnu.2015.04.013
27. Gerber M. Omega-3 fatty acids and cancers: a systematic update review of epidemiological studies. *Br J Nutr* (2012) 107(Suppl 2):S228–39. doi:10.1017/S0007114512001614
28. Serini S, Ottes Vasconcelos R, Fasano E, Calviello G. How plausible is the use of dietary n-3 PUFA in the adjuvant therapy of cancer? *Nutr Res Rev* (2016) 29(1):102–25. doi:10.1017/S0954422416000044
29. Cockbain AJ, Toogood GJ, Hull MA. Omega-3 polyunsaturated fatty acids for the treatment and prevention of colorectal cancer. *Gut* (2012) 61(1):135–49. doi:10.1136/gut.2010.233718
30. Serini S, Ottes Vasconcelos R, Fasano E, Calviello G. Epigenetic regulation of gene expression and M2 macrophage polarization as new potential omega-3 polyunsaturated fatty acid targets in colon inflammation and cancer. *Expert Opin Ther Targets* (2016) 20(7):843–58. doi:10.1517/14728222.2016.1139085
31. Cheshmehkani A, Senatorov IS, Kandi P, Singh M, Britt A, Hayslett R, et al. Fish oil and flax seed oil supplemented diets increase FFAR4 expression in the rat colon. *Inflamm Res* (2015) 64(10):809–15. doi:10.1007/s00011-015-0864-3
32. Robbins GT, Nie D. PPAR gamma, bioactive lipids, and cancer progression. *Front Biosci (Landmark Ed)* (2012) 17:1816–34. doi:10.2741/4021
33. Shaikh SR. Biophysical and biochemical mechanisms by which dietary N-3 polyunsaturated fatty acids from fish oil disrupt membrane lipid rafts. *J Nutr Biochem* (2012) 23(2):101–5. doi:10.1016/j.jnutbio.2011.07.001
34. Snodgrass RG, Huang S, Choi IW, Rutledge JC, Hwang DH. Inflammation-mediated secretion of IL-1beta in human monocytes through TLR2 activation; modulation by dietary fatty acids. *J Immunol* (2013) 191(8):4337–47. doi:10.4049/jimmunol.1300298
35. Yan Y, Jiang W, Spinetti T, Tardivel A, Castillo R, Bourquin C, et al. Omega-3 fatty acids prevent inflammation and metabolic disorder through inhibition of NLRP3 inflammasome activation. *Immunity* (2013) 38(6):1154–63. doi:10.1016/j.immuni.2013.05.015
36. Williams-Bey Y, Boullaran C, Vural A, Huang NN, Hwang IY, Shan-Shi C, et al. Omega-3 free fatty acids suppress macrophage inflammasome activation by inhibiting NF-kappaB activation and enhancing autophagy. *PLoS One* (2014) 9(6):e97957. doi:10.1371/journal.pone.0097957
37. Oh DY, Talukdar S, Bae EJ, Imamura T, Morinaga H, Fan W, et al. GPR120 is an omega-3 fatty acid receptor mediating potent anti-inflammatory and insulin-sensitizing effects. *Cell* (2010) 142(5):687–98. doi:10.1016/j.cell.2010.07.041
38. Monk JM, Liddle DM, De Boer AA, Brown MJ, Power KA, Ma DW, et al. Fish-oil-derived n-3 PUFAs reduce inflammatory and chemotactic adipokine-mediated cross-talk between co-cultured murine splenic CD8+ T cells and adipocytes. *J Nutr* (2015) 145(4):829–38. doi:10.3945/jn.114.205443
39. Li S, Sun Y, Liang CP, Thorp EB, Han S, Jehle AW, et al. Defective phagocytosis of apoptotic cells by macrophages in atherosclerotic lesions of ob/ob mice and reversal by a fish oil diet. *Circ Res* (2009) 105(11):1072–82. doi:10.1161/CIRCRESAHA.109.199570
40. Chang HY, Lee HN, Kim W, Surh YJ. Docosahexaenoic acid induces M2 macrophage polarization through peroxisome proliferator-activated receptor gamma activation. *Life Sci* (2015) 120:39–47. doi:10.1016/j.lfs.2014.10.014
41. Titos E, Rius B, Gonzalez-Periz A, Lopez-Vicario C, Moran-Salvador E, Martinez-Clemente M, et al. Resolvin D1 and its precursor docosahexaenoic acid promote resolution of adipose tissue inflammation by eliciting macrophage polarization toward an M2-like phenotype. *J Immunol* (2011) 187(10):5408–18. doi:10.4049/jimmunol.1100225
42. De Boer AA, Monk JM, Robinson LE. Docosahexaenoic acid decreases pro-inflammatory mediators in an in vitro murine adipocyte macrophage co-culture model. *PLoS One* (2014) 9(1):e85037. doi:10.1371/journal.pone.0085037
43. Mizota T, Fujita-Kambara C, Matsuya N, Hamasaki S, Fukudome T, Goto H, et al. Effect of dietary fatty acid composition on Th1/Th2 polarization in lymphocytes. *JPN J Parenter Enteral Nutr* (2009) 33(4):390–6. doi:10.1177/0148607108325252
44. Yessoufou A, Ple A, Moutairou K, Hichami A, Khan NA. Docosahexaenoic acid reduces suppressive and migratory functions of CD4+CD25+ regulatory T-cells. *J Lipid Res* (2009) 50(12):2377–88. doi:10.1194/jlr.M900101-JLR200
45. Kong W, Yen JH, Ganea D. Docosahexaenoic acid prevents dendritic cell maturation, inhibits antigen-specific Th1/Th17 differentiation and suppresses experimental autoimmune encephalomyelitis. *Brain Behav Immun* (2011) 25(5):872–82. doi:10.1016/j.bbi.2010.09.012
46. Berger H, Vegran F, Chikh M, Gilardi F, Ladoire S, Bugaut H, et al. SOCS3 transactivation by PPARgamma prevents IL-17-driven cancer growth. *Cancer Res* (2013) 73(12):3578–90. doi:10.1158/0008-5472.CAN-12-4018
47. Monk JM, Hou TY, Turk HF, McMurray DN, Chapkin RS. n3 PUFAs reduce mouse CD4+ T-cell ex vivo polarization into Th17 cells. *J Nutr* (2013) 143(9):1501–8. doi:10.3945/jn.113.178178
48. Teague H, Rockett BD, Harris M, Brown DA, Shaikh SR. Dendritic cell activation, phagocytosis and CD69 expression on cognate T cells are suppressed by n-3 long-chain polyunsaturated fatty acids. *Immunology* (2013) 139(3):386–94. doi:10.1111/imm.12088
49. Carlsson JA, Wold AE, Sandberg AS, Ostman SM. The polyunsaturated fatty acids arachidonic acid and docosahexaenoic acid induce mouse dendritic cells maturation but reduce T-cell responses in vitro. *PLoS One* (2015) 10(11):e0143741. doi:10.1371/journal.pone.0143741
50. Han SC, Koo DH, Kang NJ, Yoon WJ, Kang JG, Kang HK, et al. Docosahexaenoic acid alleviates atopic dermatitis by generating tregs and IL-10/TGF-beta-modified macrophages via a TGF-beta-dependent mechanism. *J Invest Dermatol* (2015) 135(6):1556–64. doi:10.1038/jid.2014.488
51. Lian M, Luo W, Sui Y, Li Z, Hua J. Dietary n-3 PUFA protects mice from con a induced liver injury by modulating regulatory T cells and PPAR-gamma expression. *PLoS One* (2015) 10(7):e0132741. doi:10.1371/journal.pone.0132741
52. Lee JY, Zhao L, Hwang DH. Modulation of pattern recognition receptor-mediated inflammation and risk of chronic diseases by dietary fatty acids. *Nutr Rev* (2010) 68(1):38–61. doi:10.1111/j.1753-4887.2009.00259.x
53. Tuncer S, Banerjee S. Eicosanoid pathway in colorectal cancer: recent updates. *World J Gastroenterol* (2015) 21(41):11748–66. doi:10.3748/wjg.v21.i41.11748
54. Roy J, Le Guennec JY, Galano JM, Thireau J, Bultel-Ponce V, Demion M, et al. Non-enzymatic cyclic oxygenated metabolites of omega-3 polyunsaturated fatty acid: bioactive drugs? *Biochimie* (2016) 120:56–61. doi:10.1016/j.biochi.2015.06.010
55. Furstemberger G, Krieg P, Muller-Decker K, Habenicht AJ. What are cyclooxygenases and lipoxygenases doing in the driver's seat of carcinogenesis? *Int J Cancer* (2006) 119(10):2247–54. doi:10.1002/ijc.22153
56. Calviello G, Di Nicuolo F, Gragnoli S, Piccioni E, Serini S, Maggiano N, et al. n-3 PUFAs reduce VEGF expression in human colon cancer cells modulating the COX-2/PGE2 induced ERK-1 and -2 and HIF-1alpha induction pathway. *Carcinogenesis* (2004) 25(12):2303–10. doi:10.1093/carcin/bgh265
57. Habel P, Weylandt KH, Lichopoj K, Nowak J, Purschke M, Wang JD, et al. Docosahexaenoic acid suppresses arachidonic acid-induced proliferation of LS-174T human colon carcinoma cells. *World J Gastroenterol* (2009) 15(9):1079–84. doi:10.3748/wjg.15.1079
58. Dupertuis YM, Meguid MM, Pichard C. Colon cancer therapy: new perspectives of nutritional manipulations using polyunsaturated fatty acids. *Curr Opin Clin Nutr Metab Care* (2007) 10(4):427–32. doi:10.1097/MCO.0b013e3281e2c9d4
59. Serhan CN, Gotlinger K, Hong S, Arita M. Resolvins, docosatrienes, and neuroprotectins, novel omega-3-derived mediators, and their aspirin-triggered endogenous epimers: an overview of their protective roles in catabasis. *Prostaglandins Other Lipid Mediat* (2004) 73(3–4):155–72. doi:10.1016/j.prostaglandins.2004.03.005
60. Poorani R, Bhatt AN, Dwarakanath BS, Das UN. COX-2, aspirin and metabolism of arachidonic, eicosapentaenoic and docosahexaenoic acids and their physiological and clinical significance. *Eur J Pharmacol* (2016) 785:116–32. doi:10.1016/j.ejphar.2015.08.049
61. Giovannucci E, Egan KM, Hunter DJ, Stampfer MJ, Colditz GA, Willett WC, et al. Aspirin and the risk of colorectal-cancer in women. *N Engl J Med* (1995) 333(10):609–14. doi:10.1056/Nejm199509073331001
62. Fluckiger A, Dumont A, Derangere V, Rebe C, de Rosny C, Causse S, et al. Inhibition of colon cancer growth by docosahexaenoic acid involves autocrine

- production of TNF $\alpha$ . *Oncogene* (2016) 35(35):4611–22. doi:10.1038/onc.2015.523
63. Cho Y, Turner ND, Davidson LA, Chapkin RS, Carroll RJ, Lupton JR. A chemoprotective fish oil/pectin diet enhances apoptosis via Bcl-2 promoter methylation in rat azoxymethane-induced carcinomas. *Exp Biol Med (Maywood)* (2012) 237(12):1387–93. doi:10.1258/ebm.2012.012244
  64. Cho Y, Turner ND, Davidson LA, Chapkin RS, Carroll RJ, Lupton JR. Colon cancer cell apoptosis is induced by combined exposure to the n-3 fatty acid docosahexaenoic acid and butyrate through promoter methylation. *Exp Biol Med (Maywood)* (2014) 239(3):302–10. doi:10.1177/1535370213514927
  65. Tsoukas MA, Ko BJ, Witte TR, Dincer F, Hardman WE, Mantzoros CS. Dietary walnut suppression of colorectal cancer in mice: mediation by miRNA patterns and fatty acid incorporation. *J Nutr Biochem* (2015) 26(7):776–83. doi:10.1016/j.jnutbio.2015.02.009
  66. Janakiram NB, Mohammed A, Rao CV. Role of lipoxins, resolvins, and other bioactive lipids in colon and pancreatic cancer. *Cancer Metastasis Rev* (2011) 30(3–4):507–23. doi:10.1007/s10555-011-9311-2
  67. Kehlet H, Dahl JB. Anaesthesia, surgery, and challenges in postoperative recovery. *Lancet* (2003) 362(9399):1921–8. doi:10.1016/S0140-6736(03)14966-5
  68. Calder PC. n-3 fatty acids, inflammation, and immunity – relevance to post-surgical and critically ill patients. *Lipids* (2004) 39(12):1147–61. doi:10.1007/s11745-004-1342-z
  69. Sorensen LS, Thorlacius-Ussing O, Schmidt EB, Rasmussen HH, Lundbye-Christensen S, Calder PC, et al. Randomized clinical trial of perioperative omega-3 fatty acid supplements in elective colorectal cancer surgery. *Br J Surg* (2014) 101(2):33–42. doi:10.1002/bjs.9361
  70. Senkal M, Haaker R, Linseisen J, Wolfram G, Homann HH, Stehle P. Preoperative oral supplementation with long-chain omega-3 fatty acids beneficially alters phospholipid fatty acid patterns in liver, gut mucosa, and tumor tissue. *JPEN J Parenter Enteral Nutr* (2005) 29(4):236–40. doi:10.1177/0148607105029004236
  71. Sorensen LS, Rasmussen HH, Aardestrup IV, Thorlacius-Ussing O, Lindorff-Larsen K, Schmidt EB, et al. Rapid incorporation of omega-3 fatty acids into colonic tissue after oral supplementation in patients with colorectal cancer: a randomized, placebo-controlled intervention trial. *JPEN J Parenter Enteral Nutr* (2014) 38(5):617–24. doi:10.1177/0148607113491782
  72. Sorensen LS, Thorlacius-Ussing O, Rasmussen HH, Lundbye-Christensen S, Calder PC, Lindorff-Larsen K, et al. Effects of perioperative supplementation with omega-3 fatty acids on leukotriene B(4) and leukotriene B(5) production by stimulated neutrophils in patients with colorectal cancer: a randomized, placebo-controlled intervention trial. *Nutrients* (2014) 6(10):4043–57. doi:10.3390/nu6104043
  73. Calviello G, Serini S, Piccioni E, Pessina G. Antineoplastic effects of n-3 polyunsaturated fatty acids in combination with drugs and radiotherapy: preventive and therapeutic strategies. *Nutr Cancer* (2009) 61(3):287–301. doi:10.1080/01635580802582777
  74. van der Meij BS, Langius JA, Smit EF, Spreeuwenberg MD, von Blomberg BM, Heijboer AC, et al. Oral nutritional supplements containing (n-3) polyunsaturated fatty acids affect the nutritional status of patients with stage III non-small cell lung cancer during multimodality treatment. *J Nutr* (2010) 140(10):1774–80. doi:10.3945/jn.110.121202
  75. Finocchiaro C, Segre O, Fadda M, Monge T, Scigliano M, Schena M, et al. Effect of n-3 fatty acids on patients with advanced lung cancer: a double-blind, placebo-controlled study. *Br J Nutr* (2012) 108(2):327–33. doi:10.1017/S0007114511005551
  76. Xue H, Le Roy S, Sawyer MB, Field CJ, Dieleman LA, Baracos VE. Single and combined supplementation of glutamine and n-3 polyunsaturated fatty acids on host tolerance and tumour response to 7-ethyl-10-[4-(1-piperidino)-1-piperidino]carbonyloxy-camptothecin (CPT-11)/5-fluorouracil chemotherapy in rats bearing Ward colon tumour. *Br J Nutr* (2009) 102(3):434–42. doi:10.1017/S0007114508199482
  77. Silva Jde A, Trindade EB, Fabre ME, Menegotto VM, Gevaerd S, Buss Zda S, et al. Fish oil supplement alters markers of inflammatory and nutritional status in colorectal cancer patients. *Nutr Cancer* (2012) 64(2):267–73. doi:10.1080/01635581.2012.643133
  78. Mocellin MC, Pastore e Silva Jde A, Camargo Cde Q, Fabre ME, Gevaerd S, Naliwaiko K, et al. Fish oil decreases C-reactive protein/albumin ratio improving nutritional prognosis and plasma fatty acid profile in colorectal cancer patients. *Lipids* (2013) 48(9):879–88. doi:10.1007/s11745-013-3816-0
  79. Camargo Cde Q, Mocellin MC, Pastore Silva Jde A, Fabre ME, Nunes EA, Trindade EB. Fish oil supplementation during chemotherapy increases posterior time to tumor progression in colorectal cancer. *Nutr Cancer* (2016) 68(1):70–6. doi:10.1080/01635581.2016.1115097
  80. Cockbain AJ, Volpato M, Race AD, Munarini A, Fazio C, Belluzzi A, et al. Anticolorectal cancer activity of the omega-3 polyunsaturated fatty acid eicosapentaenoic acid. *Gut* (2014) 63(11):1760–8. doi:10.1136/gutjnl-2013-306445
  81. Stuckey DW, Shah K. Stem cell-based therapies for cancer treatment: separating hope from hype. *Nat Rev Cancer* (2014) 14(10):683–91. doi:10.1038/nrc3798
  82. De Carlo F, Witte TR, Hardman WE, Claudio PP. Omega-3 eicosapentaenoic acid decreases CD133 colon cancer stem-like cell marker expression while increasing sensitivity to chemotherapy. *PLoS One* (2013) 8(7):e69760. doi:10.1371/journal.pone.0069760
  83. Yang T, Fang S, Zhang HX, Xu LX, Zhang ZQ, Yuan KT, et al. N-3 PUFAs have antiproliferative and apoptotic effects on human colorectal cancer stem-like cells in vitro. *J Nutr Biochem* (2013) 24(5):744–53. doi:10.1016/j.jnutbio.2012.03.023

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Miccadei, Masella, Mileo and Gessani. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.