



# The Multiple Functions of Common Microbial Carbon Polymers, Glycogen and PHB, during Stress Responses in the Non-Diazotrophic Cyanobacterium *Synechocystis* sp. PCC 6803

Ramon Damrow<sup>1</sup>, Iris Maldener<sup>2</sup> and Yvonne Zilliges<sup>1,3\*</sup>

<sup>1</sup> Section of Plant Biochemistry, Institute of Biology, Humboldt-Universität zu Berlin, Berlin, Germany, <sup>2</sup> Section of Organismic Interactions, Interfaculty Institute of Microbiology and Infection Medicine, Eberhard Karls Universität Tübingen, Tübingen, Germany, <sup>3</sup> Section of Biophysics and Photosynthesis, Institute of Physics, Freie Universität Berlin, Berlin, Germany

## OPEN ACCESS

### Edited by:

Robert Eugene Blankenship,  
Washington University in St. Louis,  
USA

### Reviewed by:

Alberto A. Iglesias,  
Universidad Nacional del Litoral,  
Consejo Nacional de Investigaciones  
Científicas y Técnicas, Argentina  
Jason Warren Cooley,  
University of Missouri, USA

### \*Correspondence:

Yvonne Zilliges  
yvonne.zilliges@fu-berlin.de

### Specialty section:

This article was submitted to  
Microbial Physiology and Metabolism,  
a section of the journal  
Frontiers in Microbiology

**Received:** 01 March 2016

**Accepted:** 03 June 2016

**Published:** 21 June 2016

### Citation:

Damrow R, Maldener I and Zilliges Y  
(2016) The Multiple Functions  
of Common Microbial Carbon  
Polymers, Glycogen and PHB, during  
Stress Responses  
in the Non-Diazotrophic  
Cyanobacterium *Synechocystis* sp.  
PCC 6803. *Front. Microbiol.* 7:966.  
doi: 10.3389/fmicb.2016.00966

Classical microbial carbon polymers such as glycogen and polyhydroxybutyrate (PHB) have a crucial impact as both a sink and a reserve under macronutrient stress conditions. Most microbial species exclusively synthesize and degrade either glycogen or PHB. A few bacteria such as the phototrophic model organism *Synechocystis* sp. PCC 6803 surprisingly produce both physico-chemically different polymers under conditions of high C to N ratios. For the first time, the function and interrelation of both carbon polymers in non-diazotrophic cyanobacteria are analyzed in a comparative physiological study of single- and double-knockout mutants ( $\Delta glgC$ ;  $\Delta phaC$ ;  $\Delta glgC/\Delta phaC$ ), respectively. Most of the observed phenotypes are explicitly related to the knockout of glycogen synthesis, highlighting the metabolic, energetic, and structural impact of this process whenever cells switch from an active, photosynthetic 'protein status' to a dormant 'glycogen status'. The carbon flux regulation into glycogen granules is apparently crucial for both phycobilisome degradation and thylakoid layer disassembly in the presence of light. In contrast, PHB synthesis is definitely not involved in this primary acclimation response. Moreover, the very weak interrelations between the two carbon-polymer syntheses indicate that the regulation and role of PHB synthesis in *Synechocystis* sp. PCC 6803 is different from glycogen synthesis.

**Keywords:** glycogen, PHB, metabolic spilling, cyanobacteria, nitrogen chlorosis, thylakoid disassembly

## INTRODUCTION

All microorganisms accumulate carbon biopolymers, namely glycogen and/or poly- $\beta$ -hydroxybutyrate (PHB), which act as cellular sinks as well as stable and yet readily accessible reservoirs for carbon and energy, to acclimate and to cope with starvation conditions, in particular nitrogen starvation leading to high C-to-N ratios of nutrients (Allen, 1984). Glycogen is a homogeneous, water-soluble polyglucan composed of 9–13 (1–4)-linked  $\alpha$ -D-glucose residues that are interlinked via (1–6)- $\alpha$ -D-glucosidic linkages, forming a highly branched and rigid

granule structure of about  $10^7$ – $10^8$  Da in mass and 42 nm in size (Ball et al., 2011). The first decisive step of glycogen biosynthesis is the synthesis of the ADP-glucose by ADP-glucose pyrophosphorylase (AGPase, GlgC; for review, see Zilliges, 2014). The glucose moiety of ADP-glucose is transferred to the non-reducing end of a linear  $\alpha$ -(1-4) glucan chain, a reaction catalyzed by glycogen synthase (GlgA). The branching enzyme (GlgB) introduces symmetrically distributed  $\alpha$ -(1-6) glucosidic linkages according to a binary branching principle. In contrast, PHB is a non-water-soluble, conformationally amphiphilic, linear, and highly flexible polyester consisting of (*R*)-3-hydroxybutyrate units, forming 200–500 nm large inclusions (Reusch, 2012; Jendrossek and Pfeiffer, 2014). The first step in PHB synthesis is the condensation of two molecules of acetyl-coenzyme A (CoA) to acetoacetyl-CoA, as catalyzed by  $\beta$ -ketothiolase (PhaA). The subsequent reduction by acetoacetyl-CoA reductase (PhaB) forms the monomeric precursor D-3-hydroxybutyryl-CoA, which is finally polymerized to PHB by PHB synthase (PhaC/PhaE; Hein et al., 1998). The PHB granule is surrounded by a membranous surface layer containing phasin protein, which is involved in granule formation and granule attachment to cellular components (Hauf et al., 2015).

Most microorganisms synthesize either glycogen or PHB. The enzymatic settings for the metabolism of glycogen, the prevalent cyanobacterial carbon polymer, are conserved in all cyanobacteria (Beck et al., 2012). A few cyanobacteria such as *Synechocystis* sp. PCC 6803 additionally synthesize PHB (Allen, 1984; Stal, 1992; Beck et al., 2012). This dual capability is also present in some symbiotic rhizobia and some phototrophic purple bacteria, and it is an exception to the principle that only one kind of carbon-polymer metabolism acts as both a sink and an accessible reserve (De Philippis et al., 1992). In recent years, a deeper insight into the physiological role of the two physico-chemically different carbon polymers has been gained by mutagenesis approaches targeting ADP-glucose pyrophosphorylase (GlgC) and/or glycogen synthase (GlgA, for the block of glycogen synthesis; for overview, see Zilliges, 2014), and targeting ketothiolase (PhaA), and/or PHB synthase (PhaE/C), or associated granule-formation polypeptides (for the block of PHB synthesis; Taroncher-Oldenburg and Stephanopoulos, 2000; Tsang et al., 2013; van der Woude et al., 2014; Hauf et al., 2015). A variety of metabolic principles in cyanobacteria are linked to glycogen metabolism: the maintenance of photosynthetic efficiency in light and of viability in periods of starvation, such as in darkness and/or macronutrient depletion, and the acclimation to macronutrient deficiency (for overview, see Zilliges, 2014). In contrast, the role of PHB synthesis is still unclear due to the lack of a significant phenotype. The primary focus of the current study is the direct comparison of the physiological and metabolic properties of single-knockout mutants ( $\Delta$ glgC;  $\Delta$ phaC) with a very sensitive double-knockout mutant ( $\Delta$ glgC/ $\Delta$ phaC, generated in a phototrophic model organism for the first time), followed by a conclusive discussion of the individual and mutual function of both glycogen and PHB in carbon-flux regulation and in

acclimation to macronutrient stress, especially during nitrogen chlorosis.

## MATERIALS AND METHODS

### Mutagenesis

#### $\Delta$ phaC Mutant

For deletion–insertion mutagenesis, two equipartial, distant sequence stretches of the *phaC* ORF (locus *slr1830*) were fused by an overlap-extension PCR (primer sequences are shown in Supplementary Table S1) and ligated into the pIC20H vector (Marsh et al., 1984). A kanamycin resistance cassette from pUC4K (GE Healthcare) was finally inserted via the *Bam*HI site. The constructs were used to transform *Synechocystis* wild type as described by Ermakova et al. (1993). Transformants were restreaked six times with successively increasing antibiotic pressure. Segregation status was checked and confirmed by PCR (Supplementary Figure S1; Table S1).

#### $\Delta$ glgC Mutant

The deletion–insertion mutagenesis of the *glgC* ORF (locus *slr1176*) is described by Gruendel et al. (2012).

#### $\Delta$ glgC::*glgC* Complementation

For complementation of the *glgC* knockout, the *glgC* ORF (locus *slr1176*) was amplified in full length (including the promoter-region) by PCR (Supplementary Table S1). The PCR product was integrated via *Sal*I and *Pst*I into the self-replicating plasmid pVZ325. The plasmid DNA was transferred into *Synechocystis* wild-type cells and  $\Delta$ glgC-mutant cells by triparental mating as described by Wilde and Dienst (2011). Exconjugants were restreaked six times with successively increasing antibiotic pressure, analyzed and confirmed by PCR (Supplementary Figure S2).

#### $\Delta$ glgC/ $\Delta$ phaC Mutant

For double-knockout mutagenesis of both glycogen and PHB synthesis, the  $\Delta$ glgC-knockout plasmid (Gruendel et al., 2012) was used for transformation of the  $\Delta$ phaC-mutant cells. Transformants were restreaked six times with successively increasing antibiotic pressure, analyzed and confirmed by PCR (see Supplementary Figure S1).

## Physiological Analysis

### Bacterial Strains and Growth Conditions

All comparative growth experiments were strictly performed without antibiotic pressure. Liquid cultures of *Synechocystis* wild-type and mutant strains were grown at 28°C under continuous illumination with white light in a range of 45–120  $\mu$ mol photons  $m^{-2} s^{-1}$  in BG11 medium containing 20 mM HEPES buffer (pH 8.0) and 17.6 mM sodium nitrate as the nitrogen source (+N) or in BG11<sub>0</sub>, which lacks sodium nitrate (–N; Stanier et al., 1971). The growth experiments were implemented in Erlenmeyer flasks on a rotary shaker or were performed in elongated, slim flasks, which were continuously aerated in a range of 0.03–3% v/v CO<sub>2</sub>. Nitrogen deprivation was achieved by a

repeated washing/centrifugation procedure and resuspension in BG11<sub>0</sub> medium as described by Gruendel et al. (2012). The cell suspensions obtained were split, and nitrate was added to the control cell culture. Additionally, 10 mM sodium acetate (with a final concentration of 0.4% w/v) was added to the cultures in order to induce a significant PHB accumulation (after 7 days) as described by Hein et al. (1998).

Growth characteristics of the strains (in liquid cultures) have been recorded by both daily measurement of the optical density at 750 nm (OD<sub>750</sub>) and of the chlorophyll a content (determination according to the method as described by Tandeau de Marsac and Houmard, 1988), recorded by Specord200 plus spectrophotometer (Analytic Jena AG, Germany) (Supplementary Figure S3).

To monitor the bleaching process (under nitrogen deprivation), whole-cell absorption spectra have been recorded in the wavelength range from 400 to 750 nm on a Specord200 plus spectrophotometer (Analytic Jena AG, Germany) equipped with a scattering position and were normalized at 750 nm (Supplementary Figure S4).

### Viability Tests (Spot Assay Experiments)

For spot assay experiments, cells were grown in Erlenmeyer flasks to an OD<sub>750 nm</sub> of 0.4–0.6 and adjusted to a chlorophyll content of 5 μg ml<sup>-1</sup> (chlorophyll determination according to the method as described by Tandeau de Marsac and Houmard (1988)). Seven microliters aliquots of three different dilutions (1:1, 1:10, 1:100) were dropped on BG11 agar plates with (+N) or without NaNO<sub>3</sub> (-N) and with or without K<sub>2</sub>HPO<sub>4</sub> (+P<sub>1</sub> or -P<sub>1</sub>), respectively. The plates were incubated at 28°C under alternating light/dark cycles or continuous illumination with a light intensity of 75 μmol photons m<sup>-2</sup> s<sup>-1</sup> for 7 days (Figures 1–3).


## Determination of Metabolite Concentrations

### Quantitative Determination of Glycogen (Glc<sub>n</sub>)

The intracellular glycogen levels were determined according to the protocol as described by Gruendel et al. (2012) (Figure 4A).

### Quantitative Determination of PHB

Poly-β-hydroxybutyrate was extracted from dried cell pellets and simultaneously hydrolyzed into its monomers (3-hydroxybutyrate) by shaking in 0.5 M NaOH (300–400 μl) (1 h incubation at 85°C). After cooling, the solution was neutralized by the addition of 1 M HCl (volume ratio NaOH/HCl 4:1). Finally, the PHB content was spectrophotometrically quantified in an enzymatic assay using 3-hydroxybutyrate dehydrogenase from *Rhodobacter spheroides* (Roche Diagnostics) in the presence of nicotinamide adenosine dinucleotide (NAD). The NADH synthesis was coupled to a phenazine methosulphate-p-iodonitrotetrazolium violet (PMS-INT) colorimetric assay (to prevent a backward reaction) as described by Lim and Buttery (1977). Here, the generated NADH transfers its hydrogen through the PMS-INT system to produce a red formazan, which is stable and absorbs light at 505 nm. The PHB quantification (Figure 4B) was performed via the calibration of NADH series.



**FIGURE 1 | Viability test under alternating light-dark conditions.** Growth and viability of the wild type (wt) and mutants defective in glycogen synthesis ( $\Delta glgC$ ), PHB synthesis ( $\Delta phaC$ ), or both carbon-polymer syntheses ( $\Delta glgC/\Delta phaC$ ) were tested via spot assays (7 days) under continuous and alternating light/dark (16 h/8 h, 12 h/12 h, or 8 h/16h) conditions (see Materials and Methods). The effect was verified by three independent spot assays.

### Quantitative Determination of Cyanophycin (CyPh) Content

Cyanophycin was extracted as described by Li et al. (2001) using a french press instead of a sonifier. The cyanophycin pellet was hydrolyzed as described by Maheswaran et al. (2006) and quantified enzymatically as described by Bergmeyer et al. (1974) (Figure 4C).

### Quantitative Determination of Extracellular Pyruvate and 2-Oxoglutarate


The extracellular pyruvate and 2-oxoglutarate levels were determined according to the protocol as described by Gruendel et al. (2012) (Table 1).

### Quantitative Determination of Dry Weight

For the determination of dry weight, cell pellets of 10–50 ml culture were washed twice with sterile water, dried completely at 80°C in a heating device, and finally weighed.

## Transmission Electron Microscopy

Samples for Transmission electron microscopy (TEM) were prepared as described previously for *Anabaena* sp. PCC 7120 (Fiedler et al., 1998). Fixation and post-fixation of the cells were performed using 2.5% (v/v) glutaraldehyde and 2% (w/v)



**FIGURE 2 | Viability test under nitrogen and phosphate depletion.**


Growth and viability of the wild type (wt) and mutants defective in glycogen synthesis ( $\Delta glgC$ ), PHB synthesis ( $\Delta phaC$ ), or both carbon-polymer syntheses ( $\Delta glgC/\Delta phaC$ ) were tested via spot assays under continuous illumination (7 days) on BG11 agar plates lacking the respective nitrogen (-N) or phosphate (-P) source (see Materials and Methods). The effect was verified by four independent spot assays.

potassium permanganate. Fixed cells were immobilized in agarose and dehydrated by a series of increasing ethanol. After embedding in EPON and sectioning, the ultrathin sections (60–80 nm) were stained with uranyl acetate and lead citrate. The sections were examined with a Philipps Tecnai electron microscope at 80 kV (Figure 5).

## RESULTS

### Double Mutants Lacking Glycogen and PHB

To completely abolish carbon-polymer biosynthesis, knockout mutants of either ADP-glucose pyrophosphorylase (AGPase, GlgC) or glycogen synthase (GlgA) for glycogen biosynthesis, or of either  $\beta$ -ketothiolase (PhaA), Acetoacetyl-CoA reductase (PhaB), or PHB synthase (PhaC/PhaE) for PHB synthesis were successfully created in different cyanobacterial strains previously (Taroncher-Oldenburg and Stephanopoulos, 2000; Xie et al., 2011; Tsang et al., 2013; van der Woude et al., 2014; Zilliges, 2014). In this study, we describe for the first time the generation of a double-knockout mutant lacking both AGPase (GlgC) and PHB synthase (PhaC) (see Supplementary Figure S1). The resulting strain ( $\Delta glgC/\Delta phaC$ ) is not able to synthesize neither glycogen nor PHB (see below). The direct comparison of this mutant with the respective single-knockout mutant and the wild




**FIGURE 3 | Recovery capability.** The recovery capability of nitrogen-depleted (-N, 2 days) wild type (wt) and mutants defective in glycogen synthesis ( $\Delta glgC$ ), PHB synthesis ( $\Delta phaC$ ), or both carbon-polymer syntheses ( $\Delta glgC/\Delta phaC$ ) were tested via spot assays under continuous illumination on BG11 agar plates replenished with the nitrogen source (+N; see Materials and Methods). The effect was verified by six independent spot assays.

type (as well as a complemented variant, see Supplementary Figure S2 for *glgC*) provide a deeper understanding of the individual and the mutual physiological roles as well as the metabolic interrelations of the two carbon-polymer syntheses.

### Energy and Carbon Sources for the Maintenance Metabolism in Darkness

All cyanobacteria are able to perform a maintenance metabolism in the absence of light by predominantly using internal polymer reserves. Various studies showed the contribution of the carbon storage polymer glycogen to the maintenance metabolism in darkness. In contrast, the involvement of PHB metabolism has not been reported so far (for review, see Zilliges, 2014). In order to directly compare and demonstrate putative interrelations between the two carbon polymer syntheses, the growth and viability of a glycogen-deficient mutant ( $\Delta glgC$ ), of a PHB-deficient mutant ( $\Delta phaC$ ), and of a mutant deficient in the synthesis of both compounds ( $\Delta glgC/\Delta phaC$ ) have been tested via spot assays as well as in liquid culture systems under different alternating light/dark regimes (Figure 1 and Supplementary Figures S3A,B). The  $\Delta phaC$  mutant shows similar viability and growth as the wild type under all tested conditions (Figure 1; Supplementary Figures S3A,B). In contrast, the lack of glycogen in the  $\Delta glgC$  strain has a negative effect on viability and growth with increasing length of periodic darkness (Figure 1; Supplementary Figures S3A,B). Only under continuous low-light regimes, the viability and growth of the  $\Delta glgC$  mutant and the wild type are similar (Figure 1 and Supplementary



Figures S3A,B). In contrast, the double-knockout mutant  $\Delta glgC/\Delta phaC$  shows reduced viability even under continuous low-light conditions (Figure 1). Moreover, the negative effect on the viability and growth with increasing length of periodic darkness is more pronounced in the  $\Delta glgC/\Delta phaC$  mutant than in the  $\Delta glgC$  single mutant. Cells deficient in both PHB and glycogen synthesis are not able to grow in conditions of periodic 12 h/12 h light/dark regimes (Figure 1). In conclusion, both carbon-polymer syntheses are presumably essential for the photoautotrophic lifestyle of *Synechocystis* sp. PCC 6803. However, the glycogen pool (and not the PHB pool) might be the main energy and carbon source for the maintenance metabolism in darkness and the PHB pool is dispensable.

### Carbon-Polymer Synthesis as an Adaptation and Survival Strategy to Macronutrient Stress Conditions

Unbalanced growth conditions such as macronutrient starvation generally lead to an accumulation of reserve material as well as to a rapid degradation after replenishing conditions in all microorganisms including cyanobacteria (Schwarz and

Forchhammer, 2005). To understand the role and interrelations between the two carbon polymer metabolisms in such stress responses, the growth and viability of the double-knockout mutant  $\Delta glgC/\Delta phaC$  in comparison to both single-knockout mutants  $\Delta glgC$  and  $\Delta phaC$  have been tested in response to nitrogen and phosphate depletion (Figure 2 and Supplementary Figure S3C) as well as after nitrogen-replenishing conditions (Figure 3). The  $\Delta glgC$  mutant is significantly impaired in both its viability (Figure 2) and its recovery capability (Figure 3) in prolonged starvation condition, as previously reported by Gruendel et al. (2012).

The single, PHB-deficient mutant  $\Delta phaC$  behaves similar to the wild type. Upon nitrogen starvation both strains duplicate once more and enter chlorosis, being visualized by the yellow color due to the loss of pigments (Figure 2; Supplementary Figure S3C). Upon Pi depletion, the strains show reduced growth and only faint yellow-green pigmentation (Figure 2). The  $\Delta phaC$  mutant recovers after starvation stress like the wild type, tested after 7 days of nitrogen depletion (Figure 3).

In line with this finding, the  $\Delta glgC/\Delta phaC$  mutant shows a similar phenotype as the  $\Delta glgC$  mutant, indicating that the impaired growth, viability, and recovery capacity result rather from the block of glycogen synthesis than from the block of PHB synthesis in this double-knockout mutant (Figures 2 and 3). In conclusion, the glycogen synthesis (and not the PHB synthesis) appears to be the key player in *Synechocystis*'s short-term acclimation response toward macronutrient stress conditions, such as nitrogen starvation.

**TABLE 1 | Exclusive spilling of 2-oxoglutaric acid and pyruvic acid in glycogen-deficient mutants.**

	wt	$\Delta phaC$	$\Delta glgC$	$\Delta glgC:: glgC$	$\Delta glgC/\Delta phaC$
2-OG	n.d.	n.d.	0.1–0.2	n.d.	0.2–0.3
Pyr	n.d.	n.d.	0.3–0.6	n.d.	0.3–0.8

The extracellular levels of 2-oxoglutaric acid (2-OG) and pyruvic acid (Pyr; in mM concentrations; n.d. not detectable) were determined in the supernatant of the wild type (wt), of mutants defective in glycogen synthesis ( $\Delta glgC$ ), PHB synthesis ( $\Delta phaC$ ), or both carbon-polymer syntheses ( $\Delta glgC/\Delta phaC$ ), as well as of the  $\Delta glgC$  mutant, complemented with the  $glgC$  gene in trans ( $\Delta glgC:: glgC$ ), after 48 h of nitrogen depletion (see MATERIALS AND METHODS). Data from determination of extracellular pyruvate and 2-oxoglutarate represent variances from 6 different growth experiments á 3 biological replicates.

### Metabolic Consequences of Blocked Carbon-Polymer Synthesis upon Macronutrient-Stress Response

As previously shown, the block of glycogen biosynthesis as in the  $\Delta glgC$ - or  $\Delta glgA_n$ -mutant strains leads to three very characteristic, co-occurring phenotypes as short-term response to nitrogen starvation: (i) the incapability of carbon-excess

redirection into glycogen-granule formation (see also **Figures 4** and **5**), (ii) impaired chlorosis (non-bleaching phenotype; (**Figures 2, 3**; Supplementary Figure S4), and (iii) the spilling of partially oxidized carbonic acids, such as 2-oxoglutarate and pyruvate (**Table 1**; for overview, see Zilliges, 2014). All three characteristics are also observed with a similar magnitude under the tested conditions in the double-knockout mutant  $\Delta glgC/\Delta phaC$  (**Figures 2–5**; Supplementary Figure S4; **Table 1**). This phenotype is fully complemented by *in-trans* expression of the *glgC* gene under the control of its own promoter in the  $\Delta glgC$ -mutant background (data shown for overflow metabolism only; see **Table 1**:  $\Delta glgC::glgC$ ) and is in agreement with results from Carrieri et al. (2012) using a similar complementation approach. In contrast, the  $\Delta phaC$  mutant responds to nitrogen-starvation conditions in a similar way as the wild type, which is in agreement with previous data obtained with a  $\Delta phaA$  mutant (van der Woude et al., 2014). Both wild-type and  $\Delta phaC$  mutant accumulate similar amounts of glycogen (approx. 25% of DW in 48 h) (**Figure 4**), degrade their phycobilisomes in a similar period (approx. 48 h; Supplementary Figure S4), and do not spill any of the partially oxidized carbonic acids such as 2-oxoglutarate (**Table 1**). Under the nitrogen-replenishing conditions tested, the wild type and the  $\Delta phaC$  mutant show a similar glycogen-degradation pattern (data not shown) and a similar cyanophycin accumulation (**Figure 4**).


In contrast, the glycogen-deficient mutants  $\Delta glgC$  and  $\Delta glgC/\Delta phaC$  are apparently not able to redirect the replenished

nitrogen sources into cyanophycin biosynthesis, which is not detectable in cell extracts under the tested conditions (**Figure 4**). However, in response to nitrogen-starvation conditions, the single-knockout mutant  $\Delta glgC$  significantly (**Figure 4**: *p*-value below 0.01) redirect more of the carbon excess into PHB biosynthesis than the respective wild type, which is in agreement with data from Wu et al. (2001).

In conclusion, blocks in glycogen synthesis (but not blocks in PHB synthesis) lead to overflow reactions into other polymer syntheses (and apparently not *vice versa*) as well as to a spilling of certain metabolites such as 2-oxoglutarate in response to macronutrient starvation.

### Effect on Cellular Ultra-Structure by Lack of Carbon-Polymer Biosyntheses

Four major structural and morphological changes are observed in wild-type cells in response to nitrogen starvation (**Figure 5**): (i) a massive accumulation of electron-dense glycogen inclusions (approx. 40 nm in diameter) between the thylakoid layers (**Figure 5E**), (ii) the degradation of the antenna complexes (phycobilisomes; Supplementary Figure S4), (iii) the disassembling of the thylakoid membrane layers including a reduction by number and packing density (**Figure 5E**), and (iv) the formation of distinct electron-transparent PHB granules (approx. 400–500 nm in diameter; **Figure 5E**). Similar structural and morphological changes, except for the accumulation of PHB



**FIGURE 5 | Ultra-structural changes in response to nitrogen starvation.** Structural changes occurring in wild type (wt) (**A, E**) and mutant strains defective in glycogen synthesis ( $\Delta glgC$ ) (**B, F**), in PHB synthesis ( $\Delta phaC$ ) (**C, G**), or in both carbon-polymer syntheses ( $\Delta glgC/\Delta phaC$ ) (**D, H**) in response to nitrogen deprivation were monitored by transmission electron microscopy. Cells deprived of nitrogen (2 days) (**E–H**) are compared with respective, untreated controls (**A–D**) (see Materials and Methods). T, thylakoids; GG, glycogen granule; PHB, polyhydroxybutyrate granule; C, carboxysome; PP, polyphosphate granule; V, vesicle; PG, peptidoglycan layer; CM, cytoplasmic membrane; OM, outer membrane.

inclusions, occur in the PHB synthase-deficient mutant  $\Delta phaC$  (Figures 5C,G and Supplementary Figure S4). This finding is in agreement with previously published data by Tsang et al. (2013). Some of the electron-transparent structures in both cell types are presumably polyphosphate granules (indicated in Figure 5A) as previously highlighted by Tsang et al. (2013) and Nagarajan et al. (2014).

In contrast, AGPase-deficient mutant cells ( $\Delta glgC$ ; Figures 5B,F; Supplementary Figure S4) as well as glycogen synthase-deficient cells ( $\Delta glgA1/\Delta glgA2$ ; data not shown) are not able neither to degrade their antenna complexes (phycobilisomes), nor to reduce the number and packing density of the thylakoid membrane-layer system, nor to accumulate glycogen granules in response to nitrogen depletion. Instead, several undefined vesicles (approx. 90 nm in diameter) appear inside the thylakoid stacks (Figures 5B,F). Preparative artifacts could be excluded due to equal treatment of all samples (Figure 5; compare to wild type and  $\Delta phaC$  mutant). A similar effect of structural and morphological changes as in  $\Delta glgC$ -mutant cells, except for the formation of definitive PHB granules, has been observed in the double-knockout mutant  $\Delta glgC/\Delta phaC$  (Figures 5D,H).

In conclusion, the glycogen synthesis (and not the PHB synthesis) has a substantial influence on the adaptation response to macronutrient starvation. The formation of glycogen granules appears as an important key player in the rearrangement and disassembling of the photosynthetic machinery such as phycobilisome degradation and thylakoid density and packing alterations. These changes are necessary to acclimate and trophically convert from an active photosynthetic cell to a dormant status.


## DISCUSSION

### Glycogen Synthesis as a Metabolic and Structural Key Process in Nitrogen Acclimation

The nitrogen-starvation response is characterized by a cascade of structural and metabolic events that finally switch the cellular mode from photosynthetically active to dormant status in non-diazotrophic cyanobacteria. These substantial changes are tightly related to active glycogen synthesis. The removal of external nitrate or ammonia sources stops the primary nitrogen assimilation processes mainly mediated by glutamine synthetase and glutamine/2-oxoglutarate aminotransferase. The reduction of cellular nitrogen is accompanied by the rapid rise of 2-oxoglutarate (2-OG) levels due to the absence of 2-OG dehydrogenase (2-OGDH; Muro-Pastor et al., 2001). Even in 2-OGDH-containing bacteria and plants, the enzyme is almost always inhibited during nitrogen starvation, which leads in turn to a concomitant overflow of the carbon excess into either glycogen or PHB polymer synthesis, amino acid biosynthesis, or alternative TCA shunts (e.g., GABA shunt; Walshaw et al., 1997). Our comparative study of knockout mutants reveals unambiguously that the primary carbon excess (from terminating

photosynthetic reactions and starting amino acid release by protein degradation (Hasunuma et al., 2013; Depraetere et al., 2015) is mainly directed into glycogen synthesis and not into PHB synthesis (Figure 4). Only blocks of glycogen synthesis in *Synechocystis* sp. PCC 6803 lead to a prompt spilling of partially oxidized substrates such as 2-OG (to get rid of carbon excess) under unbalanced growth conditions, independent of the absence or presence of an active PHB synthesis (Table 1). However, this spilling effect is similar to PHB-synthesis knockouts in other bacteria naturally lacking glycogen metabolism (Steinbuechel and Schlegel, 1989; Russell, 2007). The directing of carbon excess into glycogen synthesis (and not into PHB synthesis) is crucial for an adequate acclimation from an active photosynthetic cell to a dormant cell in the presence of light. In the absence of external nitrogen sources, both light-dependent processes, the progression of chlorosis (Gorl et al., 1998) and the glycogen synthesis (Gruendel et al., 2012), are inextricably linked with each other (Figure 6; Supplementary Figure S4). Statistical data analysis indicates a significant linear correlation (coefficient of determination  $R^2$  is 0.74; Pearson coefficient is 0.86;  $p$ -value is  $1.52 \times 10^{-25}$ ) between the progression of both glycogen accumulation and phycobilisome degradation in wild-type cells (Figure 6; Supplementary Figure S4) and in  $\Delta phaC$  cells (data not shown) as a short-term response to nitrogen depletion. Cells deficient in glycogen synthesis are not able to perform the switch and remain in a metabolic arrest (Figures 5B,F).

We hypothesize two mechanisms that do not necessarily exclude each other: a metabolic mechanism (i) and/or a structural mechanism (ii). (i) The concomitant directing of carbon excess (mainly from protein degradation) into glycogen synthesis allows the adjustment to an adequate cellular 2-OG concentration under conditions of nitrogen depletion, which in turn leads to an activation of the nitrogen signaling cascade, in particular of the PII signaling protein (Muro-Pastor et al., 2001; Schwarz and Forchhammer, 2005). An insufficient activation of enzymes and associated proteins of the nitrogen signaling cascade most likely prevents nitrogen chlorosis, similar as shown for *nblA* in *Synechococcus elongatus* PCC 7942 by Hickman et al. (2013). (ii) The replacement of phycobilisomes (around 35–40 nm in size) by rigid glycogen granules (around 42 nm in size) between the thylakoid layers is a crucial structural component for the subsequent, controlled shut-down process of photosynthetic activity including thylakoid disassembly. Likewise, the AGPase-deficient mutant ( $\Delta glgC$ ) of *Synechococcus* sp. PCC 7002 still maintained an efficient energy transfer from phycobilisomes to photosystem II reaction centers after prolonged nitrogen starvation (Jackson et al., 2015). The increased abundance of yet undefined vesicles between the thylakoid layers in the  $\Delta glgC$  mutant of *Synechocystis* sp. PCC 6803 (Figure 5B) additionally supports the new role of glycogen synthesis/granules for the integrity of the thylakoid layer system. Such vesicles (around 90 nm in size) might represent spilling products (to remove destabilizing catabolites from the thylakoid membrane) or plastoglobuli-like structures, which have similar functions in the thylakoid biogenesis/degradation processes in plants (Brehelin and Kessler, 2008; Cunningham et al., 2010; Besagni and Kessler, 2013). A potential confusion with lipid droplets is not likely due



to size and shape of the vesicles (Peramuna and Summers, 2014; Perez et al., 2015).

## The Classical Storage Role and Potential Interrelations of Both Carbon Polymers

This comparative study shows that the glycogen and PHB biosyntheses do not negatively affect each other (Figure 4) and thus potentially do not interrelate under conditions of nitrogen starvation. Only a small proportion of primary carbon excess is redirected into PHB synthesis in cells blocked in glycogen synthesis (Figure 4). Most excessive carbon proportions are promptly spilled (Table 1). In this respect, it is not clarified whether parts of the carbon excess were directed into lipid biosynthesis, another abundant form of carbon storage in *Synechocystis* sp. PCC 6803 (Monshupanee and Incharoensakdi, 2014). Lipid droplets as observed in akinetes of *Nostocales* were not detected in electron micrographs of wild type and mutant cells (Figure 5; compare Perez et al., 2015). Moreover, this and several other physiological studies show that massive glycogen synthesis starts promptly with the onset of nitrogen starvation. Significant amounts of glycogen are already detectable after one day, peaking around 3 days of nitrogen starvation, whereas PHB does not peak until after five days of the nitrogen-starvation conditions tested (Lehmann and Wober, 1976; De Philippis et al., 1992; Miyake et al., 1997; Panda et al., 2006; Monshupanee and Incharoensakdi, 2014; Hauf et al., 2015). This temporal shift in the nitrogen response of the two carbon-polymer syntheses is in agreement with transcriptional data (Krasikov et al., 2012; Huang et al., 2013; Kopf et al., 2014; Depraetere et al., 2015). A concomitant expression has only been shown for enzymes involved in glycogen degradation, such as the glycogen phosphorylase (GlgP2) and debranching enzyme (GlgX), as well as for PHB synthesis enzymes, such as

ketothiolase (PhaA), acetyl-coenzyme A reductase (PhaB), and PHB synthase (PhaCE; Azuma et al., 2011; Osanai et al., 2013; Nakaya et al., 2015). Both catabolic enzymes (GlgP2, GlgX) are potentially also involved in glycogen modification, which changes the accessibility of the polymer for short- or long-term storage (Suzuki et al., 2007). Finally, we cannot conclusively answer whether the carbon polymers glycogen and PHB serve as storage compounds for *Synechocystis*' maintenance metabolism during macronutrient-starvation periods.

The recovery incapability of the glycogen-deficient mutants (Figure 3) possibly results from the variety of negative pleiotropic (metabolic and structural) effects, summarized in an impaired acclimation response, than from the lack of carbon source. Solely PHB-deficient cells are neither affected under the conditions tested (Figures 1–3), which is in agreement with Tsang et al. (2013), van der Woude et al. (2014), and Hauf et al. (2015). A longer starvation exposure (above seven days), the determination of polymer modifications under nitrogen starvation and replenished conditions will most certainly bring new indications regarding the physiological function. Moreover, it cannot be conclusively answered why a few cyanobacterial species such as *Synechocystis* sp. PCC 6803 (and some purple bacteria) massively synthesize both types of the physically and chemically different carbon polymers, glycogen, and PHB, in response to macronutrient deficiency (Figure 4). Obviously, PHB synthesis does not act as a backup of glycogen synthesis and does not compensate knockouts in glycogen metabolism.

These observations are in line with the general metabolic principle that only one kind of carbon-polymer metabolism acts as both a sink and an accessible reserve. So far, the only indication for the specific role of PHB in *Synechocystis* sp. PCC 6803 is given by the pronounced sensitivity and reduced growth rate of the double-knockout mutant  $\Delta glgC/\Delta phaC$  (certain loss of robustness) compared to the single knockouts even



under standard growth conditions (Figure 1). In this respect, some already discussed functions of PHB are conceivable: (i) as a transient electron sink (Stal, 1992) and/or (ii) as storage component (however questioned by the lack of specific enzymes for PHB degradation in *Synechocystis* sp. PCC 6803 and the non-compensatory effect in glycogen-deficient knockouts), and/or (iii) as a structural component during the equal separation of the multiple bacterial nucleoids and PHB granules in cell division processes (Jendrossek and Pfeiffer, 2014), that might be essential for an immediate re-start of cell proliferation after replenished nitrogen conditions.

## AUTHOR CONTRIBUTIONS

RD, IM, and YZ made substantial contributions to the conception and the design of the study, the acquisition of data, and the analysis and interpretation of data; RD, IM, and YZ participated in drafting the article and revising it critically for important intellectual content; RD, IM, and YZ gave final approval of the version to be submitted. RD, IM, and YZ agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

## REFERENCES

- Allen, M. M. (1984). Cyanobacterial cell inclusions. *Annu. Rev. Microbiol.* 38, 1–25. doi: 10.1146/annurev.mi.38.100184.000245
- Azuma, M., Osanai, T., Hirai, M. Y., and Tanaka, K. (2011). A response regulator Rre37 and an RNA polymerase sigma factor SigE represent two parallel pathways to activate sugar catabolism in a cyanobacterium *Synechocystis* sp PCC 6803. *Plant Cell Physiol.* 52, 404–412. doi: 10.1093/pcp/pcq204
- Ball, S., Colleoni, C., Cenci, U., Raj, J. N., and Tirtiaux, C. (2011). The evolution of glycogen and starch metabolism in eukaryotes gives molecular clues to understand the establishment of plastid endosymbiosis. *J. Exp. Bot.* 62, 1775–1801. doi: 10.1093/jxb/erq411
- Beck, C., Knoop, H., Axmann, I., and Steuer, R. (2012). The diversity of cyanobacterial metabolism: genome analysis of multiple phototrophic microorganisms. *BMC Genomics* 13:56. doi: 10.1186/1471-2164-13-56
- Bergmeyer, H. U., Bernt, E., Möllering, H., and Pfeleiderer, G. (1974). “L-Aspartate and L-Asparagine,” in *Methods of Enzymatic Analysis (Second English Edition)*, eds H. U. Bergmeyer and K. Gawehn (Cambridge: Academic Press), 1696–1700.
- Besagni, C., and Kessler, F. (2013). A mechanism implicating plastoglobules in thylakoid disassembly during senescence and nitrogen starvation. *Planta* 237, 463–470. doi: 10.1007/s00425-012-1813-9
- Brehelin, C., and Kessler, F. (2008). The plastoglobule: a bag full of lipid biochemistry tricks. *Photochem. Photobiol.* 84, 1388–1394. doi: 10.1111/j.1751-1097.2008.00459.x
- Carriero, D., Paddock, T., Maness, P. C., Seibert, M., and Yu, J. P. (2012). Photocatalytic conversion of carbon dioxide to organic acids by a recombinant cyanobacterium incapable of glycogen storage. *Energy Environ. Sci.* 5, 9457–9461. doi: 10.1039/c2ee23181f
- Cunningham, F. X. Jr., Tice, A. B., Pham, C., and Gantt, E. (2010). Inactivation of genes encoding plastoglobulin-like proteins in *Synechocystis* sp PCC 6803 leads to a light-sensitive phenotype. *J. Bacteriol.* 192, 1700–1709. doi: 10.1128/JB.01434-09
- De Philippis, R., Ena, A., Guastiini, M., Sili, C., and Vincenzini, M. (1992). Factors affecting poly- $\beta$ -hydroxybutyrate accumulation in cyanobacteria and in purple non-sulfur bacteria. *FEMS Microbiol. Lett.* 103, 187–194. doi: 10.1016/0378-1097(92)90309-C

## FUNDING

We thank the Deutsche Forschungsgemeinschaft (DFG) for financial support (to YZ) in the framework of the Collaborative Research Center 1078 on ‘Protonation Dynamics in Protein Function’ (SFB 1078, project A4/Holger Dau).

## ACKNOWLEDGMENTS

We particularly acknowledge Wolfgang Lockau for his commitment and valuable discussions on this topic and for his sustained support of our research. We gratefully thank Gisa Baumert (Humboldt-Universität zu Berlin, Germany) and Claudia Menzel (Eberhard-Karls-Universität Tübingen, Germany) for their excellent technical assistance.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2016.00966>

- Depraetere, O., Deschoenmaeker, F., Badri, H., Monsieurs, P., Foubert, I., Leys, N., et al. (2015). Trade-Off between growth and carbohydrate accumulation in nutrient-limited *Arthrospira* sp PCC 8005 studied by integrating transcriptomic and proteomic approaches. *PLoS ONE* 10:0132461. doi: 10.1371/journal.pone.0132461
- Ermakova, S. Y., Elanskaya, I. V., Kallies, K. U., Weihe, A., Borner, T., and Shestakov, S. V. (1993). Cloning and sequencing of mutant psbB genes of the cyanobacterium *Synechocystis* PCC 6803. *Photosynthesis Res.* 37, 139–146. doi: 10.1007/BF02187472
- Fiedler, G., Arnold, M., Hannus, S., and Maldener, I. (1998). The DevBCA exporter is essential for envelope formation in heterocysts of the cyanobacterium *Anabaena* sp. strain PCC 7120. *Mol. Microbiol.* 27, 1193–1202. doi: 10.1046/j.1365-2958.1998.00762.x
- Gorl, M., Sauer, J., Baier, T., and Forchhammer, K. (1998). Nitrogen-starvation-induced chlorosis in *Synechococcus* PCC 7942: adaptation to long-term survival. *Microbiology* 144, 2449–2458. doi: 10.1099/00221287-144-9-2449
- Gruendel, M., Scheunemann, R., Lockau, W., and Zilliges, Y. (2012). Impaired glycogen synthesis causes metabolic overflow reactions and affects stress responses in the cyanobacterium *Synechocystis* sp PCC 6803. *Microbiology* 158, 3032–3043. doi: 10.1099/mic.0.062950-0
- Hasunuma, T., Kikuyama, F., Matsuda, M., Aikawa, S., Izumi, Y., and Kondo, A. (2013). Dynamic metabolic profiling of cyanobacterial glycogen biosynthesis under conditions of nitrate depletion. *J. Exp. Bot.* 64, 2943–2954. doi: 10.1093/jxb/ert134
- Hauf, W., Watzel, B., Roos, N., Klotz, A., and Forchhammer, K. (2015). Photoautotrophic polyhydroxybutyrate granule formation is regulated by cyanobacterial Phasin PhaP in *Synechocystis* sp strain PCC 6803. *Appl. Environ. Microbiol.* 81, 4411–4422. doi: 10.1128/AEM.00604-15
- Hein, S., Tran, H., and Steinbuechel, A. (1998). *Synechocystis* sp. PCC6803 possesses a two-component polyhydroxyalkanoic acid synthase similar to that of anoxygenic purple sulfur bacteria. *Arch. Microbiol.* 170, 162–170. doi: 10.1007/s002030050629
- Hickman, J. W., Kotovic, K. M., Miller, C., Warrenner, P., Kaiser, B., Jurista, T., et al. (2013). Glycogen synthesis is a required component of the nitrogen stress response in *Synechococcus elongatus* PCC 7942. *Algal Res.* 2, 98–106. doi: 10.1016/j.algal.2013.01.008

- Huang, S., Chen, L., Te, R., Qiao, J., Wang, J., and Zhang, W. (2013). Complementary iTRAQ proteomics and RNA-seq transcriptomics reveal multiple levels of regulation in response to nitrogen starvation in *Synechocystis* sp PCC 6803. *Mol. Biosyst.* 9, 2565–2574. doi: 10.1039/c3mb70188c
- Jackson, S. A., Eaton-Rye, J. J., Bryant, D. A., Posewitz, M. C., and Davies, F. K. (2015). dynamics of photosynthesis in a glycogen-deficient *glgC* Mutant of *Synechococcus* sp. Strain PCC 7002. *Appl. Environ. Microbiol.* 81, 6210–6222. doi: 10.1128/AEM.01751-15
- Jendrossek, D., and Pfeiffer, D. (2014). New insights in the formation of polyhydroxyalkanoate granules (carbonosomes) and novel functions of poly(3-hydroxybutyrate). *Environ. Microbiol.* 16, 2357–2373. doi: 10.1111/1462-2920.12356
- Kopf, M., Klaehn, S., Scholz, I., Matthiessen, J. K. F., Hess, W. R., and Voss, B. (2014). Comparative analysis of the primary transcriptome of *Synechocystis* sp PCC 6803. *DNA Res.* 21, 527–539. doi: 10.1093/dnares/dsu018
- Krasikov, V., von Wobeser, E. A., Dekker, H. L., Huisman, J., and Matthijs, H. C. P. (2012). Time-series resolution of gradual nitrogen starvation and its impact on photosynthesis in the cyanobacterium *Synechocystis* PCC 6803. *Physiol. Plant* 145, 426–439. doi: 10.1111/j.1399-3054.2012.01585.x
- Lehmann, M., and Woher, G. (1976). Accumulation, mobilization and turn-over of glycogen in blue-green bacterium *Anacystis nidulans*. *Arch. Microbiol.* 111, 93–97. doi: 10.1007/BF00446554
- Li, H., Sherman, D. M., Bao, S. L., and Sherman, L. A. (2001). Pattern of cyanophycin accumulation in nitrogen-fixing and non-nitrogen-fixing cyanobacteria. *Arch. Microbiol.* 176, 9–18. doi: 10.1007/s002030100281
- Lim, H. H., and Buttery, J. E. (1977). Determination of ethanol in serum by an enzymatic PMS-INT colorimetric method. *Clin. Chim. Acta* 75, 9–12. doi: 10.1016/0009-8981(77)90494-6
- Maheswaran, M., Ziegler, K., Lockau, W., Hagemann, M., and Forchhammer, K. (2006). P-II-regulated arginine synthesis controls accumulation of cyanophycin in *Synechocystis* sp strain PCC 6803. *J. Bacteriol.* 188, 2730–2734. doi: 10.1128/JB.188.7.2730-2734.2006
- Marsh, J. L., Erfle, M., and Wykes, E. J. (1984). The pIC plasmid and phage vectors with versatile cloning sites for recombinant selection by insertional inactivation. *Gene* 32, 481–485. doi: 10.1016/0378-1119(84)90022-2
- Miyake, M., Kataoka, K., Shirai, M., and Asada, Y. (1997). Control of poly-beta-hydroxybutyrate synthase mediated by acetyl phosphate in cyanobacteria. *J. Bacteriol.* 179, 5009–5013.
- Monshupanee, T., and Incharoensakdi, A. (2014). Enhanced accumulation of glycogen, lipids and polyhydroxybutyrate under optimal nutrients and light intensities in the cyanobacterium *Synechocystis* sp. PCC 6803. *J. Appl. Microbiol.* 116, 830–838. doi: 10.1111/jam.12409
- Muro-Pastor, M. I., Reyes, J. C., and Florencio, F. J. (2001). Cyanobacteria perceive nitrogen status by sensing intracellular 2-oxoglutarate levels. *J. Biol. Chem.* 276, 38320–38328.
- Nagarajan, S., Srivastava, S., and Sherman, L. A. (2014). Essential role of the plasmid hik31 operon in regulating central metabolism in the dark in *Synechocystis* sp. PCC 6803. *Mol. Microbiol.* 91, 79–97. doi: 10.1111/mmi.12442
- Nakaya, Y., Iijima, H., Takanobu, J., Watanabe, A., Hirai, M. Y., and Osanai, T. (2015). One day of nitrogen starvation reveals the effect of sigE and rre37 overexpression on the expression of genes related to carbon and nitrogen metabolism in *Synechocystis* sp. PCC 6803. *J. Biosci. Bioeng.* 120, 128–134. doi: 10.1016/j.jbiosc.2014.12.020
- Osanai, T., Numata, K., Oikawa, A., Kuwahara, A., Iijima, H., Doi, Y., et al. (2013). Increased bioplastic production with an rna polymerase sigma factor SigE during nitrogen starvation in *Synechocystis* sp PCC 6803. *DNA Res.* 20, 525–535. doi: 10.1093/dnares/dst028
- Panda, B., Jain, P., Sharma, L., and Mallick, N. (2006). Optimization of cultural and nutritional conditions for accumulation of poly-beta-hydroxybutyrate in *Synechocystis* sp PCC 6803. *Biores. Technol.* 97, 1296–1301. doi: 10.1016/j.biortech.2005.05.013
- Peramun, A., and Summers, M. L. (2014). Composition and occurrence of lipid droplets in the cyanobacterium *Nostoc punctiforme*. *Arch. Microbiol.* 196, 881–890. doi: 10.1007/s00203-014-1027-6
- Perez, R., Forchhammer, K., Salerno, G., and Maldener, I. (2015). Clear differences in metabolic and morphological adaptations of akinetes of two Nostocales living in different habitats. *Microbiology* 162, 214–223. doi: 10.1099/mic.0.000230
- Reusch, R. N. (2012). Physiological importance of Poly-(R)-3-hydroxybutyrate. *Chem. Biodivers.* 9, 2343–2366. doi: 10.1002/cbdv.201200278
- Russell, J. B. (2007). The energy spilling reactions of bacteria and other organisms. *J. Mol. Microbiol. Biotechnol.* 13, 1–11. doi: 10.1159/000103591
- Schwarz, R., and Forchhammer, K. (2005). Acclimation of unicellular cyanobacteria to macronutrient deficiency: emergence of a complex network of cellular responses. *Microbiology* 151, 2503–2514. doi: 10.1099/mic.0.27883-0
- Stal, L. J. (1992). Poly(hydroxyalkanoate) in cyanobacteria: an overview. *FEMS Microbiol. Lett.* 103, 169–180. doi: 10.1111/j.1574-6968.1992.tb05835.x
- Stanier, R. Y., Kunisawa, R., Mandel, M., and Cohen-Bazire, G. (1971). Purification and properties of unicellular blue-green algae (order Chroococcales). *Bacteriol. Rev.* 35, 171–205.
- Steinbuechel, A., and Schlegel, H. G. (1989). Excretion of pyruvate by mutants of *Alcaligenes eutrophus*, which are impaired in the accumulation of the Poly(beta-hydroxybutyric acid) (PHB), under conditions permitting synthesis of PHB. *Appl. Microbiol. Biotechnol.* 31, 168–175. doi: 10.1007/BF00262457
- Suzuki, E., Umeda, K., Nihei, S., Moriya, K., Ohkawa, H., Fujiwara, S., et al. (2007). Role of the GlgX protein in glycogen metabolism of the cyanobacterium, *Synechococcus elongatus* PCC 7942. *Biochim. Biophys. Acta* 1770, 763–773. doi: 10.1016/j.bbagen.2007.01.006
- Tandeu de Marsac, N., and Houmard, J. (1988). Complementary chromatic adaptation: physiological conditions and action spectra. *Methods Enzymol.* 167, 318–328.
- Taroncher-Oldenburg, G., and Stephanopoulos, G. (2000). Targeted, PCR-based gene disruption in cyanobacteria: inactivation of the polyhydroxyalkanoic acid synthase genes in *Synechocystis* sp PCC 6803. *Appl. Microbiol. Biotechnol.* 54, 677–680. doi: 10.1007/s002530000450
- Tsang, T. K., Roberson, R. W., and Vermaas, W. F. J. (2013). Polyhydroxybutyrate particles in *Synechocystis* sp PCC 6803: facts and fiction. *Photosynth. Res.* 118, 37–49. doi: 10.1007/s11120-013-9923-1
- van der Woude, A. D., Angermayr, S. A., Veetil, V. P., Osnato, A., and Hellingwerf, K. J. (2014). Carbon sink removal: increased photosynthetic production of lactic acid by *Synechocystis* sp PCC 6803 in a glycogen storage mutant. *J. Biotechnol.* 184, 100–102. doi: 10.1016/j.jbiotec.2014.04.029
- Walshaw, D. L., Wilkinson, A., Mundy, M., Smith, M., and Poole, P. S. (1997). Regulation of the TCA cycle and the general amino acid permease by overflow metabolism in *Rhizobium leguminosarum*. *Microbiology* 143, 2209–2221. doi: 10.1099/00221287-143-7-2209
- Wilde, A., and Dienst, D. (2011). “Tools for genetic manipulation of cyanobacteria,” in *Bioenergetic Processes of Cyanobacteria*, eds G. A. Peschek, C. Obinger, and G. Renger (Amsterdam: Springer), 685–703.
- Wu, G. F., Shen, Z. Y., Wu, Q. Y., and Zhao, N. M. (2001). Molecular cloning and construction of *agg* gene deletion-mutant in cyanobacterium *Synechocystis* sp. PCC 6803. *Acta Bot. Sin.* 43, 512–516.
- Xie, J., Zhou, J., Zhang, H., and Li, Y. (2011). Increasing reductant NADPH content via metabolic engineering of PHB synthesis pathway in *Synechocystis* sp. PCC 6803. *Chin. J. Biotechnol.* 27, 998–1004.
- Zilliges, Y. (2014). “Glycogen, a dynamic cellular sink and reservoir for carbon,” in *The Cell Biology of Cyanobacteria*, eds E. Flores and A. Herrero (Norfolk, VA: Caister Academic Press), 189–210.

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Damrow, Maldener and Zilliges. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.