



Syddansk Universitet

Ultrasonography in trauma

Weile, Jesper; Nielsen, Klaus; Primdahl, Stine C; Frederiksen, Christian A; Laursen, Christian Borbjerg; Sloth, Erik ; Kirkegaard, Hans

Published in:
Critical Ultrasound Journal

DOI:
[10.1186/s13089-017-0071-2](https://doi.org/10.1186/s13089-017-0071-2)

Publication date:
2017

Document version
Publisher's PDF, also known as Version of record

Document license
CC BY

Citation for pulished version (APA):
Weile, J., Nielsen, K., Primdahl, S. C., Frederiksen, C. A., Laursen, C. B., Sloth, E., & Kirkegaard, H. (2017). Ultrasonography in trauma: a nation-wide cross-sectional investigation. *Critical Ultrasound Journal*, 9, [16]. DOI: [10.1186/s13089-017-0071-2](https://doi.org/10.1186/s13089-017-0071-2)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ORIGINAL ARTICLE

Open Access



Ultrasonography in trauma: a nation-wide cross-sectional investigation

Jesper Weile^{1,2*} , Klaus Nielsen³, Stine C. Primdahl⁴, Christian A. Frederiksen⁵, Christian B. Laursen⁶, Erik Sloth⁷ and Hans Kirkegaard⁸

Abstract

Background: The Focused Assessment with Sonography in Trauma (FAST) protocol is considered beneficial in emergent evaluation of trauma patients with blunt or penetrating injury and has become integrated into the Advanced Trauma Life Support (ATLS) protocol. No guidelines exist as to the use of ultrasonography in trauma in Denmark. We aimed to determine the current use of ultrasonography for assessing trauma patients in Denmark.

Methods: We conducted a nation-wide cross-sectional investigation of ultrasonography usage in trauma care. The first phase consisted of an Internet-based investigation of existing guidelines, and the second phase was a series of structured interviews of orthopedic surgeons, anesthesiologists, and radiologists on call in all hospitals receiving traumatized patients in Denmark.

Results: Guidelines were obtained from all 22 hospitals receiving traumatized patients in Denmark. Twenty-one (95.5%) of the guidelines included and recommended FAST as part of trauma assessment. The recommended person to perform the examination was the radiologist in $n = 11$ (50.0%), the surgeon in $n = 6$ (27.3%), the anesthesiologist in $n = 1$ (4.5%), and unspecified in $n = 3$ (13.6%) facilities. FAST indications varied between circulatory instability $n = 8$ (36.4%), team leader's discretion $n = 6$ (27.3%), abdominal trauma $n = 3$ (13.6%), and not specified $n = 6$ (27.3%). Telephone interviews revealed that exams were always $n = 8$ (36.4%) or often $n = 4$ (18.2%) registered in the patients' charts. The remaining $n = 10$ (45.5%) facilities either never registered $n = 2$ (9.1%), it was not possible to register $n = 1$ (4.5%), or unknown by the trauma leaders $n = 7$ (31.8%). Images were often stored in $n = 1$ (4.5%), never stored in $n = 10$ (45.5%), not possible to store in $n = 2$ (9.1%), and unknown in $n = 9$ (40.9%) facilities.

Conclusion: Ultrasonography was used in a non-uniform fashion by multiple specialties in Danish trauma facilities. Very few images from FAST examinations were stored and documentation was scanty. National guidelines on application and documentation of ultrasonography in trauma are called for.

Keywords: Trauma, Ultrasonography, Focused Assessment with Sonography in Trauma, FAST, Emergency Medicine


Background

The Focused Assessment with Sonography in Trauma (FAST) protocol is considered beneficial in emergent evaluation of trauma patients with blunt or penetrating injury [1]. FAST is a focused abdominal and cardiac ultrasound examination designed for assessment of free fluid in the peritoneum and the pericardium. The examination was initially introduced in the early 1990's and has

spread throughout the world [2]. The protocol includes three views of the abdomen and one view of the heart, as shown in Fig. 1. The abdominal views are right upper quadrant, left upper quadrant, and pelvis for assessment of hemorrhage into the peritoneal cavity. The subcostal view of the heart is used for detecting blood in the pericardium. The parasternal long axis view can be used as an alternative if the subcostal view cannot be obtained [3]. Previous studies of sensitivity and specificity of FAST have found these to be 63–100% and 95–100%, respectively [4]. The likelihood ratio of positive test has been found to be >7.9 [5]. The FAST exam has been shown

*Correspondence: jesper.weile@clin.au.dk

² Research Center for Emergency Medicine, Aarhus University Hospital, Nørrebrogade 44, Building 1B, 8000 Aarhus C, Denmark
Full list of author information is available at the end of the article



to decrease time to intervention, complication rate, and hospital length of stay [6]. The FAST protocol is reproducible and not correlated to adverse effects from radiation[6].

In 2004, Kirkpatrick et al. presented an extension adding anterior and dorsolateral thoracic images of the pleura to the FAST exam [7]. See Fig. 1. The intention of the Extended Focused Assessment with Sonography in Trauma (eFAST) was to add detection of pneumothorax or hemothorax to the basic FAST. Ultrasonography has a higher sensitivity detecting pneumothorax than does antero-posterior chest X-ray in the supine position (88–98.1% vs. 50–75.5%) and can detect very small amounts of intrathoracic fluid [6–16].

In Denmark, there are no strict recommendations for trauma facilities to implement FAST, eFAST, or other ultrasound examinations as standard of care for traumatized patients. According to the Advanced Trauma Life Support (ATLS) protocol, FAST scanning should be performed as an extension to the primary survey [17], and the American College of Emergency Physicians (ACEP) recommends ultrasonography as initial diagnostic modality for blunt and penetrating abdominal trauma in hemodynamic unstable patients[18]. A previous Danish national study found that anesthesiologists or surgeons, present as consultants in the emergency department, perform ultrasonography in 43 and 18% of the cases, respectively [19]. It is unknown whether these are FAST examinations performed on trauma patients or other point-of-care examinations. Application and

development of ultrasonography in trauma care are difficult if little is known about the existing usage of ultrasonography in the trauma facilities.

This study aims to clarify the current use of ultrasonography in existing trauma facilities, including examination indication, ultrasound protocol choice, examiner characteristics, and examination documentation.

Methods

Inclusion criteria

All 22 hospitals in Denmark receiving acutely injured patients via ambulance services (air, ship, or ground) with a predefined multidisciplinary trauma team, were included in this study. We defined the trauma patient as a patient activating the trauma call where a multidisciplinary trauma team cares for the patient in the trauma room.

We conducted a cross-sectional observational study performed in two phases.

Phase 1

- We gathered all local guidelines for trauma patient care from these 22 hospitals. This was done by searching publicly available resources on the Internet. If guidelines could not be accessed online, the Emergency Department was contacted by e-mail, and it sent the guideline.
- The guidelines from trauma facilities were investigated for use of ultrasonography in trauma. All mentions of “ultrasonography,” “ultrasound,” or “FAST” were recorded.

Phase 2

- The anesthesiologist, the orthopedic surgeon, and the radiologist on call in every trauma facility were interviewed via telephone by use of a structured questionnaire regarding the use of ultrasonography during treatment of traumatized patients. The questionnaire can be found in the Additional file 1: Appendix.
- All phone calls were conducted during weekdays between 09:00 am to 08:00 pm. A phone call was repeated seven times if not answered. After the 7th missed call, an e-mail containing the same questionnaire as the structured telephone interview was sent to the department. The department was asked to forward the questionnaire to a doctor frequently involved in trauma care. A second reminder e-mail was sent, and if the department did not respond, it was designated as “not responding.”

Endpoints

Our primary endpoint was to establish the proportion of trauma facilities recommending ultrasonography in

local trauma care manuals. Secondary endpoints were to establish the specialty of the physician conducting the FAST examination, the indication(s) for performing FAST, and whether other ultrasonography examinations were recommended in trauma care. Lastly, we examined the proportion of facilities documenting examinations in the patient records and the frequency of trauma facilities storing images from sonography.

Statistics


Descriptive statistics was used to present actual numbers and percentages unless otherwise indicated in the text. We calculated Cohen's weighted kappa for multiple observers on interview answers as to who was performing the examinations. When interpreting Cohen's kappa, we used Landis and Koch's guidelines from 1977 [20]. The interobserver agreement was also expressed in percentage of agreement. Calculations were performed using Stata 13 (Statacorp, USA).

Results

Data were collected from August 2016 to December 2016. Trauma care manuals were obtained from all 22 facilities receiving traumatized patients in Denmark. A total of 64 (97.0%) out of 66 possible interviews were performed. Anesthesiologists, radiologists, and orthopedic surgeons on call on 22 hospitals were eligible for inclusion via telephone interviews. Anesthesiologists from all 22 hospitals participated in the study, while one eligible orthopedic department and one department of radiology failed to participate.


The FAST protocol was mentioned in 21 (95.5%) manuals for traumatized patients. One (4.5%) manual did not mention ultrasonography of any kind for trauma care.

The specialty of the performing physicians and the distribution of specialties are shown in Fig. 2. The results are from the 21 facilities mentioning ultrasonography in the trauma care manual, and from the interviews with trauma leaders from the 22 facilities. We interviewed orthopedic surgeons, anesthesiologists, and radiologists on call on the subject; Fig. 2 expresses the answer from the trauma leader. The numbers on the y axis are the number of hospitals. "Depending on competency" meant that the doctor present from either radiology, anesthesiology, or surgery with the highest level of competency performed the examination. In 8 (36.4%), all three interviewed physicians agreed on the person performing the FAST examination. In 6 (27.3%), two agreed, and in the remaining 8 (36.4%) there was total disagreement. Cohen's kappa on interrater agreement among radiologists, anesthesiologists, and orthopedic surgeons was $K = 0.35, p < 0.0001$ and interpreted as fair.



Indications for ultrasonography were either not mentioned, at the discretion of the trauma leader, abdominal trauma, or circulatory instability. The distribution is shown in Fig. 3. Because two trauma manuals stated indications as being abdominal trauma or circulatory instability, the total number of FAST indications ($n = 23$) was higher than the number of guidelines mentioning FAST ($n = 21$). The remaining facilities each stated a single indication as shown. One facility stated the indication as all circulatory unstable patients with suspected abdominal trauma; it was included as "circulatory instability" in the figure. In one manual, FAST exam was mentioned as a useful modality for "Trauma in Children." As this indication was unspecific, it was not included in the figure. It was the only manual mentioning ultrasonography specifically for children. In 16 (72.7%) manuals, the FAST exam was part of the primary survey, and in 2 (9.1%), it was part of the secondary survey. The remaining 3 (13.6%) trauma facilities did not specify when the examination was to be performed.

The use of the eFAST protocol for ruling out pneumothorax and hemothorax was mentioned in 6 (27.3%) facilities. Additional uses for ultrasound examinations were mentioned in 7 (31.8%) trauma manuals, categorized as follows. Ultrasonography of arterial flow in extremities was mentioned in 5 (22.7%) manuals, but none of the manuals specify which specialty should perform the examination. Ultrasonography of scrotal trauma performed by radiologist was mentioned in two (9.1%) manuals, and ultrasonography of internal organs by radiologist was mentioned in two (9.1%) manuals. One (4.5%)



manual mentioned focused cardiac ultrasonography in the primary survey, but it did not mention indication or who was to perform the examination. Lastly, one (4.5%) manual mentioned unspecified thoracic ultrasonography for penetrating trauma.


Trauma leaders were questioned in regards to the frequency of use of ultrasonography in trauma care. Answers are shown in Fig. 4.

Lastly, trauma leaders were questioned about the frequency of storing images and entering descriptions of the ultrasound examinations into medical charts. Results are shown in Fig. 5.

Discussion

The FAST exam was recommended in 21 out of 22 local guidelines. In addition, our study revealed three main findings. First, different specialties seem to perform ultrasound examinations in different trauma facilities. Second, we found various indications for performing FAST, ranging from all “abdominal trauma” to “trauma leader’s discretion.” Third, we found that documentation of ultrasonography examinations and storing of images was sparse.


An e-mail survey from 2008 in the U.S. showed that 85% of hospitals reported using the FAST examination for trauma care [2]. Our investigation showed a recommendation of FAST in 95.5% of all facilities receiving



traumatized patients in Denmark. The higher percentage in our study probably reflects the growing interest in ultrasonography over time. Repetition of the American study might reveal a higher percentage today, which is why we consider the difference as a reflection of general development in the area.

Our study revealed differing local recommendations regarding which specialties should perform the ultrasound examinations. We have not come across evidence in the literature that one specialty should outperform others. We found that fewer surgeons and radiologists, and more anesthesiologists, perform FAST examination according to interviews than according to the local trauma care guidelines. Earlier work has shown that the vast majority of physicians performing ultrasonography in the Emergency Department are anesthesiologists [19]. Our study suggests that in trauma it will more often be the person with the greatest skill who performs the FAST examination rather than the person recommended in the guidelines. These findings might reflect that anesthesiology often is the specialty present with the most experience, while radiologists may be on call but not present in the trauma room.

The FAST exam is user dependent, and the specificity and sensitivity are related to the experience of the provider. The examination has low sensitivity for organ injury without hemoperitoneum and low sensitivity for retroperitoneal bleeding [6]. To reach an overall sensitivity of 67% and specificity of 100%, extensive experience



of more than 100 FAST examinations is necessary [21]. A reliable and validated assessment tool for assessment of competency in point-of-care ultrasonography exists [22]. This evidence calls for a specification in the existing guidelines recommending the most experienced person to perform the examination. Preferably, the experience should be quantified, and an expected minimum of experience or a minimum required competency level should be outlined.

The indications for performing FAST differ between facilities, and, including extended FAST, views of the thorax are also heterogenic. Some facilities recommend other use of sonography beyond FAST, such as sonography of the scrotum. The addition of extended FAST views appears helpful in the literature as life-threatening injuries such as tension pneumothorax or massive hemothorax can be diagnosed at the bedside [11, 23]. Cardiac evaluation alone in penetrating trauma to the chest has been shown to decrease mortality [24]. Although the eFAST exam for trauma has not been shown to reduce mortality, it still has several advantages. Knowledge of thoracic injury with pneumothorax, hemothorax, or pericardial effusion at the time of arrival can be used to guide initial treatment even before a CT scan is performed. Future trials should seek to illuminate the clinical strength of the full eFAST over the FAST alone.

To our knowledge, the frequency of ultrasonography images being stored and documented has not previously been investigated in the trauma setting. Recommendations have been published on how to store images and document examinations [25], but these are not followed. Our findings, which show a surprisingly low frequency of storage and documentation, call for attention to the problem in the local guidelines. Without proper documentation, developing the field is immensely difficult. Furthermore, lack of image storing may present legal issues in the future.

Our study has limitations. First, the study is national and limited to the Danish hospitals receiving traumatized patients. However, our findings of particular problems in heterogeneity between facilities and scanty documentation undoubtedly exist abroad, and our call for national guidelines can only inspire other countries to do the same. A second limitation is ascribing value to interviews of the trauma team leaders and other doctors on call. We do not know whether the results would have been different if the phone call had been made on a different day or whether the answers were influenced by recollection bias.

Conclusion

Ultrasonography is applied in a heterogeneous manner by multiple specialties on multiple indications in trauma care in Denmark. Storage and documentation of examinations is sparse and desultory. Multispecialty national evidence-based guidelines, as well as unified implementation of existing guidelines, are called for.

Additional file

[Additional file 1.](#) Appendix 1

Abbreviations

ATLS: Advanced Trauma Life Support; eFAST: extended Focused Assessment with Sonography in Trauma; FAST: Focused Assessment with Sonography in Trauma.

Authors' contributions

JW, KN, ES, HK, CAF, and CBL contributed to conception and design the study. JW, SCP, and KN performed the telephone interviews. JW collected all trauma care manuals and drafted the manuscript. All authors critically revised drafts for important intellectual content and approved the final manuscript submitted. All authors read and approved the final manuscript.

Author details

¹ Emergency Department, Regional Hospital Herning, Herning, Denmark. ² Research Center for Emergency Medicine, Aarhus University Hospital, Nørrebrogade 44, Building 1B, 8000 Aarhus C, Denmark. ³ Department of Medicine, Section of Respiratory Medicine, University Hospital Hvidovre, Hvidovre, Denmark. ⁴ Aarhus University, Aarhus, Denmark. ⁵ Department of Cardiology, Aarhus University Hospital, Aarhus, Denmark. ⁶ Department of Respiratory Medicine, Odense University Hospital, Odense, Denmark. ⁷ Department of Anesthesiology and Intensive Care, Aarhus University Hospital, Aarhus, Denmark. ⁸ Research Center for Emergency Medicine, Aarhus University Hospital, Aarhus, Denmark.

Acknowledgements

The authors would like to thank all departments and physicians participating in the interviews. Without this great help, this study could not have been performed. The authors especially wish to thank medical student Stig Holm Jensen for contributing to the data collection.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

For access to full dataset please contact corresponding author by e-mail.

Ethics

The Central Denmark Region Committees on Health Research Ethics granted a waiver for the study cf. inquiry 111/2015. All participants in the phone interviews gave oral informed consent or consented by e-mail if an e-mail questionnaire was forwarded.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 7 May 2017 Accepted: 13 June 2017

Published online: 20 June 2017

References

- Melniker LA, Leibner E, McKenney MG, Lopez P, Briggs WM, Mancuso CA (2006) Randomized controlled clinical trial of point-of-care, limited ultrasonography for trauma in the emergency department: the first sonography outcomes assessment program trial. *Ann Emerg Med* 48:227–235
- Scaife ER, Fenton SJ, Hansen KW, Metzger RR (2009) Use of focused abdominal sonography for trauma at pediatric and adult trauma centers: a survey. *J Pediatr Surg* 44:1746–1749
- American Institute of Ultrasound in (2014) M, American College of Emergency P. AIUM practice guideline for the performance of the focused assessment with sonography for trauma (FAST) examination. *J Ultrasound Med* 33:2047–2056
- Smith ZA, Wood D (2014) Emergency focussed assessment with sonography in trauma (FAST) and haemodynamic stability. *Emerg Med J* 31:273–277
- Quinn AC, Sinert R (2011) What is the utility of the Focused Assessment with Sonography in Trauma (FAST) exam in penetrating torso trauma? *Injury* 42:482–487
- Ollerton JE, Sugrue M, Balogh Z, D'Amours SK, Giles A, Wyllie P (2006) Prospective study to evaluate the influence of FAST on trauma patient management. *J Trauma* 60:785–791
- Kirkpatrick AW, Sirois M, Laupland KB, Liu D, Rowan K, Ball CG et al (2004) Hand-held thoracic sonography for detecting post-traumatic pneumothoraces: the Extended Focused Assessment with Sonography for Trauma (EFAST). *J Trauma* 57:288–295
- Knudtson JL, Dort JM, Helmer SD, Smith RS (2004) Surgeon-performed ultrasound for pneumothorax in the trauma suite. *J Trauma* 56:527–530
- Lichtenstein D, Meziere G, Biderman P, Gepner A (2000) The "lung point": an ultrasound sign specific to pneumothorax. *Intensive Care Med* 26:1434–1440
- Rowan KR, Kirkpatrick AW, Liu D, Forkheim KE, Mayo JR, Nicolaou S (2002) Traumatic pneumothorax detection with thoracic US: correlation with chest radiography and CT—initial experience. *Radiology* 225:210–214
- Blaivas M, Lyon M, Duggal S (2005) A prospective comparison of supine chest radiography and bedside ultrasound for the diagnosis of traumatic pneumothorax. *Acad Emerg Med* 12:844–849
- Whitson MR, Mayo PH (2016) Ultrasonography in the emergency department. *Crit Care* 20:227
- Abu-Zidan F (1997) The role of ultrasonography and ultrasound-guided aspiration in blunt abdominal trauma. *Ann Saudi Med* 17:569–570
- Rozycki GS, Feliciano DV, Davis TP (1998) Ultrasound as used in thoraco-abdominal trauma. *Surg Clin North Am* 78:295–310
- Ding W, Shen Y, Yang J, He X, Zhang M (2011) Diagnosis of pneumothorax by radiography and ultrasonography: a meta-analysis. *Chest* 140:859–866
- Alrajhi K, Woo MY, Vaillancourt C (2012) Test characteristics of ultrasonography for the detection of pneumothorax: a systematic review and meta-analysis. *Chest* 141:703–708
- Subcommittee A, American College of Surgeons' Committee on T, International Awg (2013) Advanced trauma life support (ATLS(R)): the ninth edition. *J Trauma Acute Care Surg* 74:1363–1366.
- American College of Emergency P (2009) Emergency ultrasound guidelines. *Ann Emerg Med* 53:550–570
- Nielsen K, Lauridsen JR, Laursen CB, Brabrand M (2015) Physicians using ultrasound in Danish emergency departments are mostly summoned specialists. *Scand J Trauma Resusc Emerg Med* 23:51
- Landis JR, Koch GG (1977) An application of hierarchical kappa-type statistics in the assessment of majority agreement among multiple observers. *Biometrics* 33:363–374
- Gracias VH, Frankel HL, Gupta R, Malczynski J, Gandhi R, Collazzo L et al (2001) Defining the learning curve for the Focused Abdominal Sonogram for Trauma (FAST) examination: implications for credentialing. *Am Surg* 67:364–368
- Todsen T, Tolsgaard MG, Olsen BH, Henriksen BM, Hillingsø JG, Konge L et al (2015) Reliable and valid assessment of point-of-care ultrasonography. *Ann Surg* 261:309–315
- Lee C, Revell M, Porter K, Steyn R (2007) Faculty of Pre-hospital Care RCoSoE. The prehospital management of chest injuries: a consensus statement. Faculty of Pre-hospital Care, Royal College of Surgeons of Edinburgh. *Emerg Med J* 24:220–224
- Arntfield RT, Millington SJ (2012) Point of care cardiac ultrasound applications in the emergency department and intensive care unit—a review. *Curr Cardiol Rev* 8:98–108
- Laursen CB, Nielsen K, Riishede M, Tiwald G, Mollekaer A, Aagaard R et al (2014) A framework for implementation, education, research and clinical use of ultrasound in emergency departments by the Danish Society for Emergency Medicine. *Scand J Trauma Resusc Emerg Med* 22:25

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com