

Information and Communication Technologies for the Activities of Daily Living in Older Patients with Dementia: A Systematic Review

Grazia D'Onofrio^{a,b,*}, Daniele Sessa^a, Francesco Di Giordano^c, Francesco Panico^{a,d,e}



[Metadata, citation and similar papers at core.ac.uk](https://www.core.ac.uk)

Provided by Archivio della ricerca della Scuola Superiore Sant'Anna

^aIRCCS “Casa Sollievo della Sofferenza”, San Giovanni Rotondo, Foggia, Italy

^bThe BioRobotics Institute, Scuola Superiore Sant'Anna, Pontedera, Italy

^cICT, Innovation & Research Unit, IRCCS “Casa Sollievo della Sofferenza”, San Giovanni Rotondo, Foggia, Italy

^dDepartment of Basic Medicine, Neurodegenerative Disease Unit, Neuroscience, and Sense Organs, University of Bari Aldo Moro, Bari, Italy

^eDepartment of Clinical Research in Neurology, University of Bari Aldo Moro, “Pia Fondazione Cardinale G. Panico”, Tricase, Lecce, Italy

Accepted 7 February 2017

Abstract.

Background: Significant innovations have been introduced in recent years in the application of information and communication technologies (ICTs) to support healthcare for patients with dementia.

Objective: In the present systematic review, our goal is to keep track of ICT concepts and approaches to support the range of activities of daily living for people with dementia and to provide a snapshot of the effect that technology is having on patients' self-reliance.

Methods: We reviewed the literature and identified systematic reviews of cohort studies and other authoritative reports. Our selection criteria included: (1) activities of daily living, (2) ICT, and (3) dementia.

Results: We identified 56 studies published between 2000 and 2015, of which 26 met inclusion criteria. The present systematic review revealed many ICT systems that could purportedly support the range of activities of daily living for patients with dementia. The results showed five research bodies: 1) technologies used by patients with dementia, 2) technologies used by caregivers, 3) monitoring systems, 4) ambient assistive living with ICTs, and 5) tracking and wayfinding.

Conclusions: There is a potential for ICTs to support dementia care at home and to improve quality of life for caregivers, reducing healthcare costs and premature institutional care for these patients.

Keywords: Basic activities of daily living, dementia, information and communication technologies, instrumental activities of daily living

INTRODUCTION

In recent years, there have been significant innovations in the application of information and communication technologies (ICT) to support healthcare for patients with Alzheimer's disease (AD) and

*Correspondence to: Grazia D'Onofrio, Geriatric Unit & Gerontology-Geriatrics Research Laboratory, IRCCS Casa Sollievo della Sofferenza, Viale Cappuccini 1, 71013 San Giovanni Rotondo, Foggia, Italy. Tel./Fax: +39 0882 410271; E-mail: g.donofrio@operapadrepio.it.

dementia. These technologies can be used by patients with dementia and caregivers, and can run automatically; in this latter case we referred to them with the term “ambient intelligence” [1]. Many ICT systems are currently being developed and could potentially support a patient with dementia in the activities of daily living [2].

The activities of daily living can be included in two broad classes: 1) Basic Activities of Daily Living (BADL) [3, 4] and 2) Instrumental Activities of Daily Living (IADL) [5].

BADL are physical tasks essential to maintaining one’s independence and include the ability to go to the toilet, to feed, to dress, to groom, to bathe, and to ambulate. IADL are typically more cognitively demanding than BADL, and include the ability to successfully use the telephone, do shopping, prepare food, do the housekeeping and laundry, manage medications and finances, and use the transportation means outside of the home (e.g., driving a car, using public transit, or riding in a taxi).

In the early stage of dementia, most people are independent with BADL, but they start to need help with some IADL, especially complex tasks requiring multiple steps or extensive planning [6]. As dementia progresses to the moderate stage, IADL such as working, medication management, and keeping track of personal finances become difficult or even impossible to perform, and a person may begin to need help with BADL. In the moderate phase of dementia, cooking, housework, and shopping may require direct assistance, BADL require assistance for set-up and safety, and completing BADL may be disrupted by neuropsychiatric symptoms (NPS) such as anger, frustration, and difficulty in communicating needs [7]. As dementia enters the severe stage, independence is gradually lost and caregivers must provide consistent direct care with most, if not all, BADL. At this stage, a person must be directly assisted with simple BADL such as eating, bathing, transfers, and walking [8]. Safety issues and wandering require constant monitoring [9], and there is the need to create a safe environment [10, 11].

Therefore, people with dementia experience progressive cognitive impairments, and more innovative approaches need to be developed to help promote independence and maximize quality of life. In this context, ICTs may offer a lot of potential and can make a significant difference in the lives of people with dementia and to their primary caregivers. Indeed, it has been noted that these technologies

should be part of a home package and should be provided in a thoughtful, sensitive, and ethical way [12]. However, the overall opportunities that technology could create for people with dementia have not been fully clarified to date. The objective of the present systematic review is to investigate the use of ICTs to support the range of activities of daily living for people with dementia, considering “ambient intelligence” technologies that can be used by patients and caregivers.

MATERIAL AND METHODS

The search analysis was informed by the study’s aims, previous reviews using qualitative data, and best practice recommendations in the research literature [13, 14]. We reviewed clinical and epidemiological reports from the international literature published from January 2000 to October 2015. This systematic review was based upon searches of the US National Library of Medicine (PubMed), Ovid MEDLINE, EMBASE, Google Scholar, Web of Science, and Scopus databases. The search queries included the following terms (Information and Communication Technologies OR ICT) combined with terms to determine the outcomes of interest (cognitive AND [impairment OR decline OR disorders] OR Alzheimer’s disease OR dementia) and were limited to human studies. Only English language articles were included, due to a lack of resources for translation. Reference lists of included articles and relevant review articles were examined to identify every study which the electronic search strategy may have missed.

Study selection

A single reviewer (GDO) examined abstracts retrieved by the electronic search to identify articles deserving a full review. Full length articles were then reviewed before data were extracted from relevant papers. The inclusion/exclusion criteria used for our review protocol are the following: inclusion criteria were: 1) age ≥ 60 years; 2) diagnosis of dementia according to the National Institute on Aging-Alzheimer’s Association criteria [15], and 3) widely accepted clinical measures of cognitive impairment, disability, quality of life, and global clinical assessments. Exclusion criteria were: 1) no English language (as we lacked resources for translation) and 2) no diagnosis of dementia.

Data extraction

For the present systematic review, from the 170 records identified through database searching, 57 published studies were deemed relevant. Of these, 12 studies were excluded because they were not focused on the activities of daily living. After assessment of the remaining full-text articles, we excluded other 18 studies, 10 of which did not evaluate patients with dementia, and 8 that were not focused on the use of ICT. Thus, the final list included 27 published studies that were eligible for the current systematic review (Fig. 1).

An inductive approach to the analysis was used. The results section was divided in five categories: 1) technologies used by patients with dementia, 2) technologies used by caregivers, 3) monitoring systems, 4) ambient assistive living (AAL) with ICT, and 5) tracking and wayfinding. Other reviewers (DSA, FR, FC, and FG) provided a detailed summary of each study, including its strengths and weaknesses, as well as an overall review of the category [16, 17]. Through this process, the following items were systematically extracted from the articles: approach, methodology, transparency, strengths, and weaknesses. These were then organized according to theme, in order to provide an overview of the state of the field as a whole. Quality




Fig. 1. Flow diagram outlining the selection procedure to identify articles which were included in the systematic review of information and communication technologies (ICTs) and activities of daily living and in patients with dementia.

of study reporting was assessed using the Standards for the Reporting of Diagnostic accuracy studies in dementia (STARDdem) [18].

RESULTS

The potential ICTs that support the range of activities of daily living for people with dementia are shown in Table 1, summarizing the 26 published studies selected for the current systematic review [1, 19–38].

Technologies used by patients with dementia

In the early stages of dementia, ICTs can directly support the patients. Several studies have assessed online resources for patients with dementia focused on maintaining cognitive and functional skills, learning new things, maintaining social interactions, and looking for information [19]. Examples include electronic applications providing reminders (e.g., medication management prompting devices), social contact (e.g., cell phones, online chat groups), safety (e.g., alarm systems and action triggered lighting), and daily activities (e.g., music players).

Technologies used by caregivers

Currently, ICT devices are available to support formal and informal caregivers. An example was the internet-based Savvy Caregiver [20] for informal caregivers. It was based on a face-to-face caregiver-training program and curriculum [21] and consists of a customized computer–telephone integration system that provides a psychoeducational intervention for informal caregivers to identify resources and strategies to enhance safety, self-care, social support, communication, and behavior management [22]. It was shown that this system significantly decreased caregiver burden and depression in informal caregivers [18]. In a pilot study, another type of in-home caregiver support linked informal caregivers with expert guidance for managing challenging care situations using video monitoring, and demonstrated improved behavior management and caregiver communication [23]. Caregivers were trained to capture behaviors that were a problem via computer video recording, which was then wirelessly uploaded to a team of experts to review and provide feedback. Support for caregivers can also be provided through text-based chat forums and web-based video conferencing [24]. A recent study had administered a questionnaire to Canadian unpaid caregivers

Table 1
Current use of information and communication technologies (ICTs) for activities of daily living in patients with dementia according to the identified research areas

Research areas identified	Studies	Methods	Outcomes
Technologies used by patients with dementia	Rosenberg et al., 2011 [19]	AT	To explore actions and driving forces of the actors involved in the process of bringing AT into the life of a person with dementia
Technologies used by caregivers	Hepburn et al., 2003 [20]	SCP	Better caregiver well-being
	Finkel et al., 2009 [22]	E-care	Significant decrease in post-intervention burden
	Lewis et al., 2010 [21]	IBSC	To strengthen family caregivers' confidence in caring for persons with dementia
	Marziali et al., 2011 [24]	2-IBIP	To improve mental health status and improve personal characteristics with associated lower caregiver stress response
	Williams et al., 2013 [23]	IHMSDC	To improve communication and behavior management and ease of use
Monitoring system	Czarnuch et al., 2016 [25]	BADLS	To improve ICT applicability and acceptance for unpaid caregivers.
	Carswell et al., 2009 [27]	AT	To enhance wellbeing and maintain activity levels
	Lexis et al., 2013 [29]	AMT	Support of frail older people to live longer independently
	Doctor et al., 2014 [28]	AI	To better approximate the learnt behaviors and to identify long-term macro-level behavior changes
Ambient assistive living with ICT	Peetoom et al., 2015 [26]	MT	To demonstrate the potential to prolong independent living of elderly persons
	Cook et al., 2009 [1]	AI + VR	To trigger broad empowerment processes induced by a strong sense of presence, leading to greater agency and control over one's actions and environment
	Rashidi et al., 2013 [30]	AAL	To improve safe and independent aging
	McKenzie et al., 2013 [31]	SHP	To ameliorate some of the stress and burden associated with providing care for persons with dementia
	Rantz et al., 2013 [32]	ST	Monitoring individuals' health status, detecting emergency situations, and notifying health care providers
	Jekel et al., 2016 [33]	PFOAT	Smart home technologies offer the chance for an objective and ecologically valid assessment of IADL.
	Realdon et al., 2016 [34]	TEMD	To ensure the continuity of care from clinical practice to the patient's home, enabling also cost effectiveness and the empowerment of patient and caregiver in the care process.
	Lazarou et al., 2016 [35]	IHMS	To monitor problematic daily living activity areas and design personalized interventions based on system feedback and clinical observations.
	Esposito et al., 2015 [36]	SR	To improve the quality of assistive services.
	Wang et al., 2017 [37]	RS	To decrease frustration, stress, relationship strain, and increase social interaction via the robot
Tracking and wayfinding	Chang et al., 2010 [38]	RFID	To serve as good context for triggering navigation prompts, although individual differences in effectiveness varied
	Lancioni et al., 2011 [41]	BOT	To orient their travel and find the rooms correctly
	Grierson et al., 2011 [40]	TWFD	To make the individuals' houses bound and able to perform daily activities without significant frustrations
	Pot et al., 2012 [43]	GPS	Useful for people in early stages of dementia and their informal caregivers
	Pulli et al., 2012 [39]	MATbSNC	Increased awareness of the target users and their relatives or other people in charge
	Tchalla et al., 2013 [44]	HBTEC-TS	Reduced incidence of primary indoor falling needing GP intervention or attendance at an emergency room among elderly people with AD and mild-to-moderate dementia
	Lancioni et al., 2014 [42]	AT	Increased positive performance of patients with dementia

AAL, ambient-assisted living; AI, ambient intelligent; AMT, activity monitoring technology; AT, assistive technology; BADLS, Bristol Activities of Daily Living Scale; BOT, basic orientation technology; GPS, global positioning system; HBTEC-TS, home-based technologies coupled with teleassistance service; IADL, Instrumental Activities of Daily Living; IBSC, Internet-Based Savvy Caregiver; IHMS, Intelligent Home Monitoring System; IHMSDC, In-home monitoring support for dementia caregivers; MATbSNC, Mobile Augmented Teleguidance-based Safety Navigation Concept; MT, monitoring technologies; PFOAT, Proxy-Free Objective Assessment Tool; RFID, radio-frequency identification; RS, robotic solution; SCP, savvy caregiver program; SHP, Safe Home Program; SR, service robotics; ST, sensor technology; TEMD, technology-enhanced multi-domain; TWFD, Tactile Way-Finding Device; VR, virtual reality; 2-IBIP, 2 Internet-based intervention programs.

208 of patients with dementia in order to develop a
209 model that uses 13 parameters to predict a per-
210 son with dementia's ability [25]. The 13 parameters
211 include caregiver relation, age, marital status, place
212 of residence, language, housing type, proximity to
213 caregiver, service use, informal primary caregiver,
214 diagnosis of Alzheimer's disease or dementia, time
215 since diagnosis, and level of dependence on care-
216 giver [25]. Pursuant to the authors of this study,
217 the knowledge of task independence can inform
218 the development of ICT for people with dementia,
219 improving their applicability and acceptance [25].

220 *Monitoring systems*

221 It is recognized that caregiving leads to excessive
222 levels of stress; thus, technology must be tailored to
223 address the determinants of caregiver stressors. For
224 example, caregivers are concerned about the safety
225 and security of their patients with dementia. It is
226 stressful if the caregiver needs to leave the dementia
227 patient at home alone for any length of time or while
228 they work, due to possible wandering, accidents, or
229 other negative events that may harm the patient. ICTs
230 may monitor whether or not the patient leaves his/her
231 home, falls, or enacts other behaviors with poten-
232 tially dangerous environmental conditions (e.g., heat,
233 water on the floor, and fire). ICTs may alleviate some
234 of these concerns. Many new systems provide these
235 types of monitoring and tracking systems. Three main
236 aims of monitoring are detection of BADL, occur-
237 rence of significant events (i.e., falls), and changes in
238 health status, or a combination of these [26].

239 Monitoring systems are designed to detect changes
240 in one or more BADL or physical parameters. Detec-
241 tion of subtle changes can trigger interventions to
242 avert negative outcomes, such as hospitalizations. An
243 example of a monitoring system is a motion detector
244 that turns a bathroom light on or a wireless home secu-
245 rity system that is triggered by exiting or entering,
246 activating an alarm. This technology records infor-
247 mation through activity sequence awareness, location
248 awareness, presence awareness, and context aware-
249 ness capabilities [27].

250 Alterations in BADL and IADL performance pro-
251 vided objective data about disease progression [28].
252 Through sensor technology, health care providers
253 were able to detect cognitive decline reflected in
254 behaviors such as getting out of bed at odd hours or
255 going to bed earlier and earlier from day to day [23].
256 A commercially available example is the QuietCare
257 system, which monitors BADL [29].

Ambient assistive living with ICT

258 AAL is called "ambient intelligent" and integrates
259 computer technology, electronics, and telecommu-
260 nications to support people in carrying out their
261 activities of daily life, and maintaining independent
262 living at home [30]. AAL is characterized by "sen-
263 sors and devices interconnected through a network
264 which senses features of the users and their envi-
265 ronment, then reasons about the accumulated data,
266 and finally, selects actions to take that will ben-
267 efit the users in the environment" [1]. The focus
268 of this technology is to empower human-machine
269 interactions using adaptive, sensitive, and responsive
270 strategies to human needs in digital environments
271 [26]. AAL systems can include multiple sensor
272 devices located within an environment, such as sensor
273 technology, software agents, radiofrequency identi-
274 fication, microchip implant, Bluetooth Low Energy,
275 biometrics, and affective computing. 276

277 Examples of complex AAL systems are the smart
278 homes in which remote network monitoring and data
279 exchange at a distance are used. AAL technologies
280 can monitor ambient temperatures, gas levels, and
281 motion, notify problematic changes to remote users,
282 and enable family and health care providers to predict
283 and intervene on impending incidents [31, 32].

284 The data are collected through infrared sensors
285 (detecting motion on kitchen cabinets while prepar-
286 ing meals) and bed pneumatic sensors (assessing
287 levels of pulse, respiration, and restlessness) [29].
288 These data are analyzed by developed algorithms
289 to identify functional decline and potential illnesses.
290 Then, health care providers such as nurse practition-
291 ers provide timely and effective care to improve and
292 maintain health and functional independence [29]. A
293 recent pilot study had shown that smart home tech-
294 nologies offer the chance for an objective and eco-
295 logically valid assessment of IADL [33]. The authors
296 had suggested that one can analyze not only whether
297 a task is successfully completed but also how it is
298 completed [33]. A single blind randomized controlled
299 trial had informed on the efficacy of a technology-
300 enhanced home care service to preserve cognitive and
301 motor levels of functioning in patients with mild cog-
302 nitive impairment and Alzheimer's disease in order
303 to slow down their loss of autonomy in daily life [34].
304 The expected outcome was to ensure the continuity of
305 care from clinical practice to the patient's home, also
306 enabling cost effectiveness and the empowerment of
307 patient and caregiver in the care process, positively
308 impacting on their quality of life [34]. 309

309 A study had proposed a system for continu- 359
310 ous and objective remote monitoring of problematic 360
311 daily living activity areas and to design personalized 361
312 interventions based on system feedback and clinical 362
313 observations to improve cognitive function and 363
314 health-related quality of life [35]. The assistive tech- 364
315 nology of the proposed system, including wearable 365
316 sleep, object motion, presence, and utility usage sen- 366
317 sors, was methodically deployed at four different 367
318 home installations of people with cognitive impair- 368
319 ment. It has been proved that the proposed system 369
320 was suitable to support clinicians to reliably drive 370
321 and evaluate clinical interventions for the improve- 371
322 ment of quality of life improvement in people with 372
323 cognitive impairment and dementia [35]. 373

324 Advances in service robotics for assisted living 374
325 applications [36] have facilitated the deployment 375
326 of ICT Robotics solutions to improve the quality 376
327 of assistive services. Indeed, another recent study 377
328 explored perspectives of older adults with dementia 378
329 and their caregivers on robots that provide stepwise 379
330 prompting to complete activities in their home [37]. 380
331 In this study, positive consequences of robots in care- 381
332 giving scenarios were shown including decreased 382
333 frustration, stress, relationship strain, and increased 383
334 social interaction via the robot [37]. 384

335 *Tracking and wayfinding*

336 Dementia patients can be supported in living 385
337 safely by wayfinding technologies improving inde- 386
338 pendent mobility at home and in the community [38]. 387
339 Wayfinding systems typically use global positioning 388
340 systems (GPS) that allow dementia patients in earlier 389
341 stages to increase freedom, autonomy, and confidence 390
342 in being able to go outdoors independently and with- 391
343 out fear of being lost [39]. Orientation assistance can 392
344 be given through three modalities, visual, audio, and 393
345 tactile signals, or a combination, and they have been 394
346 tested indoors and outdoors [39]. The tactile-based 395
347 system can use four small vibrating motors attached 396
348 to a wearable belt with an integrated GPS, three-axis 397
349 compass, inertial sensor, and an algorithmic executive 398
350 processor that provides the patients with dementia 399
351 with direction-relevant cues on which way to go [40]. 400
352 Regarding this system, the usefulness is limited to 401
353 dementia patients in mild stages. 402

354 Another navigation system using scenario-based 403
355 video clips embedded with Bluetooth sensor technol- 404
356 ogy was developed and tested with some success on 405
357 cognitively intact individuals [30]. Studies were con- 406
358 ducted on navigational systems using either light or 407

auditory cues, finding consistently that those visual 359
cues were the more effective wayfinding cues for 360
dementia patients [41, 42]. In contrast, tracking 361
devices use similar technology but focus on provid- 362
ing caregivers with a way to know the location of 363
the patient with dementia and to prevent and inter- 364
vene in unsafe situations. A pilot study evaluated the 365
feasibility and acceptability of a GPS device with 366
expanded features for tracking location and tracing 367
paths through satellite, programming telephone con- 368
nection between the patients with dementia and the 369
family caregivers, and activating a loudspeaker func- 370
tion to communicate with the patients in case they 371
cannot use the phone [43]. This study found that the 372
GPS intervention increased the ability of the patients 373
to go outside independently, resulting in more free- 374
dom from the caregivers and decreased levels of 375
stress for both patients and caregivers [39]. Another 376
example using complex tracking programs was the 377
prevention of night-time falls in patients with demen- 378
tia. The HBTec-TS tracking system incorporated a 379
nightlight path device that is installed near the bed 380
and was triggered to light up automatically when the 381
person stepped out of bed. The light guided the patient 382
to the bathroom at night [44]. A recent study using 383
the HBTec-TS system reported a significant decrease 384
in night-time falls in the intervention group com- 385
pared with the control group of community-dwelling 386
dementia patients (odds ratio = 0.37, 95% confidence 387
interval = 0.15–0.88, $p = 0.02$). This system reduced 388
the relative risk of falls by 48.8% in dementia patients 389
at high risk for frequent falls [40]. 390

391 **DISCUSSION**

392 The present systematic review, using an inductive 392
393 approach to the analysis, identified 21 studies reveal- 393
394 ing that many ICT systems could purportedly support 394
395 the range of activities of daily living for dementia 395
396 patients, showing five principal research bodies: 1) 396
397 technologies used by patients with dementia, 2) tech- 397
398 nologies used by caregivers, 3) monitoring systems, 398
399 4) AAL with ICTs, and 5) tracking and wayfind- 399
400 ing. This increasing body of evidence suggested the 400
401 potential for ICTs to support dementia care at home 401
402 and to improve quality of life for caregivers, thus 402
403 reducing healthcare costs and premature institutional 403
404 care for these patients. 404

405 The purpose of the present systematic review was 405
406 to investigate the use of ICTs to support the range 406
407 of activities of daily living for people with dementia 407

408 considering the technologies of “ambient intelli-
409 gence” and the technologies that can be used by
410 patients and caregivers. As the population with
411 dementia expands and the burden on family care-
412 givers increases, ICT applications have the potential
413 to support aging in place for patients with demen-
414 tia while reducing caregiver burden and its negative
415 outcomes, improving quality of life for families expe-
416 riencing dementia and reducing health care costs.
417 Maintaining independence and a good quality of life
418 might be challenging for some patients with demen-
419 tia, as for them the activities of daily living are
420 difficult to perform. Thanks to the recent advances in
421 ICTs, they may effectively be used to support older
422 adults in these activities.

423 In the last ten years, an increasing body of evidence
424 suggested significant benefits in detecting and mon-
425 itoring early cognitive impairment in AD and other
426 forms of dementia and capturing markers of intra-
427 individual change over time, in real time, and real
428 life situations using ICT devices also in clinical tri-
429 als [45] or assessing specific cognitive domain (e.g.,
430 apathy) [46]. Furthermore, other reviews on ICT use
431 in patients with dementia focused on the activities of
432 daily living also identifying different [47] or specific
433 research areas [48]. In fact, Lauriks and colleagues,
434 in their systematic review on ICT-based services for
435 unmet needs of people with dementia and their infor-
436 mal caregivers, identified need areas including 1) the
437 need for general and personalized information; 2) the
438 need for support to manage symptoms of dementia;
439 3) the need for social contact and company; and 4) the
440 need for health monitoring and perceived safety [47].
441 More recently, Teipel and colleagues reviewed only
442 ICT-based possible solutions for supporting outdoor
443 and social activities in patients with dementia, sug-
444 gesting that ICT services should be proposed at the
445 prodromal stage of dementia and should be carefully
446 validated within the life space of users in terms of
447 quality of life, social activities, and costs. [48]. Very
448 recently, a variety of ICT-based recreational systems
449 have been also developed and evaluated to support
450 patients with dementia in engaging in recreational
451 activities such as social interactions with friends and
452 families or playing games [49].

453 From our search, we identified 21 studies regard-
454 ing ICTs that may assist some aspects of ADLs
455 and IADLs in patients with dementia and that, with
456 further interdisciplinary research and modifications,
457 may have potential applications to dementia care. The
458 higher number of ICTs was designed to assist with
459 ambulation, housekeeping, and social communica-

460 tion, whereas a smaller number of ICTs were found to
461 support dressing and toileting. Eventually, the poten-
462 tial for ICTs to support dementia care at home can
463 reduce health care expenditures, secondary to formal
464 care needs and premature institutional care. How-
465 ever, exploiting the potential for technology to meet
466 dementia care needs depends on a number of factors,
467 including raising awareness of available technologies
468 and their utility, promoting accessibility and afford-
469 ability, and overcoming challenges to acceptance and
470 use. Future research, rigorous clinical trials, and con-
471 tinuous ICT developments are required to improve
472 the use of advanced technologies to be integrated
473 in current dementia care settings. Long term exper-
474 imentations are essential to assess the real efficacy
475 of ICTs, the acceptability and correct use by patients
476 and caregivers. Moreover, to explore the burden level
477 of caregivers that use ICTs in patient management
478 could be an important indicator of ICT effective-
479 ness. The issue of acceptability and usability was
480 another important issue to explore. In fact, patients
481 and their caregivers are sometimes skeptical about
482 accepting the installation and use of ICTs in their
483 life environments. The investigation of usability and
484 acceptability aspects of ICTs may be fundamental
485 to guarantee the suitability of these solutions in real
486 daily contexts. An idea for future studies could be
487 the implementation of technologies to detect uncor-
488 rected emotions and behaviors that can lead dementia
489 patients to a rapid progression of cognitive impair-
490 ment. Emotion and behavior detection in real time
491 by ICTs could reduce episodes of delirium, onset
492 of NPS, activate rehabilitative and/or relaxing proce-
493 dures, and improve the performances of the patients
494 in their activities of daily living.

495 ACKNOWLEDGMENTS

496 The research leading to these results has received
497 funding from the European Union Horizon 2020
498 Framework Programme for Research and Innovation
499 (2014–2020) under grant agreement 643808 Project
500 MARIO ‘Managing active and healthy aging with use
501 of caring service robots’.

502 Authors’ disclosures available online (<http://j-alz.com/manuscript-disclosures/16-1145r1>).

504 REFERENCES

- 505 [1] Cook DJ, Augusto JC, Jakkula VR (2009) Ambient
506 intelligence: Technologies, applications, and opportunities.
507 *Pervasive Mob Comput* 5, 277-298.

- 508 [2] Alzheimer's, Society (2015) "Assistive technology - devices
509 to help with everyday living". https://www.alzheimers.org.uk/site/scripts/documents_info.php?documentID=109,
510 Posted April 2015. Accessed May 26, 2016.
- 511 [3] Katz S, Downs TD, Cash HR, Grotz RC (1970) Progress
512 in the development of the index of ADL. *Gerontologist* **10**,
513 20-30.
- 514 [4] Lawton MP (1990) Aging and performance of home tasks.
515 *Hum Factors* **32**, 527-536.
- 516 [5] Lawton MP, Brody EM (1969) Assessment of older people:
517 Self-maintaining and instrumental activities of daily living.
518 *Gerontologist* **9**, 179-186.
- 519 [6] Wattmo C, Wallin AK, Minthon L (2013) Progression of
520 mild Alzheimer's disease: Knowledge and prediction models
521 required for future treatment strategies. *Alzheimers Res Ther* **5**, 44.
- 522 [7] Nowrangi MA, Lyketsos CG, Rosenberg PB (2015) Principles
523 and management of neuropsychiatric symptoms in
524 Alzheimer's dementia. *Alzheimers Res Ther* **7**, 12.
- 525 [8] Fraga FJ, Falk TH, Kanda PA, Anghinah R (2013) Characterizing
526 Alzheimer's disease severity via resting-awake EEG amplitude
527 modulation analysis. *PLoS One* **8**, e72240.
- 528 [9] Futrell M, Melillo KD, Remington R, Schoenfelder DP
529 (2010) Evidence-based guideline. Wandering. *J Gerontol Nurs* **36**, 6-16.
- 530 [10] Eshkoor SA, Hamid TA, Nudin SS, Mun CY (2014) A
531 research on functional status, environmental conditions, and
532 risk of falls in dementia. *Int J Alzheimers Dis* **2014**, 769062.
- 533 [11] Volicer L, van der Steen JT (2014) Outcome measures for
534 dementia in the advanced stage and at the end of life. *Adv Geriatr* **2014**, 346485.
- 535 [12] Marshall M (2006) Foreword. In *Assistive Technology in Dementia Care*, J. Woolham, eds. Hawker Publications Ltd, London UK.
- 536 [13] Dixon-Woods M, Agarwal S, Jones D, Young B, Sutton A (2005) Synthesising qualitative and quantitative evidence: A review of possible methods. *J Health Serv Res Policy* **10**, 45-53.
- 537 [14] Hannes K, Macaitis K (2012) A move to more systematic
538 and transparent approaches in qualitative evidence synthesis: Update on a review of published papers. *Qual Res* **12**, 402-442.
- 539 [15] McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR Jr, Kawas CH, Klunk WE, Koroshetz WJ, Manly JJ, Mayeux R, Mohs RC, Morris JC, Rossor MN, Scheltens P, Carrillo MC, Thies B, Weintraub S, Phelps CH (2011) The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* **7**, 263-269.
- 540 [16] Barnett-Page E, Thomas J (2009) Methods for the synthesis
541 of qualitative research: A critical review. *BMC Med Res Methodol* **9**, 59.
- 542 [17] Grant MJ, Booth A (2009) A typology of reviews: An analysis
543 of 14 review types and associated methodologies. *Health Info Libr J* **26**, 91-108.
- 544 [18] Noel-Storr AH, McCleery JM, Richard E, Ritchie CW, Flicker L, Cullum SJ, Davis D, Quinn TJ, Hyde C, Rutjes AW, Smailagic N, Marcus S, Black S, Blennow K, Brayne C, Fioravanti M, Johnson JK, Köpke S, Schneider LS, Simmons A, Mattsson N, Zetterberg H, Bossuyt PM, Wilcock G, McShane R (2014) Reporting standards for studies of diagnostic test accuracy in dementia: The STARDdem Initiative. *Neurology* **83**, 364-373.
- 545 [19] Rosenberg L, Nygard L (2011) Persons with dementia
546 become users of assistive technology: A study of the process. *Dementia* **11**, 135-154.
- 547 [20] Hepburn KW, Lewis M, Sherman CW, Tornatore J (2003) The Savvy Caregiver program: Developing and testing a transportable dementia family caregiver training program. *Gerontologist* **43**, 908-915.
- 548 [21] Lewis ML, Hobday JV, Hepburn KW (2010) Internet-based program for dementia caregivers. *Am J Alzheimers Dis Other Dement* **25**, 674-679.
- 549 [22] Finkel S, Czaja SJ, Schulz R, Martinovich Z, Harris C, Pezuto D (2007) E-care: A telecommunications technology intervention for family caregivers of dementia patients. *Am J Geriatr Psychiatry* **15**, 443-448.
- 550 [23] Williams K, Arthur A, Niedens M, Moushey L, Hutfles L (2013) In-home monitoring support for dementia caregivers: A feasibility study. *Clin Nurs Res* **22**, 139-150.
- 551 [24] Marziali E, Garcia LJ (2011) Dementia caregivers' responses to 2 internet-based intervention programs. *Am J Alzheimers Dis Other Dement* **26**, 36-43.
- 552 [25] Czarnuch S, Ricciardelli R, Mihailidis A (2016) Predicting the role of assistive technologies in the lives of people with dementia using objective care recipient factors. *BMC Geriatr* **16**, 143.
- 553 [26] Peetoom KK, Lexis MA, Joore M, Dirksen CD, De Witte LP (2015) Literature review on monitoring technologies and their outcomes in independently living elderly people. *Disabil Rehabil Assist Technol* **10**, 271-294.
- 554 [27] Carswell W, McCullagh PJ, Augusto JC, Martin S, Mulvenna MD, Zheng H, Wang HY, Wallace JG, McSorley K, Taylor B, Jeffers WP (2009) A review of the role of assistive technology for people with dementia in the hours of darkness. *Technol Health Care* **17**, 281-304.
- 555 [28] Doctor F, Iqbal R, Naguib RNG (2014) A fuzzy ambient intelligent agents approach for monitoring disease progression of dementia patients. *J Ambient Intell Human Comput* **5**, 147-158.
- 556 [29] Lexis M, Everink I, van der Heide L, Spreuwenberg M, Willems C, de Witte L (2013) Activity monitoring technology to support homecare delivery to frail and psychogeriatric elderly persons living at home alone. *Technol Disabil* **25**, 189-197.
- 557 [30] Rashidi P, Mihailidis A (2013) A survey on ambient assisted-living tools for older adults. *IEEE J Biomed Health Inform* **17**, 579-590.
- 558 [31] McKenzie B, Bowen ME, Keys K, Bulat T (2013) Safe home program: A suite of technologies to support extended home care of persons with dementia. *Am J Alzheimers Dis Other Dement* **28**, 348-354.
- 559 [32] Rantz MJ, Skubic M, Miller SJ, Galambos C, Alexander G, Keller J, Popescu M (2013) Sensor technology to support Aging in Place. *J Am Med Dir Assoc* **14**, 386-391.
- 560 [33] Jekel K, Damian M, Storf H, Hausner L, Frölich L (2016) Development of a proxy-free objective assessment tool of instrumental activities of daily living in mild cognitive impairment using smart home technologies. *J Alzheimers Dis* **52**, 509-517.
- 561 [34] Realdon O, Rossetto F, Nalin M, Baroni I, Cabinio M, Fioravanti R, Saibene FL, Alberoni M, Mantovani F, Romano M, Nemni R, Baglio F (2016) Technology-enhanced multi-domain at home continuum of care program with respect to usual care for people with cognitive impairment: The Ability-Telerehabilitation study protocol for a randomized controlled trial. *BMC Psychiatry* **16**, 425.

- 637 [35] Lazarou I, Karakostas A, Stavropoulos TG, Tsompanidis
638 T, Meditskos G, Kompatsiaris I, Tsolaki M (2016) A novel
639 and intelligent home monitoring system for care support
640 of elders with cognitive impairment. *J Alzheimers Dis* **54**,
641 1561-1591.
- 642 [36] Esposito R, Fiorini L, Limosani R, Bonaccorsi M, Manzi
643 A, Cavallo F, Dario P (2015) Supporting active and healthy
644 aging with advanced robotics integrated in smart environ-
645 nment. In *Optimizing Assistive Technologies for Aging*
646 *Populations*, Morsi YS, Shukla A, Rathore CP, eds. IGI
647 GLOBAL, pp. 46-77.
- 648 [37] Wang RH, Sudhama A, Begum M, Huq R, Mihailidis A
649 (2017) Robots to assist daily activities: Views of older
650 adults with Alzheimer's disease and their caregivers. *Int*
651 *Psychogeriatr* **29**, 67-79.
- 652 [38] Chang YJ, Wang TY (2010) Comparing picture and video
653 prompting in autonomous indoor wayfinding for individuals
654 with cognitive impairments. *Pers Ubiquitous Comput* **14**,
655 737-747.
- 656 [39] Pulli P, Asghar Z, Siitonen M, Niskala R, Leinonen E,
657 Pitkänen A, Hyry J, Lehtonen J, Kramar V, Korhonen M
658 (2012) Mobile augmented teleguidance-based safety nav-
659 igation concept for senior citizens. *AMK-lehti/J Finnish*
660 *Univ Appl Sci* **2**, 1-9.
- 661 [40] Grierson LEM, Zelek J, Lam I, Black SE, Carnahan H
662 (2011) Application of a tactile way-finding device to facili-
663 tate navigation in persons with dementia. *Assist Technol* **23**,
664 108-115.
- 665 [41] Lancioni G, Perilli V, Singh NN, O'Reilly MF, Sigafos
666 J, Bosco A, De Caro MF, Cassano G, Pinto K, Minervini
667 M (2011) Persons with mild or moderate Alzheimer's dis-
668 ease use a basic orientation technology to travel to different
669 rooms within a day center. *Res Dev Disabil* **32**, 1895-1901.
- 670 [42] Lancioni G, Sigafos J, O'Reilly MF, Singh NN (2014)
Assistive technology for people with severe/profound
intellectual and multiple disabilities. In *Assistive Technol-
ogies for People with Diverse Abilities*, Lancioni G, Singh
NN, eds. Springer, New York, pp. 277-313.
- [43] Pot AM, Willemse BM, Horjus S (2012) A pilot study on
the use of tracking technology: Feasibility, acceptability,
and benefits for people in early stages of dementia and their
informal caregivers. *Aging Ment Health* **16**, 127-134.
- [44] Tchalla AE, Lachal F, Cardinaud N, Saulnier I, Rialle V,
Preux PM, Dantoine T (2013) Preventing and managing
indoor falls with home-based technologies in mild and
moderate Alzheimer's disease patients: Pilot study in a com-
munity dwelling. *Dement Geriatr Cogn Disord* **36**, 251-261.
- [45] Robert PH, König A, Andrieu S, Bremond F, Chemin I,
Chung PC, Dartigues JF, Dubois B, Feutren G, Guillemaud
R, Kenisberg PA, Nave S, Vellas B, Verhey F, Yesava-
ge J, Mallea P (2013) Recommendations for ICT use in
Alzheimer's disease assessment: Monaco CTAD Expert
Meeting. *J Nutr Health Aging* **17**, 653-660.
- [46] König A, Aalten P, Verhey F, Bensadoun G, Petit PD, Robert
P, David R (2014) A review of current information and com-
munication technologies: Can they be used to assess apathy?
Int J Geriatr Psychiatry **29**, 345-358.
- [47] Lauriks S, Reinersmann A, Van der Roest HG, Meiland
FJ, Davies RJ, Moelaert F, Mulvenna MD, Nugent CD,
Dröes RM (2007) Review of ICT-based services for identi-
fied unmet needs in people with dementia. *Ageing Res Rev*
6, 223-246.
- [48] Teipel S, Babiloni C, Hoey J, Kaye J, Kirste T, Burmeister
OK (2016) Information and communication technology
solutions for outdoor navigation in dementia. *Alzheimers*
Dement **12**, 695-707.
- [49] Lazar A, Thompson HJ, Demiris G (2016) Design rec-
ommendations for recreational systems involving older
adults living with dementia. *J Appl Gerontol*, doi: 10.1177-
0733464816643880.