

2P
(mix)

HIGH VOLTAGE ELECTRON MICROSCOPY AND ELECTRON DIFFRACTION OF PYROXENES IN TYPE B LUNAR SAMPLES FROM APOLLO 11.

H. Fernández-Morán and M. Ohtsuki

ELECTRON MICROSCOPY LABORATORY
DEPARTMENT OF BIOPHYSICS
PRITZKER SCHOOL OF MEDICINE

get PRA

and S. Hafner and D. Virgo


DEPARTMENT OF THE GEOPHYSICAL SCIENCES

to be presented at the

APOLLO 11 LUNAR SCIENCE CONFERENCE
NASA - MANNED SPACECRAFT CENTER

HOUSTON, TEXAS

January 5-8, 1970



(NASA-CR-127444) HIGH VOLTAGE ELECTRON
MICROSCOPY AND ELECTRON DIFFRACTION OF
PYROXENES IN TYPE B LUNAR SAMPLES FROM
APOLLO 11. H. Fernández-Moran, et al
(Chicago Univ.) 8 Dec. 1969 23 p CSCL 03B G3/30
N72-28817
Unclas
15675

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
US Department of Commerce
Springfield, VA. 22151

THE UNIVERSITY OF CHICAGO
CHICAGO, ILLINOIS

HIGH VOLTAGE ELECTRON MICROSCOPY AND ELECTRON
DIFFRACTION OF PYROXENES IN TYPE B LUNAR
SAMPLES FROM APOLLO 11

Humberto Fernández-Morán and Mitsuo Ohtsuki
Electron Microscope Laboratory
Department of Biophysics
Pritzker School of Medicine
and

Stefan S. Hafner and David Virgo
Department of the Geophysical Sciences

The University of Chicago
Chicago, Illinois 60637

**Details of illustrations in
this document may be better
studied on microfiche**

HIGH VOLTAGE ELECTRON MICROSCOPY AND ELECTRON
DIFFRACTION OF PYROXENES IN TYPE B LUNAR
SAMPLES FROM APOLLO 11

By: Humberto Fernández-Morán, Mitsuo Ohtsuki
Electron Microscope Laboratory, Department of
Biophysics and
Stefan S. Hafner and David Virgo
Department of the Geophysical Sciences
The University of Chicago
Chicago, Illinois 60637

ABSTRACT:

Lunar pyroxene 10044 specimens cleaved and sectioned by diamond knife ultramicrotomy were examined by standard (75 to 100 kV) and high voltage (200 kV) electron microscopy and diffraction.

Salient findings based on evaluation of 2000 plates show uniform 300 to 600Å-wide bands, probably corresponding to single crystal domains, with lattice spacings of 2.5Å. These dense bands, found predominantly in lunar pyroxene, are absent in terrestrial pyroxene XYZ. Lattice spacings of 6.5Å in lunar pyroxene and 18.2Å in pyroxene XYZ were directly visualized. High resolution bright and dark field images of iron-rich and magnesium-rich crystals were compared with corresponding electron diffraction patterns. Possible relations of observed structures to magnetic domains were considered.

Prepared and submitted: December 8, 1969.

Electron Microscopy of Lunar Pyroxenes

In view of the unusual variations of chemical composition within each crystal of lunar pyroxene (1) and of the well known distinct phases of exsolution phenomena observed in terrestrial and meteoritic pyroxenes (2,3,4), study of the fine structure of lunar pyroxenes as revealed by electron microscopy is essential.

Separated lunar pyroxene 10044 crystals cleaved and sectioned by diamond knife ultramicrotomy (5) and mounted directly on thin film specimen grids (without water or solvent contamination) were examined by both standard (75 to 100 kV) (6) and high voltage (200 kV) electron microscopy (fig. 1) and selected area electron diffraction techniques under conditions of higher penetration power, reduced radiation damage and negligible contamination (7) in a cryogenic vacuum (figs. 2,3,12).

Based on the examination of numerous representative samples and on the quantitative evaluation of 2000 plates, we can state the following observed characteristics:

1. Exceptionally regular, periodically spaced dense bands with uniform widths of 300 to 600Å (graphs). These straight-edged bands exhibit electron-optical phenomenon corresponding to single crystal domains (figs. 4,5,6,7,8,9,11), and they appear to be oriented with their long axis in the plane of the crystalline layers (approximately normal to crystallographic c).

Electron Microscopy of Lunar Pyroxenes

By combined high resolution dark field electron microscopy and selected area electron diffraction (fig. 6), intrinsic lattice spacings of 2.5\AA can be detected within the bands, arranged parallel to their long axis (i.e. normal to c).

Detailed studies show that these bands resemble electron-optical images of magnetic domain walls as seen in thin layers of ferromagnetic materials (8).

They are predominantly seen in iron-rich lunar pyroxene crystals, and the 2.5\AA spacings could correspond to a dense population of the iron atoms at the M positions within the bands.

The single crystal band domains are absent in both terrestrial pyroxene XYZ (figs. 4,10) and in magnesium-rich lunar pyroxene 10044 specimens. The latter show instead irregular striations along the planes of the cleaved lamellae. Dense granules (ca. 100 to 1000\AA in diameter) are also found in iron-rich pyroxene crystals (fig. 5).

2. Lattice spacings of 6.5\AA in lunar pyroxene and 18.2\AA in terrestrial pyroxene XYZ were directly visualized in high resolution bright and dark field images which could be compared with the corresponding

Electron Microscopy of Lunar Pyroxenes

electron diffraction patterns (fig. 4).

The 18.2Å spacings may tentatively be correlated with the a axis of pyroxene XYZ, and the 6.5Å spacings may correspond to the M-M interatomic distances in the cleavage planes.

These results, which are being further analyzed, are of particular importance in determining the cationic order-disorder phenomena in these silicates (9). However, more work must be carried out to establish the precise correlation with the unit cell dimensions of lunar pyroxene crystals.

The significance of present results indicates the potential contribution of correlated electron-optical and crystallographical studies to a better understanding of the intrinsic atomic organization of pyroxenes and their possible bearing on the nature and evolution of the moon.

December 8, 1969

Condensed version to be submitted for publication in Science.

Electron Microscopy of Lunar Pyroxenes

Correlation with new observations made by S. Hafner and D. Virgo on the magnetic behavior of pyroxenes in type B lunar samples as revealed by nuclear gamma ray resonance (NGR) studies.

Dr. Stefan Hafner and Dr. David Virgo (1) have independently made Mössbauer resonant absorption studies of ^{57}Fe in the same lunar pyroxene type B specimen 10044 and have demonstrated that the crystal structure in these specimens is ferrimagnetically ordered.

The lunar pyroxene crystals exhibit a sharp Curie point in the range of 10° to 20°K . The spin orientations in this lunar pyroxene are assumed to be ferrimagnetic.

This result is unusual. Chain silicate crystal structures are generally not magnetically ordered, even at very low temperatures (i.e. 1.7°K), particularly when the amount of diamagnetic cations (Mg, Ca, etc.) substituting for Fe is larger than 25 per cent as is the case in lunar augite. (G.K. Shenoy, G.M. Kalvius and S.S. Hafner, J. Appl. Phys. 40, p. 1314, 1969.)

We believe that the unusual ferrimagnetic ordering in lunar augite ($\text{Fe}_{0.34}\text{Mg}_{0.30}\text{Ca}_{0.36}\text{SiO}_3$) is due to iron-iron clustering in the ca. 300\AA wide single crystal domain bands depicted in the electron micrographs of the present report.

December 13, 1969

Electron Microscopy of Lunar Pyroxenes

REFERENCES AND NOTES

1. S. S. Hafner and D. Virgo, Science, 1970 (see special Lunar Symposium issue)
2. F. R. Boyd and G. M. Brown, Mineral Soc. Amer., Spec. paper 2, 211 (1969).
3. R. A. Binns and J. V. Long, Nature 198, 777 (1963).
4. M. B. Dube and L. T. Silver, Geochim. Cosmochim. Acta 31, 1637 (1967).
5. H. Fernández-Morán, Exper. Cell Res. 5, 255 (1953); Ind. Diam. Rev. 16, 128 (1956); J. Biophys. Biochem. Cytol. Suppl. 2, 29, (1956).
6. H. Fernández-Morán, Sixth International Congress for EM, Kyoto, 13, 27, 147 (1966).
7. H. Fernández-Morán, The Neurosciences, ed. F. O. Schmitt et al, 16, Rockefeller University Press, 281, (1967).
8. J. Silcox, Philos. Mag., 8, 7 (1963).
9. S. S. Hafner and D. Virgo, Science 165, 285 (1969)


Electron Microscopy of Lunar Pyroxenes

ACKNOWLEDGEMENTS

We thank Mr. C. L. Hough, Mr. C. Weber and Mr. G. Bowie for all the photographic reproduction; Mrs. V. Iglesias, Misses A. Hibino and M. Hanaoka, Mr. Ralph Vicario, Mr. H. Krebs, and Mr. G. Arcuri for specimen preparation; Miss S. Rowe for editorial assistance and Mrs. S. Erikson for assistance in preparation of this manuscript.



A special thanks is due Dr. George J. Jacobs, Chief, Physical Biology, Bioscience Programs, NASA Office of Space Science and Applications and Dr. Verl R. Wilmarth, Chief Lunar Scientist, NASA Manned Spacecraft Center for granting us permission to carry out extensive experiments on lunar rock samples.

Supported by the Pritzker Fund, the L. Block Fund, and the Otho Sprague Memorial Fund of the University of Chicago, by Grant NGL 14-001-012 of the National Aeronautics and Space Administration, and by Grant GM 13243 of the National Institutes of Health, General Medical Sciences.



HIGH VOLTAGE 200 K.V. H.U. 200E WITH SPECIALLY
DESIGNED LIQUID HELIUM SPECIMEN STAGE ATTACHED TO
CLOSED CYCLE SUPERFLUID LIQUID HELIUM REFRIGERATOR WITH
COLLINS/ADL HEAT EXCHANGER AND ACCESSORIES FOR SUPERCONDUCTING
HIGH VOLTAGE ELECTRON MICROSCOPE 1969

Fig.1



CORRELATED LIGHT MICROSCOPY AND HIGH VOLTAGE ELECTRON MICROSCOPY
STUDIES OF LUNAR SAMPLES RETURNED BY APOLLO 11



Reproduced from
best available copy.





Fig.2



CORRELATED LIGHT MICROSCOPY AND HIGH VOLTAGE ELECTRON MICROSCOPY STUDIES OF LUNAR SAMPLES RETURNED BY APOLLO 11

Fig.3








Fig-4



200 KV

2000 A

HIGH VOLTAGE ELECTRON MICROGRAPH OF IRON RICH LUNAR PYROXENE 10044 SHOWING UNIFORM DENSE BANDS AND IRREGULAR GRANULES



200KV


SELECTED AREA ELECTRON DIFFRACTION PATTERN OF IRON RICH LUNAR PYROXENE 10044



200 KV

2000 A


HIGH VOLTAGE ELECTRON MICROGRAPH OF MAGNESIUM RICH LUNAR PYROXENE 10044 SHOWING IRREGULAR STRIATIONS



200KV


SELECTED AREA ELECTRON DIFFRACTION PATTERN OF MAGNESIUM RICH LUNAR PYROXENE 10044

Fig.5




200KV

2000A




200KV

ELECTRON MICROGRAPH OF LUNAR PYROXENE 10044 SELECTED FOR COMPARING IMAGES OF BAND STRUCTURE WITH DIFFRACTION PATTERNS. CENTRAL BRIGHT FIELD IMAGE OF LUNAR PYROXENE TAKEN WITH SELECTOR APERTURE ON (000) AND DARK FIELD IMAGES CORRESPONDING TO ENLARGED DIFFRACTION SPOTS.



200KV



200KV

SELECTED AREA DIFFRACTION PATTERNS AND DARK FIELD IMAGES CORRESPONDING TO DENSE BAND STRUCTURES WITH CRYSTALLINE LATTICE SPACINGS OF 2.517 Å AND 0.843 Å.





Fig.7







Fig.8






Reproduced from
best available copy.

CORRELATED HIGH VOLTAGE ELECTRON MICROSCOPY AND INDEXED SELECTED AREA ELECTRON DIFFRACTION PATTERN OF CLINOPYROXENE CRYSTAL OBTAINED FROM LUNAR SAMPLE 10044 RETURNED BY APOLLO 11. SPACINGS OF 4.8 Å 2.77 Å 2.19 Å RECORDED BY MITSUO OHTSUKI ELECTRON MICROSCOPY LABORATORY H. FERNANDEZ MORAN UNIVERSITY OF CHICAGO


Fig.9



Reproduced from
best available copy.

CORRELATED HIGH VOLTAGE ELECTRON MICROSCOPY AND INDEXED SELECTED AREA ELECTRON
 DIFFRACTION OF TERRESTRIAL PYROXENE XYZ OBTAINED FROM DR. HAFNER. SPACINGS OF
 27.62 Å 7.79 Å 7.53 Å RECORDED BY MITSUO OHTSUKI. CALIBRATION WITH GOLD SINGLE
 CRYSTAL 200 PLANES 2.039 Å AND 220 PLANES 1.442 Å ELECTRON MICROSCOPY LABORATORY
 H. FERNANDEZ MORAN UNIVERSITY OF CHICAGO OCT. 1969

Fig.10



200 KV

LUNAR PYROXENE S 10044


5000 A

200 KV

LUNAR PYROXENE S 10044

5000 A

HIGH VOLTAGE BRIGHT AND DARK FIELD ELECTRON MICROGRAPHS OF CLEAVED LUNAR PYROXENE S10044 SHOWING DENSE BAND STRUCTURES



200 KV

5A

200 KV


5A

CORRESPONDING SELECTED AREA ELECTRON DIFFRACTION PATTERNS OF LUNAR PYROXENE S10044

Fig.11



SPECIAL ULTRAHIGH VACUUM CHAMBER FOR CONTROLLED MANIPULATION AND EXAMINATION OF LUNAR SAMPLES.



Reproduced from best available copy.






Fig.12



1 μ

The nucleation of domains in cobalt as the foil becomes thicker. The edge of the foil is shown as E.

Magnetic Domain Walls in Thin Films of Nickel and Cobalt

By J. SILCOX 1963

PHILOSOPHICAL MAGAZINE VIII / 7



LUNAR PYROXENE CRYSTALLINE LAYERS WITH



STRUCTURES RESEMBLING MAGNETIC DOMAIN

WALLS AS REVEALED BY HIGH VOLTAGE

ELECTRON MICROSCOPY 1969

H. FERNANDEZ MORAN AND M. OHTSUKI

LUNAR PYROXENE S 10044




200 KV

1000A


LUNAR PYROXENE CRYSTALLINE LAYERS WITH STRUCTURES RESEMBLING MAGNETIC DOMAIN WALLS AND STACKING FAULTS AS REVEALED BY SPECIAL ELECTRON MICROSCOPY TECHNIQUES

Reproduced from best available copy.




(a)

1 μ



(b)

1 μ



(c)

1 μ

Domains observed by stopping out one of the beams with the objective aperture. The position of the objective aperture in relation to the diffraction spots for each micrograph is shown in the inset. Magnification $\times 21\,000$.

Magnetic Domain Walls in Thin Films of Nickel and Cobalt

By J. SILCOX

Cavendish Laboratory, Free School Lane, Cambridge†

PHILOSOPHICAL MAGAZINE

VIII/7 1963