

N71-26011
NASA CR-115041

Louisiana State University in Shreveport

Final Technical Report on Grant NGR-19-012-001

Dr. Rex L. Matlock, Principal Investigator

The purpose of this work was to develop a model for the description of nucleon-nucleon and pion-nucleon collisions at high energy and to incorporate this model into a complete Monte-Carlo description of a Nuclear Cascade.

In the initial phase of the work an attempt was made to use the statical model originally proposed by Fermi¹. This attempt² was prompted by the success of the author (in an earlier work) to describe proton-antiproton annihilations at rest with a thermodynamic version of Fermi's model. The failure of this simple model to correctly describe the angular distribution of the particles produced in high energy nucleon-nucleon collisions led to its rejection.

The next phase of the work consisted of a study of a thermodynamic model developed by Hagedorn³. It was found that in the energy region where reliable experimental results are available that the Hagedorn model yields rather good agreement with experiment. The Hagedorn model was thus adopted as the basic model for the description of high energy particle collisions.

The effort in this research was then directed toward calculating, based on the Hagedorn model, the momentum and particle number distributions necessary for a Monte-Carlo description of a Nuclear Cascade. The required momentum distributions, for the case of proton-proton collisions, were readily available from Hagedorn's³ calculations. The momentum distributions for the pion-proton case were worked out by the author in collaboration with Dr. Hagedorn. It is worth noting here that the distributions for the pion-proton case needs further experimental verification in order to insure their validity. The particle number distributions, within the framework of the Hagedorn model, are quite a problem.

CASE FILE
COPY

This problem has not been satisfactorily resolved as of this writing. It turns out that, within the energy region where reliable experimental results are available, the particle number distributions are to a good approximation Poisson. One can then use "as a stop gap measure" the Poisson distribution together with the Hagedorn model momentum distributions to form a complete nuclear model for high energy particle collisions. The author is continuing work on the particle number distribution. When a satisfactory calculation is made, it will be made available to the Cosmic Ray Branch of NASA's Manned Space-Craft Center in Houston.

The details of how to incorporate the Hagedorn model momentum distributions into a Monte-Carlo description of a Nuclear Cascade program are supplied in the appendix to this report.

References:

1. E. Fermi, Progress of Theoretical Physics 5,570 (1950)
2. Ph. D. Dissertation by the Author (Unpublished)
3. R. Hagedorn and J. Ranft, Supplement to Nuovo Cimento 6,169 (1968)

Appendix to Technical Report on Grant

NGR-19-012-001

I. Momentum and angular distribution tables

The following momentum and angular distribution tables need to be calculated and saved. A description of how these tables are to be used is given in section 2 below.

$DMIJK(P,E)$ = Momentum distribution tabular value.

Where

I = 1 for nucleon (proton or neutron) table.

= 2 for π^+ and π^0 table.

= 3 for π^- table.

J = 1 for newly created particles.

= 2 for "Through going" particles.

K = 1 for "Thermodynamic contribution".

= 2 for "Isobar contribution".

= 3 for sum of "thermodynamic contribution" and "Isobar contribution".

E = Laboratory energy of the incident particle.

P = Magnitude of a secondary particle of type "I".

$DAIJK(\theta,E)$ = Angular distribution tabular value. Where I,J,K, and E are defined as above and θ = angle that the momentum of the secondary makes with the collision axis.

For definitions of the W(p) and Q(E) functions used below see section III of this appendix.

1. Through Going Protons (I=1, J=2)

$$DM_{123}(P, E) = 2\pi P^2 \Delta P \int_0^\pi W_p(\vec{P}) \sin\theta d\theta$$

$$DA_{123}(\theta, E) = 2\pi \sin\theta \Delta\theta \int_0^{P_{\max}} W_p(\vec{P}) P^2 dP$$

$P_{\max} = \gamma_0 \beta_0 M_p$; $M_p = 938$ Mev; See Section III for γ_0, β_0 values.

2. Newly created π^+ and π^0 (I=2, J=1)

$$DM_{211}(P, E) = 2\pi P^2 \Delta P \int_0^\pi \frac{W_{\pi^-}(\vec{P})}{Q_\pi(E)} \sin\theta d\theta$$

$$DM_{212}(P, E) = 2\pi P^2 \Delta P \int_0^\pi \left\{ \begin{array}{l} \text{Isobar decay term} \\ \text{see section III} \end{array} \right\} \sin\theta d\theta$$

$$DA211(\theta, E) = 2\pi \sin\theta \Delta\theta \int_0^{P_{\max}} \frac{W_{\pi^-}(\vec{P})}{Q_{\pi}(E)} P^2 dP$$

$P_{\max} = \gamma_0 \beta_0 M_{\pi}$; $M_{\pi} = 140$ Mev; See Section III for γ_0, β_0 values.

$$DA212(\theta, E) = 2\pi \sin\theta \Delta\theta \int_0^{P_{\max}} \left\{ \begin{array}{l} \text{isobar decay term} \\ \text{see section III} \end{array} \right\} P^2 dP$$

3. Newly Created π^- ($I=3, J=1$)

$$DM311(P, E) = 2\pi P^2 \Delta P \int_0^{\pi} \frac{W_{\pi^-}(\vec{P})}{Q_{\pi}(E)} \sin\theta d\theta$$

$$DA311(\theta, E) = 2\pi \sin\theta \Delta\theta \int_0^{P_{\max}} \frac{W_{\pi^-}(\vec{P})}{Q_{\pi}(E)} P^2 dP$$

$P_{\max} = \gamma_0 \beta_0 M_{\pi}$; $M_{\pi} = 140$ Mev; See Section III for γ_0, β_0 values.

4. Through Going π^+ and π^0 ($I=2, J=2$)

$$DM221(P, E) = 2\pi P^2 \Delta P \int_0^\pi W_{\pi^-}^*(\vec{P}) \sin\theta d\theta$$

$$DM222(P, E) = 2\pi P^2 \Delta P \int_0^\pi \left\{ \begin{array}{l} \text{Isobar decay term} \\ \text{see section III} \end{array} \right\} \sin\theta d\theta$$

$$DA221(\theta, E) = 2\pi \sin\theta \Delta\theta \int_0^{P_{\max}} W_{\pi^-}^*(\vec{P}) P^2 dP$$

$$DA222(\theta, E) = 2\pi \sin\theta \Delta\theta \int_0^{P_{\max}} \left\{ \begin{array}{l} \text{Isobar decay term} \\ \text{see section III} \end{array} \right\} P^2 dP$$

$P_{\max} = \gamma_0 \beta_0 M_\pi$; $M_\pi = 140$ Mev; See Section III for γ_0, β_0 values.

5. Through Going π^- ($I=3, J=2$)

$$DM321(P, E) = 2\pi P^2 \Delta P \int_0^\pi W_{\pi^-}^*(\vec{P}) \sin\theta d\theta$$

$$DA321(\theta, E) = 2\pi \sin\theta \Delta\theta \int_0^{P_{\max}} W_{\pi^-}^*(\vec{P}) P^2 dP$$

$P_{\max} = \gamma_0 \beta_0 M_\pi$; $M_\pi = 140$ Mev; See Section III for γ_0, β_0 values.

II. Prescription for picking momenta and angles from Distribution Tables

1. Pick an energy value, E , such that

(1) Distribution tables have been calculated for E

(2) $E_i \in [10^{-1/2}E, 10^{1/2}E)$ where E_i is the incident particles energy.

2. Through Going Nucleon case

(1) Pick a number X from the uniform distribution

$$f(x) = \begin{cases} 1/k & \text{for } 0 < x < k \\ 0 & \text{otherwise} \end{cases}$$

$$\text{Where } K = \sum_{p=0}^{P_{\text{max}}} DM123(p, E).$$

(2) Pick a momentum value, P , such that

$$X < \sum_{p=0}^P DM123(p, E) < \sum_{p=0}^{P+\Delta P} DM123(p, E)$$

(3) Pick a value, θ , such that

$$X < \sum_{\theta'=0}^{\theta} DA123(p, E) < \sum_{\theta'=0}^{\theta+\Delta\theta} DA123(p, E)$$

3. Newly created π^+ and π^0 case

(1) Pick a number X from the uniform distribution

$$f(x) = \begin{cases} 1 & \text{for } 0 < x < k \\ - & \\ k & \\ 0 & \text{otherwise} \end{cases} .$$

$$\text{Where } K = Q_{\pi}(E_i) \sum_{P=0}^{P_{\max}} \text{DM211}(p, E) + \sum_{P=0}^{P_{\max}} \text{DM212}(p, E) .$$

(2) Pick a momentum value, P , such that

$$X < Q_{\pi}(E_i) \sum_{P=0}^P \text{DM211}(p, E) + \sum_{P=0}^P \text{DM212}(p, E) < Q \sum_{P=0}^{P+\Delta P} + \sum_{P=0}^{P+\Delta P} .$$

(3) Pick a value, θ , such that

$$X < Q_{\pi}(E_i) \sum_{\theta'=0}^{\theta} \text{DA211}(p, E) + \sum_{\theta'=0}^{\theta} \text{DA212}(p, E) < Q \sum_{\theta'=0}^{\theta+\Delta\theta} + \sum_{\theta'=0}^{\theta+\Delta\theta} .$$

4. Newly created π^- case

(1) Pick a number X from the uniform distribution

$$f(x) = \begin{cases} 1 & \text{for } 0 < x < k \\ - & \\ k & \\ 0 & \text{otherwise} \end{cases} .$$

$$\text{Where } K = Q_{\pi}(E_i) \sum_{P=0}^{P_{\max}} \text{DM311}(p, E) .$$

(2) Pick a value, P , such that

$$X < Q_{\pi}(E_i) \sum_{P=0}^P \text{DM311}(p, E) < Q_{\pi}(E_i) \sum_{P=0}^{P+\Delta P} \text{DM311}(p, E) .$$

(3) Pick a value, θ , such that

$$X < Q_{\pi}(E_i) \sum_{\theta'=0}^{\theta} DA311(p,E) < Q_{\pi}(E_i) \sum_{\theta'=0}^{\theta+\Delta\theta'} DA311(p,E)$$

5. Through Going π^+ and π^0 case

(1) Pick a number X from the uniform distribution

$$f(x) = \begin{cases} \frac{1}{k} & \text{for } 0 < x < k \\ 0 & \text{otherwise} \end{cases}$$

$$\text{Where } k = \sum_{P=0}^{P_{\max}} [DM221(p,E) + DM222(p,E)].$$

(2) Pick a value, P , such that

$$X < \sum_{P=0}^P [DM221(p,E) + DM222(p,E)] < \sum_{P=0}^{P+\Delta P} [DM221(p,E) + DM222(p,E)]$$

(3) Pick a value, θ , such that

$$X < \sum_{\theta=0}^{\theta} [DA221(\theta,E) + DA222(\theta,E)] < \sum_{\theta=0}^{\theta+\Delta\theta'} [DA221(\theta,E) + DA222(\theta,E)]$$

6. Through Going π^- case

(1) Pick a number X from the uniform distribution

$$f(x) = \begin{cases} \frac{1}{k} & \text{for } 0 < x < k \\ 0 & \text{otherwise} \end{cases}$$

$$\text{Where } k = \sum_{P=0}^{P_{\max}} DM321(p,E)$$

(2) Pick a momentum value, P , such that

$$X < \sum_{P=0}^P \text{DM321}(p,E) < \sum_{P=0}^{P+\Delta P} \text{DM321}(p,E)$$

(3) Pick an angular value, θ , such that

$$X < \sum_{\theta'=0}^{\theta} \text{DA321}(\theta',E) < \sum_{\theta'=0}^{\theta+\Delta\theta} \text{DA321}(\theta',E)$$

III. Hagedorn Model Momentum Distributions

1. Newly created π^-

$$W_{\pi^-}(\vec{p}) = Q_{\pi^-}(E) \int_{-1}^1 \left\{ O \xrightarrow{\pi^-} \right\} d\lambda \quad .$$

$$O \xrightarrow{\pi^-} \equiv F(\lambda) L(\beta) \left\{ f_{\pi^-}(\xi', \lambda) \right\} \quad .$$

$$F(\lambda) \equiv (1 - |\lambda|) e^{-a|\lambda|} \left\{ \frac{1}{a^2} (a - 1 + e^{-a}) \right\}^{-1} \quad .$$

$$L(\beta) \left\{ f_{\pi^-}(\xi') \right\} = f_{\pi^-} \left[\gamma(\xi - \beta p_{||}) \right] \cdot \frac{\gamma(\xi - \beta p_{||})}{\xi} \quad .$$

$$a = 5.635 \quad . \quad \beta = \frac{\lambda}{|\lambda|} \left[\frac{1}{\gamma} \sqrt{\gamma^2 - 1} \right] \quad .$$

$$\gamma = 1 + (\gamma_0 - 1) |\lambda| \quad . \quad \gamma_0 = \sqrt{\frac{\gamma_L + 1}{2}} \quad . \quad \gamma_L = \frac{E_L}{M_p}$$

$$M_p = 938 \text{ Mev} \quad . \quad \xi = \sqrt{p^2 + m_{\pi}^2} \quad . \quad p_{||} = p \cos \theta \quad .$$

$$M_{\pi} = 140 \text{ Mev} \quad . \quad \theta \equiv \text{Direction of } \vec{p} \quad .$$

$$f_{\pi}(\xi', \lambda) = \begin{cases} \frac{V}{8\pi^3} [e^{\frac{\xi'}{T(\lambda)}} - 1]^{-1} & \text{for } \xi' \leq (\xi'_{\pi})_{\max} \\ 0 & \text{for } \xi' > (\xi'_{\pi})_{\max} \end{cases} .$$

$$T(\lambda) = \gamma T(\gamma_0, \gamma) = \gamma T[\xi(\gamma_0, \gamma)] .$$

$$\gamma = 1.027 \quad \xi(\gamma_0, \gamma) = \frac{M_p}{V} \left(\frac{\gamma_0}{\gamma} \right)$$

$$V = \gamma V_0 = \frac{4}{3} \pi (1.656) (M_{\pi})^{-3}$$

$$M_p = 938 \text{ Mev} \quad M_{\pi} = 140 \text{ Mev}$$

$T(\xi)$ will be supplied as a tabulated function. (See pg. 300 of reference 3)


$$(\xi')_{\max} = \frac{M_z^2 + M_{\pi}^2 - M_p^2}{2 M_z} .$$

$$M_z = \gamma E_{cm} - \sqrt{\gamma^2 E_{cm}^2 - E_{cm}^2 + M_p^2} .$$


$$E_{cm} = 2 \gamma_0 M_p$$

$$Q_{\pi}(E) = (0.848) E_L^{\frac{1}{4}} [\text{Gev}]^{\frac{1}{4}} \quad (\text{See page 33 of Hagedorn Model II}^3).$$

2. Newly created π^0 and π^+

$$W_{\pi^+}(\vec{P}) = W_{\pi^-}(\vec{P}) + \int_{-1}^1 \left[\text{Diagram} \right] d\lambda .$$


The second term is the isobar decay term.

$$\text{Diagram} \equiv \frac{F_0(\lambda)}{N_B(\lambda)} L(\beta) \sum_{N^*} A_{N^*} f_{\pi, N^*}^*(\xi', \lambda) .$$


$$F_0(\lambda) = \frac{(1-\lambda) \exp[-b\lambda] + \lambda d \exp[-c(1-\lambda)]}{\left(\frac{1}{b^2}\right)(b-1 + e^{-b}) + \left(\frac{b}{c^2}\right)(c-1 + e^{-c})}$$

$$b = 20.8 \quad , \quad c = 2.4 \quad , \quad d = 7.1$$

$$N_B(\lambda) = 4V \left(\frac{M_P T}{2\pi} \right)^{\frac{3}{2}} \exp\left[-\frac{M_P}{T}\right] \left\{ 1 + \frac{a_B}{4M_P^{3/2}} \exp\left[\frac{M_P}{T}\right] E_1\left(\frac{M^* T_0 - T}{T_0 T}\right) \right\} .$$

$$E_1(x) = \int_x^\infty t^{-1} e^{-t} dt \quad , \quad a_B = 2 \times 10^3 (\text{Mev})^{\frac{3}{2}} .$$

$$M^* = 1236 \text{ Mev}$$

$$T_0 = 130 \text{ Mev} .$$

$$f_{\pi^+, N^*}^*(\epsilon', \lambda) = \frac{Z_{N^*} V M^*}{16 \pi^3 \epsilon' p_{\pi^+}} T^2 \left\{ \left(1 + \frac{\epsilon^{(-)}}{T}\right) \exp\left[-\frac{\epsilon^{(-)}}{T}\right] - \left(1 + \frac{\epsilon^{(+)}}{T}\right) \exp\left[-\frac{\epsilon^{(+)}}{T}\right] \right\}.$$

$$\epsilon^{(\pm)} = \frac{M^*}{M_{\pi^+}^2} (\epsilon' \epsilon_{\pi^+} \pm p' p_{\pi^+})$$

$$p' = \sqrt{\epsilon'^2 - M_{\pi^+}^2}$$



$$p_{\pi^+} = \frac{1}{2M^*} \left\{ \left[M^{*2} - (M_{\pi^+} + M_j)^2 \right] \left[M^{*2} - (M_{\pi^+} - M_j)^2 \right] \right\}^{\frac{1}{2}}$$

$$\epsilon_{\pi^+} = \frac{1}{2M^*} (M^{*2} + M_{\pi^+}^2 - M_j^2)$$

Newly Created π Table

A_{N^*}	M^* (mev)	Z_{N^*}	M_j (mev)
57.3	1236	8	$M_p = 938.3$
1,733.	1525	4	$M_n = 939.6$
65,460.	2200	8	$M_n = 939.6$

3. "Through Going" Nucleons (proton and Neutron)

$$W_P(\vec{P}) = \int_{-1}^1 \left[\text{Diagram 1} + \text{Diagram 2} \right] d\lambda$$



$$\text{Diagram 1} \equiv \frac{F_0(\lambda)}{N_B(\lambda)} L(\beta) \left\{ f_{\kappa}(\xi', \lambda) \right\}$$

$$\text{Diagram 2} \equiv \frac{F_0(\lambda)}{N_B(\lambda)} L(\beta) \left\{ \sum_{\kappa^*} A_{\kappa^*} f_{P, \kappa^*}^*(\xi', \lambda) \right\}$$

$$F_0(\lambda) \equiv \text{See } \pi^+ \text{ case.}$$

$$N_B(\lambda) \equiv \text{See } \pi^+ \text{ case.}$$

$$f_{\kappa}(\varepsilon', \lambda) = \begin{cases} \frac{2V}{8\pi^3} \left[e^{\frac{\varepsilon'}{T}} + 1 \right]^{-1} & \text{for } \varepsilon' \leq (\varepsilon'_p)_{\text{max}} \\ 0 & \text{for } \varepsilon' > (\varepsilon'_p)_{\text{max}} \end{cases} .$$

$$(\varepsilon'_p)_{\text{max}} = \gamma (\gamma_0 M_p - \beta P_R)$$

See π^- case for a relation between γ and λ .

See π^- case for a relation between γ_0 and $\gamma_L = \frac{E_L}{M_p}$.

$$L(\beta) \left\{ f_{\kappa}(\varepsilon', \lambda) \right\} = \text{See } \pi^- \text{ case.}$$

Proton table

A_{N^*}	$M^* (\text{mev})$	Z_{N^*}	$M_j (\text{mev})$
57.3	1236	12	$M_{\pi} = 140$
1,733.	1525	8	$M_{\pi} = 140$
65,460.	2200	16	$M_{\pi} = 140$

$f_{p, N^*}(\varepsilon', \lambda)$ is the same as $2f_{\pi^+, N^*}(\varepsilon', \lambda)$ except $\pi^+ \rightarrow p$.

$F(\lambda) \equiv \text{See } \pi^- \text{ case.}$

$$N_F(\lambda) = Z_F V \left(\frac{M_F T(\lambda)}{2\pi} \right) \exp\left[-\frac{M_F}{T(\lambda)}\right] \left\{ 1 + \frac{a_F}{Z_F M_F^{3/2}} \exp\left[\frac{M_F}{T(\lambda)}\right] E_1\left(\frac{M_F(T_0 - T)}{T T_0}\right) \right\}$$

F	Z _F	M _F	a _F	M _F [*]
K	2	496 Mev	4 × 10 ³ Mev ^{3/2}	725 Mev
Y	6	1171 Mev	1.7 × 10 ³ Mev ^{3/2}	1382 Mev
B̄	4	940 Mev	2 × 10 ³ Mev ^{3/2}	1236 Mev
π	3	140 Mev	2 × 10 ³ Mev ^{3/2}	549 Mev

$$E_1(\lambda) \equiv \text{See } \pi^+ \text{ case.} \quad T(\lambda) \equiv \text{See } \pi^- \text{ case.} \quad T_0 = 160 \text{ Mev}$$

4. "Through Going" π^-

$$W_{\pi^-}^*(\vec{P}) = \int_0^1 \left[\text{Diagram: a circle with a right-pointing arrow labeled } \pi^- \text{ inside} \right] d\lambda$$


$$\text{Diagram: a circle with a right-pointing arrow labeled } \pi^- \text{ inside} \equiv \frac{F_0(\lambda)}{N_T(\lambda)} L(\beta) \left\{ f_{\pi^-}(\varepsilon; \lambda) \right\}.$$


$$F_0(\lambda) \equiv \text{See through going proton case.}$$

$$L(\beta) \left\{ f_{\pi^-}(\mathcal{E}', \lambda) \right\} \text{ See newly created } \pi^- \text{ case.}$$

$$N_{\pi}(\lambda) \equiv \text{See page (16) of this appendix.}$$

5. "Through Going" π^+ , π^0

$$W_{\pi^+}^*(\vec{P}) = W_{\pi^-}^*(\vec{P}) + \int_0^1 \left[\text{Diagram} \right] d\lambda.$$


$$\text{Diagram} \equiv \frac{F_0(\lambda)}{N_{\pi}(\lambda)} L(\beta) \left\{ \sum_{A_{\pi^*}} A_{\pi^*} f_{\pi, \pi^*}^*(\mathcal{E}', \lambda) \right\}.$$


$$f_{\pi, \pi^*}^*(\mathcal{E}', \lambda) = \frac{2Z_{\pi^*} V M^*}{16\pi^3 \mathcal{E}' \rho_{\pi}^0} \left\{ \left(1 + \frac{\mathcal{E}^{(-)}}{T}\right) \exp\left[-\frac{\mathcal{E}^{(-)}}{T}\right] - \left(1 + \frac{\mathcal{E}^{(+)}}{T}\right) \exp\left[-\frac{\mathcal{E}^{(+)}}{T}\right] \right\}.$$

$$E^{(\pm)} = \frac{M^*}{M_\pi^2} (E' E_\pi \pm P' P_\pi) \quad , \quad P' = \sqrt{E'^2 - M_\pi^2} \quad , \quad E_\pi = \frac{1}{2M^*} (M^{*2} + M_\pi^2 - M_j^2) \quad .$$

$$P_\pi = \frac{1}{2M^*} \left\{ \left[M^{*2} - (M_\pi + M_j)^2 \right] \left[M^{*2} - (M_\pi - M_j)^2 \right] \right\}^{\frac{1}{2}}$$

Through going π Table

A_{π^*}	$M^* (\text{Mev})$	Z_{π^*}	$M_j (\text{Mev})$
8.3×10^3	765	6	140
1.5×10^3	1069	1	140
1.5×10^2	1263	5	140