

CENTER FOR METEORITE STUDIES

ARIZONA STATE
UNIVERSITY
TEMPE • ARIZONA



GPO PRICE \$ _____

CFSTI PRICE(S) \$ _____

METEORITIC RUTILE
PETER B. BUSECK
and
KLAUS KEIL

Hard copy (HC) 1.00

April, 1966
Contribution No. 17

Microfiche (MF) 0.50

653 July 65

FACILITY FORM 002

N66 26487
(ACCESSION NUMBER)

11
(PAGES)

CR 95782
(NASA CR OR TMX OR AD NUMBER)

(THRU)

(CODE)

13
(CATEGORY)

METEORITIC RUTILE

Peter R. Buseck
Departments of Geology and Chemistry
Arizona State University
Tempe, Arizona

and

Klaus Keil
Space Sciences Division
National Aeronautics and Space Administration
Ames Research Center
Moffett Field, California

ABSTRACT

26687

Rutile has not been widely recognized as a meteoritic constituent. Recent microscopic and electron microprobe studies show, however, that TiO_2 is a reasonably widespread phase, albeit in minor amounts. X-ray diffraction studies confirm the TiO_2 to be rutile. It was observed in the following meteorites - Allegan, Bondoc, Estherville, Farmington, and Vaca Muerta. The rutile is associated primarily with ilmenite and chromite, in some cases as exsolution lamellae.

Rutile, as a meteoritic phase, is not widely known. In their summary of meteorite mineralogy neither Mason (1962) nor Ramdohr (1963) report rutile as a mineral occurring in meteorites, although Ramdohr did describe a similar phase from the Farmington meteorite in his list of "unidentified minerals." He suggested (correctly) that his "mineral D" might be rutile. He also observed it in several mesosiderites. The mineral was recently mentioned to occur in Vaca Muerta (Fleischer, et al., 1965) and in Odessa (El Goresy, 1965). We have found rutile in the meteorites Allegan, Bondoc, Estherville, Farmington, and Vaca Muerta; although nowhere an abundant phase, it appears to be rather widespread.

Of the several meteorites in which it was observed, rutile is the most abundant in the Farmington L-group chondrite. There it occurs in fine lamellae in ilmenite. The ilmenite is only sparsely distributed within the meteorite although wherever it does occur it is in moderately large clusters - up to 0.5 mm in diameter - and it then is usually associated with chromite as well as rutile (Buseck, et al., 1965).

Optically, the rutile has a faintly bluish tinge when viewed in reflected, plane-polarized light with immersion objectives. It is probably strongly anisotropic, although this is difficult to determine positively as it occurs within a matrix - ilmenite - that is itself so strongly anisotropic. Because of its small size, internal reflections could only be noted in rare instances. The reflectivity is low and the hardness approximately that of the ilmenite.

The initial identification of the rutile was performed by electron beam scans across the area of interest. In order to confirm the qualitative identification, quantitative electron microprobe analyses were run (Tables 1-3). The intensity readings were corrected for deadtime, background, mass absorption, fluorescence, and atomic number using methods previously described

(e.g., Keil and Andersen, 1965). The Ti content of the rutile was measured using a synthetic, spectrographically pure rutile standard.

The composition of the rutile was determined by moving the sample in 1-micron steps under a fixed electron beam. As the lamellae are very fine (on the order of a few microns in width) the electron beam is bound to overlap the host mineral ilmenite. Hence, the measured Ti content is slightly lower than expected (the ideal Ti content would be 59.9 percent). The measured value of 57.3 percent represents the highest value in a series of several hundred points. The small amount of iron which is detected when measuring the rutile lamellae is probably also a result of the overlapping. In view of this ambiguity it cannot be positively asserted that the rutile is either pure or stoichiometric. The ilmenite is fairly rich in Mg and should be called manganese-chromium-bearing magnesian ilmenite following the terminology suggested by Palache, et al., 1944. The chromite contains Ti as well as Mg and thus may be called titanium-manganese-bearing magnesian chromite. Terrestrial ilmenites and chromites commonly also contain Mg, there being solid solution series of geikielite (MgTiO_3) and magnesiochromite (MgCr_2O_4), respectively. The ilmenite and chromite were qualitatively scanned for all the elements with atomic numbers greater than that of Na; only those elements shown in Tables 1 and 2 were present in amounts greater than 0.05 weight percent.

The electron microprobe analysis clearly shows the mineral in question to be a titanium dioxide; this technique, however, cannot distinguish between the several polymorphs. Presumably because rutile is more stable under most geologic conditions than either anatase or brookite other investigators have assumed that meteoritic titanium dioxides consisted of rutile. No X-ray measurements have been reported. Unfortunately the small size of the TiO_2

TABLE 1. COMPOSITION OF MANGANESE-CHROMIUM-BEARING MAGNESIAN ILMENITE FROM THE FARMINGTON CHONDRITE AS OBTAINED BY ELECTRON MICROPROBE TECHNIQUES, AVERAGE OF 53 ANALYSES ON 5 GRAINS (IN WEIGHT PERCENT)

<u>Measured</u>	<u>Calculated</u>	<u>Ideal</u>
Fe = 25.8	FeO 33.19	47.34
Ti = 34.4	TiO ₂ 57.38	52.66
Mg = 4.11	MgO 6.82	
Mn = 0.59	MnO 0.76	
Cr = $\frac{0.51}{65.31}$	Cr ₂ O ₃ $\frac{0.75}{98.90}$	<u>100.00</u>

TABLE 2. COMPOSITION OF TITANIUM-MANGANESE-BEARING MAGNESIAN CHROMITE FROM THE FARMINGTON CHONDRITE AS OBTAINED BY ELECTRON MICROPROBE TECHNIQUES, AVERAGE OF 46 ANALYSES ON 5 GRAINS (IN WEIGHT PERCENT)

<u>Measured</u>	<u>Calculated</u>	<u>Ideal</u>
Fe = 23.8	FeO 30.62	32.09
Cr = 40.2	Cr ₂ O ₃ 58.75	67.91
Mg = 3.61	MgO 5.99	
Ti = 1.90	TiO ₂ 3.17	
Mn = $\frac{0.48}{69.99}$	MnO $\frac{0.62}{99.15}$	<u>100.00</u>

TABLE 3. COMPOSITION OF RUTILE FROM THE FARMINGTON CHONDRITE AS OBTAINED BY ELECTRON MICROPROBE TECHNIQUES (IN WEIGHT PERCENT)

<u>Measured</u>	<u>Calculated</u>	<u>Ideal</u>
Ti = 57.3	TiO ₂ 95.58*	100.00

*The difference to 100 percent is iron. The iron is probably not a constituent of the rutile but originates from the electron beam overlapping ilmenite.

lamellae and scarcity of the Farmington material precludes separation of the phase and thus X-ray diffraction measurements.

The presumptive rutile was also identified with the microprobe in the Bondoc meteorite. In an attempt to concentrate the titanium dioxide severely weathered outer crust of Bondoc was taken and treated first with HCl and then with $\text{HNO}_3 + \text{HF}$ and then again with HCl in order to dissolve all but the most resistant minerals. Further concentration of the very small amount of insoluble residue proved necessary to get a good X-ray pattern. Therefore, first magnetic splits and then specific gravity splits were made. The resulting material is pure rutile (Table 4).

The occurrence of rutile in the Farmington meteorite is interesting for not only are there rutile lamellae within the ilmenite, but chromite lamellae occur in the same area. Some of the chromite grains look like exsolution lamellae but others are continuous with adjacent and larger, granular chromite grains (Figures 2, 3). Rutile and chromite lamellae sometimes occur together in one and the same ilmenite grain as is shown in the center of Figure 3. This grain exhibits a rather peculiar relationship between the rutile and chromite lamellae. The rutile seems to crosscut the chromite and also seems to offset it in places. However, the edges of the host ilmenite grain do not show any marked physical displacements. Note also that the left chromite lamella seems to penetrate somewhat into the rutile lamella.

In most of the other meteorites, rutile is associated with chromite. In Estherville, a mesosiderite fall, chromite is widespread within the silicate areas. Ilmenite commonly occurs with the chromite, either in euhedral crystals or thin lamellae. The rutile is less abundant and it is restricted to occurring in long, thin lamellae within the chromite. These

TABLE 4

X-RAY DIFFRACTION PATTERN ($Fe_{K\alpha}$ RADIATION) OF POWDER HAVING A SPECIFIC GRAVITY GREATER THAN 3.33; TAKEN FROM THE ACID-INSOLUBLE, NONMAGNETIC FRACTION OF THE BONDOC METEORITE. COLUMNS 2, 3, AND 4 ARE FOR COMPARISON AND SHOW STANDARD PATTERNS OF RUTILE, ANATASE, AND BROOKITE, FROM THE A.S.T.M. X-RAY POWDER DATA FILE (A.S.T.M. SPEC. TECH. PUB. 48-J, PHILADELPHIA, 1960).

<u>Meteoritic Rutile (measured)</u>		<u>Rutile</u>			<u>Anatase</u>		<u>Brookite</u>	
<u>d</u>	<u>I/I₀</u>	<u>d</u>	<u>I/I₀</u>	<u>hkl</u>	<u>d</u>	<u>I/I₀</u>	<u>d</u>	<u>I/I₀</u>
-		-			3.51	100	-	
-		-			-		3.47	100
3.246	M*	3.245	100	110	-		-	
-		-			-		2.90	85
2.487	ST	2.489	41	101	-		2.48	50
-		-			2.435	9	-	
-		-			-		2.41	25
-		-			2.379	22	-	
-		-			2.336	9	-	
-		2.297	7	200	-		-	
-		-			-		2.21	25
2.184	W	2.188	22	111	-		-	
-		-			-		2.11	25
2.061	W	2.054	9	210	-		-	
-		-			-		1.97	50
-		-			1.891	33	1.88	75
-		-			-		1.84	50
-		-			1.699	21	-	
1.686	M	1.687	50	211	-		1.68	50
-		-			1.665	19	-	
-		-			-		1.65	60
1.624	W	1.624	16	220	-		-	
-		-			-		1.59	50
-		-			1.494	4	-	
1.478	W	1.480	8	002	1.480	13	1.48	25
1.459	W	1.453	6	310	-		-	
-		-			-		1.44	25
-		-			1.367	5	-	

*ST = strong, M = medium, W = weak

lamellae appear to be crystallographically oriented and presumably resulted by exsolution from chromite (Fig. 4). Its greyish-blue color and higher reflectivity than chromite or ilmenite serve to identify it, even though it is commonly at the limit of resolution at maximum magnification (1265X). It is reasonably widespread.

Rutile has recently been mentioned as occurring in Vaca Muerta (Fleischer, et al., 1965) as well as in other mesosiderites (Ramdohr, 1964). In Vaca Muerta - a mesosiderite find - the rutile occurs together with chromite, both in lamellae in the chromite as well as in individual grains. It is interesting to note that the rutile does not only occur in lamellae in the chromite, but also that chromite lamellae occur within large rutile grains (Fig. 5).


The chromite in Allegan - an H-group chondrite fall - contains 1.20 weight percent TiO_2 (Merrill and Stokes, 1900). Being moderately Ti-rich it was thought that it might also contain rutile. In fact, Allegan does contain rutile as a very minor accessory mineral, but it is disseminated as discrete grains within the silicate matrix, rather than as lamellae in chromite.

Bondoc is an extremely unusual meteorite (Nininger, 1963). It consists of fist-size, rounded pods of metal within a predominantly silicate matrix. The metal, consisting largely of kamacite, is in sizeable grains between which there are inclusions of nonmetallic minerals, mainly silicates and some apatite. The rutile occurs within these non-metallic segregations either together with chromite or as discrete small, angular grains not far away from it.

The mineralogical association in the vicinity of the rutile in meteorites is rather interesting as it includes kamacite, troilite, chromite, and sometimes ilmenite. Rutile has been seen in contact with ilmenite, chromite, and kamacite. Unfortunately, the regions of interest in the Fe-Ti-Cr-O system


have not been adequately described; otherwise this complex assemblage would very likely provide enough information to determine the pressure-temperature conditions at the time of formation.

In Farmington the rutile appears to have been derived from the ilmenite. This is commonly the case in terrestrial materials, although the reactions are slightly different. In the terrestrial case ilmenite breaks down when oxidized, first by oxidation of the ferrous to ferric iron, with concurrent addition of water and then decomposes further to yield goethite plus rutile, commonly as amorphous TiO_2 (Kukovskii and Kononov, 1959). The latter readily recrystallizes. In meteorites, having a relatively low f_{O_2} , this reaction is unlikely - reduction appears much more probable. The reaction



has been suggested for ilmenite breakdown (Ramdohr, 1964).

Experimental measurements indicate that ilmenite and rutile cannot coexist stably at very high temperatures as they react to form $FeTi_2O_5$, a mineral in the pseudobrookite solid solution series. On cooling, however, the $FeTi_2O_5$ decomposes to rutile plus ilmenite at $1140 \pm 10^\circ C$ (Lindsley, 1965),



clearly indicating that the assemblage observed in the Farmington meteorite formed below that temperature.

A limited amount of thermochemical data is available. In order to use the existing data it must be assumed that minor constituents in the minerals do not appreciably affect the thermodynamic properties and that the phases are stoichiometric (if they are not the calculations must be revised - in the case of rutile see, e.g., Moser, et al., 1965; Taylor and Schmalzried point out that stoichiometric ilmenite cannot coexist stably with Fe at $1300^\circ C$ - it

is inferred, however, that it is stoichiometric at lower temperatures). Thus, applying free energy values given by Taylor and Schmalzried (1964) to the reduction reaction (eq. 1) it can be shown that rutile will be the stable phase only for oxygen fugacities less than $10^{-21.1}$ atm and less than $10^{-14.6}$ atm at 800° and 1100°C , respectively, these being the temperature limits of the applicable thermodynamic data. These results are in agreement with calculations made from the virial coefficients given by Kubaschewski and Evans (1958). They are presumably not unreasonable for conditions within a meteorite body, although it must be emphasized that they apply with certainty to only very small areas within the meteorite. There are surely insufficient amounts of these minerals to buffer the entire meteorite mass.

Bulk chemical composition, including TiO_2 determination, are available for several of the meteorites in which rutile was found. The stony fraction of Allegan contains 0.10 weight percent TiO_2 (Merrill and Stokes, 1900), the Bondoc pyroxene contains 0.22 weight percent TiO_2 (Wiik, personal communication, 1965), and the Farmington meteorite contains 0.17 weight percent TiO_2 (Buseck, et al., 1965). These values may be compared to the average values of superior analyses of 53 stony meteorites, which is 0.11 weight percent TiO_2 (Urey and Craig, 1953). This agrees with the average value of 0.11 weight percent TiO_2 for 13 chondrites analyzed by Wiik and Wahl (Moore and Brown, 1961, Table 2) and for 86 chondrites analyzed spectrographically by Moore and Brown (1961). It is slightly lower than the mean of 0.15% given by Mason (1965). Thus, on the basis of limited compositional data, the rutile-bearing meteorites do not appear to be exceptionally Ti-rich. Presumably it is likely that more careful optical examination will show that rutile is present in many meteorites.

Although texturally much of the rutile appears to be the result of exsolution from ilmenite, it has been reported (e.g., Edwards, 1960), that based on experimental evidence, "ilmenite and rutile do not form solid solution to any marked degree." This is confirmed by recent unpublished measurements of Taylor (written communication to PRB, 1965). The fact that rutile lamellae seem to crosscut chromite lamellae within ilmenite of Farmington supports this contention, as do the small rutile grains in Bondoc, which occur isolated from either chromite or ilmenite. It appears likely that the rutile has several origins - a) reduction from ilmenite (Farmington), b) exsolution from chromite (Estherville), and c) primary crystallization as a minor accessory mineral (Allegan, Bondoc).

ACKNOWLEDGEMENT

We are grateful to Dr. H. Whitehurst for making the TiO_2 standard available to us, to Mr. J. Erlichman for his assistance in the electron microprobe work, to Mr. C. Lewis for sample preparations, to Dr. H. B. Wiik for help with the chemical separations, and to Dr. R. W. Taylor for making available unpublished data. This work was supported in part by NASA Grant NsG-399.

FIGURE TITLES


FIG. 1.- Electron beam scanning pictures of a rutile lamella (R) in ilmenite (Il), associated with chromite (Chr) and troilite (Tr) from the Farmington chondrite. The pictures were obtained by scanning the electron beam over the microscopically selected area in the sample and recording the signals from the counter on the cathode ray tube of an oscilloscope, the cathode ray beam scanning the tube in synchronism with the electron beam scanning the sample.

FIG. 2.- Farmington chondrite. Polycrystalline ilmenite (Il) showing various shades of gray due to bi-reflectance, with rutile lamellae (TiO_2), associated with chromite (Chr), nickel-iron and troilite (white). Black is silicate matrix.


FIG. 3.- Farmington chondrite. Lamellae of rutile (TiO_2) within ilmenite (Il), associated with chromite (Chr), troilite (Tr), and nickel-iron (NiFe). Black is silicate matrix. Chromite also occurs as lamellae in ilmenite. Note the peculiar relationship between the rutile and chromite lamellae occurring together in the ilmenite grain shown in the center of the figure.

FIG. 4.- Estherville mesosiderite. Chromite crystal (Chr) with rutile lamellae (TiO_2), associated with ilmenite (Il), in a silicate matrix. Oil immersion.

FIG. 5.- Vaca Muerta mesosiderite. Chromite (Chr) with lamellae of troilite (Tr) and of rutile (TiO_2), and rutile with lamellae of chromite, in contact with metallic nickel-iron (NiFe). Dark matrix is constituted of silicates. Oil immersion.




SKETCH OF GRAIN BOUNDARIES



Fe $\kappa\alpha$




Ti $\kappa\alpha$




Cr $\kappa\alpha$

10 μ


A-35975




A-35978



A-35977



A-35976



REFERENCES

- BUSECK, P. R., MASON, B. and WIIK, H. B. (1966) The Farmington meteorite - mineralogy and chemistry. Geochim. et Cosmochim. Acta, 30, 1-8.
- EDWARDS, A. B. (1960) Textures of the ore minerals and their significance. Aust. Inst. Min. and Met., Melbourne.
- EL GORESY, A. (1965) Mineralbestand und Strukturen der Graphit - und Sulfideinschlüsse in Eisenmeteoriten. Geochim. et Cosmochim. Acta, 29, 1131-1151.
- FLEISCHER, R. L., NAESER, C. W., PRICE, P. B., WALKER, R. M., and MARVIN, U. B. (1965) Fossil particle tracks and uranium distributions in minerals of the Vaca Muerta meteorite. Science, 148, 629-631.
- KEIL, K. and ANDERSEN, C. A. (1965) Electron microprobe study of the Jajh deh Kot Lalu enstatite chondrite. Geochim. et Cosmochim. Acta, 29, 621-632.
- KUBASCHEWSKI, O. and EVANS, E. LL. Metallurgical Thermochemistry. Pergamon Press, London, 1958.
- KUKOVSKII, E. G. and KONONOV, Yu. V. (1959) X-ray diffraction study of the products of rutilization of ilmenite. Rentgenogr. Mineral'n. Syr'ya, Gl. Upr. Geol. i Okhrany Nedr pri Sov. Min. Ukr. SSR, Tr. Pervogo Vses. Soveshch., Kiev, 1, 120-127 (CA, 59, 11105d).
- LINDSLEY, D. H. (1956) Lower thermal stability of FeTi_2O_5 - Fe_2TiO_5 (pseudobrookite) solid solution series (abst.) Geol. Soc. Am. Spec. Paper (in press).
- MASON, B. (1962) Meteorites. John Wiley and Sons, 274.
- MASON, B. (1965) The chemical composition of olivine-bronzite and olivine-hypersthene chondrites. Amer. Mus. Novitates, 2223, 38.

- MERRILL, G. P. and STOKES, H. N. (1900) A new stony meteorite from Allegan, Michigan. Proc. Washington Acad. Sciences, 2, 41-51.
- MOORE, C. B. and BROWN, H. (1962) The distribution of manganese and titanium in stony meteorites. Geochim. et Cosmochim. Acta, 26, 495-502.
- MOSER, J. B., BLUMENTHAL, R. N. and WHITMORE, D. H. (1965) Thermodynamic study of nonstoichiometric rutile (TiO_{2-x}). J. Am. Cer. Soc., 48, 384.
- NININGER, H. H. (1963) Meteorite with unique features. Science, 139, 345-347.
- PALACHE, C., BERMAN, H. and FRONDEL, C. E. (1944) Dana's system of mineralogy. John Wiley and Sons, Inc.
- RAMDOHR, P. (1963) The opaque minerals in stony meteorites. J. Geophys. Res., 68, 2011-2036.
- RAMDOHR, P. (1964) Opaque minerals in stony meteorites. Carnegie Institution, Annual Report of the Director of the Geophysical Laboratory.
- TAYLOR, R. W., and SCHMALZRIED, H. (1964) The free energy of formation of some titanates, silicates and magnesium aluminate from measurements made with galvanic cells involving solid electrolytes. J. Phys. Chem., 68, 2444-2449.
- UREY, H. C. and CRAIG, H. (1953) The composition of the stone meteorites and the origin of the meteorites. Geochim. et Cosmochim. Acta, 4, 36-82.