

RESEARCH

Open Access



Ecuzumab treatment: stochastic occurrence of C3 binding to individual PNH erythrocytes

Michela Sica¹, Tommaso Rondelli¹, Patrizia Ricci², Maria De Angioletti^{1,3}, Antonio M. Risitano² and Rosario Notaro^{1*} 

Abstract

Background: C5 blockade by ecuzumab prevents complement-mediated intravascular hemolysis in paroxysmal nocturnal hemoglobinuria (PNH). However, C3-bound PNH red blood cells (RBCs), arising in almost all treated patients, may undergo extravascular hemolysis reducing clinical benefits. Despite the uniform deficiency of CD55 and of CD59, there are always two distinct populations of PNH RBCs, with (C3+) and without (C3-) C3 binding.

Methods: To investigate this paradox, the phenomenon has been modeled in vitro by incubating RBCs from ecuzumab untreated PNH patients with compatible sera containing ecuzumab, and by assessing the C3 binding after activation of complement alternative pathway.

Results: When RBCs from untreated patients were exposed in vitro to activated complement in the context of C5-blockade, there was the prompt appearance of a distinct C3+ PNH RBC population whose size increased with time and also with the rate of complement activation. Eventually, all PNH RBCs become C3+ to the same extent, without differences between old and young (reticulocytes) PNH RBCs.

Conclusions: This study indicates that the distinct (C3+ and C3-) PNH RBC populations are not intrinsically different; rather, they result from a stochastic *all-or-nothing* phenomenon linked to the time-dependent cumulative probability of each individual PNH red cell to be exposed to levels of complement activation able to trigger C3 binding. These findings may envision novel approaches to reduce C3 opsonization and the subsequent extravascular hemolysis in PNH patients on ecuzumab.

Keywords: Paroxysmal nocturnal hemoglobinuria, Complement blockade, Intravascular hemolysis, Extravascular hemolysis, Complement C3, Complement C5

Background

Paroxysmal nocturnal hemoglobinuria (PNH) is an acquired disorder of hematopoiesis characterized by a somatic mutation in the *PIGA* gene that prevents or impairs the synthesis of glycosylphosphatidylinositol (GPI) anchors [1, 2]. The deficiency on red blood cells (RBCs) of GPI-anchored proteins [3, 4], including the complement regulators CD55 [5, 6] and CD59 [7], results in

chronic intravascular hemolysis with recurrent exacerbations, anemia, smooth muscle cell dystonia, and high risk of thrombosis [4, 8–10].

The blockade of terminal complement pathway by ecuzumab [11], a monoclonal antibody (moAb) against complement component 5 (C5), abrogates intravascular hemolysis with the consequent normalization of lactate dehydrogenase (LDH) levels in almost all patients suffering from PNH. This treatment has proven to be safe and clinically effective in hemolytic PNH patients [12–14], except those in which bone marrow failure is the major cause of anemia [15, 16].

The persistence (or the recurrence) of intravascular hemolysis is observed only in few conditions: (i) Japanese patients carrying a rare polymorphism of C5 [17],

* Correspondence: rosario.notaro@ittumori.it

Preliminary results of this work have been presented at the XV Meeting of the European Hematology Association (Barcelona, Spain; 2010) and at the XI Congress of the Italian Society of Experimental Hematology (Torino, Italy; 2010).

¹Laboratory of Cancer Genetics and Gene Transfer, Core Research Laboratory – Istituto Toscano Tumori (CRL-ITT), AOU Careggi, viale Pieraccini 6, 50139 Florence, Italy

Full list of author information is available at the end of the article

(ii) patients with an increased eculizumab turnover requiring extra-dosage (pharmacokinetic breakthrough) [18], and (iii) patients who occasionally experience transient episodes of intravascular hemolysis because of massive complement activation during infections or inflammatory disorders (pharmacodynamic breakthrough) [18–21]. Despite these small and infrequent limits, the treatment with eculizumab has radically changed the natural history of PNH since in most patients it reduces anemia [12, 13] and thrombosis [22], and improves quality of life and survival [14].

However, the abrogation of intravascular hemolysis is not the only relevant change in PNH pathophysiology associated with eculizumab treatment. In fact, at variance with PNH patients not treated with eculizumab, a population of GPI-negative (PNH) RBCs bound with fragment of complement component 3 (C3) appears in almost all patients on eculizumab [23] and, in some patients, also the less-sensitive direct antiglobulin test may turn positive [24]. The PNH RBCs bound with C3 become apparent because PNH RBCs, spared from hemolysis by the blockade of the terminal complement cascade, remain unable to control the early steps of the ongoing complement activation. Eventually, these PNH RBCs, once opsonized with complement, become potential targets of phagocytosis by macrophages, with consequent variable degrees of extravascular hemolysis [23]. Accordingly, in PNH patients on eculizumab, the extent of C3 binding correlates with reticulocyte count, the in vivo half-life of ^{51}Cr -labeled RBCs is reduced and there is an excess of spleen and liver ^{51}Cr uptake [23]. This extravascular hemolysis is clinically relevant because it can limit the efficacy of eculizumab to the point that some patients may remain transfusion-dependent [14, 22–25]. The heterogeneity of mechanisms controlling C3 binding and/or removal of C3+ PNH RBCs are likely to account for the variable extent of C3 binding and of the consequent extravascular hemolysis. Part of this variability may result from genetic diversity in genes coding for endogenous regulators of complement: indeed, we have recently shown that polymorphisms of complement receptor 1 (*CR1*) gene are associated with the transfusion need of patients on eculizumab [26].

The most intriguing biological feature of C3 binding is that two distinct populations of PNH RBCs are always present in patients on eculizumab, one with (C3+) and one without (C3-) C3 binding, despite the uniform deficiency of the GPI-anchored proteins on all PNH RBCs [23]. The understanding of this phenomenon is of more than academic importance because, depending on the answer, one might explore different ways to overcome the consequent clinical problems. In this paper, by reproducing in vitro C3 binding in the presence of C5 blockade, we provide evidence suggesting a stochastic

model for the emergence of these two distinct, C3+ and C3-, populations of PNH RBC.

Methods

Patients and samples

Peripheral blood samples were collected from 31 healthy donors and from 31 patients with frank hemolytic PNH. Nine of these patients were on eculizumab, 13 were not and 9 have been studied before and on eculizumab. Signed informed consent was obtained according to an IRB-approved protocol. RBCs and sera were promptly separated from freshly collected samples. RBCs were washed three times in saline just before use. Sera were stored at $-80\text{ }^{\circ}\text{C}$.

In vitro complement activation

The approach we have used derives from the diagnostic Ham test traditionally used to test the susceptibility of PNH RBCs to complement-mediated lysis by activation of the complement alternative pathway (CAP) [27–29]. Briefly, 2% RBC suspensions were incubated at $37\text{ }^{\circ}\text{C}$ in saline with 84% of pooled ABO-compatible sera from either healthy donors (normal) or from PNH patients on eculizumab (with eculizumab). Sera with eculizumab were collected 1 h after infusion from PNH patients stably treated with eculizumab (900 mg each 2 weeks) and with normal LDH during the last 3 months (in these samples the expected concentration of eculizumab is between 200 and 500 $\mu\text{g/ml}$: [12, 30] a concentration that is 5–14 times higher than the 35 $\mu\text{g/ml}$ reported as the minimal effective level of eculizumab [12]. In selected experiments, eculizumab, obtained from remnants in the infusion lines, was added (400 $\mu\text{g/ml}$).

CAP was activated by either spontaneous activation (incubation in sealed tubes with 100% atmospheric air) [31] or by mild acidification (HCl 0.016 M: the final pH was between 6.8 and 6.5) [31–34]. All the procedures were performed in sterile conditions. The hemolysis of normal RBCs has been quantified by a direct spectrophotometric method [35]. In both experimental models, the lysis of RBCs from healthy donors was less than 2%. The hemolysis of PNH RBCs has been quantified by comparing the relative variation of the percentage of normal RBCs (GPI-positive resistant to complement) with that of PNH RBCs (GPI-negative sensitive to complement) [36].

Flow cytometry

C3 binding was assessed with anti-C3d-neoantigen (A250, Quidel, USA) secondary stained with polyclonal rabbit-anti-mouse antibodies (Dako Cytomation, Denmark). In selected experiments, RBCs were stained also with anti-C3b (H11, Serotec), anti-iC3b (A209, Quidel), anti-C3c (MCA2605, Serotec), anti-Bb (A227, Quidel), and the

proper secondary polyclonal rabbit-anti-mouse antibodies (Dako Cytomation). PNH RBCs have been identified with anti-CD59 (Mem43, Serotec, UK). Intact and lysed (ghost) RBCs have been identified by physical parameters as previously described [37]; in selected experiments, the identity of intact and of ghost RBCs has been confirmed by staining with anti-Glycophorin A moAb (GA-R2, BD Becton Dickinson, NJ, USA).

Reticulocytes have been stained with the polymethine dye (Ret Search II, Sysmex) that, upon excitation at 633 nm, emits between 650 and 670 nm. Briefly, 1 volume of RBC suspension was mixed in 4 volumes of "Ret Search" dilution buffer (Sysmex) and stained by adding the polymethine dye just before flow cytometry analysis [38].

Flow cytometry has been performed with either FACS-Canto (BD) or Accuri C6 (BD). Fluorescent-activated cell sorting (FACS) has been performed with FACSARIA III (BD).

Statistical analysis

Statistical analysis was performed using non-parametric tests: Wilcoxon signed paired and Friedman rank sum tests as suitable. Statistical significance was accepted for any $P < 0.05$.

Results

Analysis of C3 fragments bound to PNH red blood cells in patients on eculizumab

In most of the PNH patients undergoing eculizumab treatment, there is a relative and an absolute increase of PNH RBCs because complement blockade prevents their complement-mediated lysis. In parallel, the percentage of PNH RBCs with C3 binding increases progressively from the time of starting eculizumab treatment (a representative patient in Fig. 1a), until a relatively stable plateau is reached after 10–12 weeks: $5.8 \pm 3.4\%$ at 1 week ($n = 8$), $16.9 \pm 13.3\%$ at 4–5 weeks ($n = 8$), $20.5 \pm 6.6\%$ at 10–




Fig. 1 In vivo C3 binding on red cells of PNH patients on eculizumab. **a** Kinetics of C3 binding on red cells from a representative PNH patient during eculizumab treatment. Before treatment (0 week), no red cell binds C3, whereas starting from 1 week of treatment three populations of red cells are displayed: one population of normal red cells (CD59 + C3-) and two distinct populations of PNH (CD59-negative) red cells, one with (C3+) and one without (C3-) fragments bound on their surface. **b** Characterization of C3 fragment bound to erythrocytes from PNH patient on eculizumab. The *left panel* shows the normal (CD59-positive) and PNH (CD59-negative) populations (gated by the *elliptic mark*). The PNH (CD59-negative) population (see gate in the *left panel*) has been analyzed with anti-C3d and either anti-C3b (*middle panel*) or anti-iC3b (*left panel*). SSC: side scatter

12 weeks ($n = 8$). In a series of 41 PNH patients on stable eculizumab treatment, we have previously observed $27.2 \pm 18.7\%$ of C3+ PNH RBCs [23]. PNH RBCs with bound C3 were stained by a moAb that recognizes a specific C3d neo-epitope [39], but not by an anti-C3b (unable to bind the C3d fragment) and an anti-iC3b (Fig. 1b); in addition, these C3 bound PNH RBCs were not stained by anti-Bb and anti-C3c antibodies (Additional file 1: Figure S1).

Modeling in vitro the in vivo binding of C3 to PNH red blood cells

In order to investigate what happens in vivo in PNH patients on eculizumab, we have designed a novel experimental setting [31], whereby RBCs from PNH patients are exposed to the mild activation of complement that mimics the in vivo spontaneous activation of CAP known as C3 “tickover”. C3 “tickover” occurs because of the spontaneous continuous and low-rate hydrolysis of C3, followed by its interaction with Factor B, which eventually leads to the formation of the initial fluid phase C3 convertase (C3:H₂OBB) [39–44].

When we incubated RBCs from a PNH patient in normal sera at 37 °C, we observed after 5 days a significant degree of hemolysis [36]. The ghosts arising from hemolysis were exclusively GPI-negative (ghosts of PNH RBCs), and nearly all of them bound C3 fragments (Additional file 1: Figure S2). In contrast, none of the intact RBCs have bound C3 fragments confirming that we are mimicking the continuous chronic intravascular hemolysis [32, 37] occurring in vivo in PNH patients [8, 45].

When we repeated these experiments using sera with eculizumab, there was almost no lysis of PNH RBCs (0–10% at day 5) and we observed, already after 24 h, the appearance of a discrete population of C3+ PNH RBCs coexisting with PNH RBCs without C3 and with normal RBCs (Fig. 2a). In addition, the size of the C3+ PNH RBC population increased significantly with time ($P < 0.04$) up to $29 \pm 23\%$ after 5 days (Fig. 2b). This parallels exactly what happens in vivo in patients on eculizumab (Fig. 1a). No C3 binding was observed with sera that were previously heat-inactivated (data not shown), confirming that the phenomenon was entirely complement-mediated.




Fig. 2 Effect of spontaneous complement activation on PNH red cells. **a** Kinetics of C3 binding on PNH red cells from a representative PNH patient not on eculizumab after spontaneous complement activation in presence of eculizumab. The *first panel* shows that no C3 binding is present on red cells from patients not on eculizumab. The *second panel* shows that there is no C3 binding on red cells after 5 days of incubation at 37 °C in ABO-compatible sera without eculizumab. C3 binding (C3+) appears on PNH (CD59-negative) red cells after 3 (*third panel*) and after 5 (*fourth panel*) days of incubation at 37 °C in ABO-compatible sera with eculizumab. The number in the *upper right angle* of each diagram indicates the time in days (d); 0d indicates the sample analyzed before any in vitro treatment. ECU: eculizumab. **b** Kinetics of C3 binding on PNH red cells from a series of PNH patients not on eculizumab after spontaneous complement activation in presence of eculizumab. The bar diagram shows the average (+SD) proportion of PNH (CD59-negative) red cells bound with C3 in a series of patients not on eculizumab after 3 and after 5 days of incubation at 37 °C in ABO-compatible sera with eculizumab. *n*: number of different patients studied at the indicated time points. *Empty bars* show the experiments without addition of MgCl₂ whereas the *filled bars* show the experiments with the addition of MgCl₂ (1.25 mM). C3 binding on PNH red cells increased significantly along the time (Wilcoxon signed paired test: $P < 0.04$ for the series without MgCl₂ and $P < 0.05$ for the series with addition MgCl₂). At both 3 and 5 days C3 binding of samples with MgCl₂ was significantly higher than that of the same samples without MgCl₂: $P < 0.008$, Wilcoxon signed paired test

Then we have carried out similar experiments in which CAP was activated by acidification (as in the Ham test) [27, 28, 32]. When RBCs from patients not on eculizumab were incubated with acidified normal sera, there was the lysis of almost all PNH RBCs after 2 h (Fig. 3a). When we used sera with eculizumab, the hemolysis of PNH RBCs was largely reduced but not completely prevented ($42.6 \pm 12.6\%$ after 24 h; $n = 11$). The level of this “residual” hemolysis was only slightly and not significantly reduced by adding an excess of eculizumab (400 $\mu\text{g}/\text{ml}$) that is expected to double its final concentration: 47.1 ± 12.0 vs. $43.0 \pm 10.0\%$ ($n = 5$).

In addition, when complement was activated by acidification in the presence of eculizumab, we observed the prompt appearance of a discrete population of C3+ PNH RBCs within 5 min from serum acidification ($5.9 \pm 3.6\%$) and its size increased with time up to $71 \pm 17\%$ within 24 h (Fig. 3a, b; $P < 0.001$). Again, no C3 binding was observed with sera that were previously heat-inactivated. With the addition of 1.25 mM MgCl_2 (which is known to optimize complement activation [46]), there was a further significant increase of C3+ PNH RBCs in both experimental settings (Figs. 2b, 3b) and also an increase of residual hemolysis ($66.7 \pm 14.7\%$ after 24 h; $n = 11$).

Furthermore, when RBCs from patients on eculizumab, who already had both populations of C3+ and C3- PNH RBCs, were incubated with their own acidified sera, nearly all the C3- PNH RBCs eventually became C3+ (Fig. 4a, b). This held true also for the C3- RBCs physically sorted from these patients: in fact, upon complement activation, also these sorted C3- PNH RBCs became C3+ (Fig. 4c).

C3 binding to young (reticulocytes) and to mature PNH red blood cells

Since in vivo not all PNH RBCs are bound with C3, one may surmise that this could be due to intrinsic difference between the PNH RBCs that become C3+ and those that remain C3-. For instance, it could be that young RBCs have some extra mechanism for complement regulation, which would inhibit C3 binding and might be lost with aging: i.e., the surface density of glycophorin A [47], CD44, CD147 [48], and CR1 [49] decreases with RBC aging. In order to test this hypothesis, we have compared the proportion of C3+ cells among young (reticulocytes) and mature (non-reticulocytes) PNH RBCs. In samples obtained ex vivo from patients on eculizumab ($n = 18$), the proportion of C3+ PNH RBCs was much lower in reticulocytes




Fig. 3 Effect of complement activation by mild acidification on PNH red cells. **a** Kinetics of C3 binding on PNH red cells from a representative untreated PNH patient after complement activation in vitro by mild acidification in presence of eculizumab. Red cells from a patient not on eculizumab incubated with acidified ABO-compatible sera without eculizumab show almost complete lysis of PNH (CD59-negative) red cells by 2 h: compare the first panel (0 h: before any acidification) and the second panel (2 h, no ECU). C3 binding (C3+) on PNH (CD59-negative) red cells is shown before (time 0) and after 5 min, 2 h and 24 h from complement activation in ABO-compatible sera with eculizumab. **b** Kinetics of C3 binding on PNH red cells from a series of PNH patients not on eculizumab after complement activation in vitro by mild acidification in presence of eculizumab. The line diagram shows the average (+SD) proportion of PNH (CD59-negative) red cells bound with C3 in a series of untreated patients at different time points (from 5 min to 24 h) since complement activation in ABO-compatible sera with eculizumab (continuous line without MgCl_2 ; dotted line with 1.25 mM MgCl_2). n : number of different patients studied at the indicated time points. C3 binding on PNH red cells increased significantly along the time (Friedman rank sum test: $P < 0.001$ for both series with and without MgCl_2). At any time C3 binding of samples with MgCl_2 was significantly higher than that of the same samples without MgCl_2 (Wilcoxon signed paired test: $P < 0.03$ at 5 min and 1 h; $P < 0.005$ at 30 min and 2, 6, and 24 h)




Fig. 4 Effect of complement activation by mild acidification on PNH red cells from patients on eculizumab. **a** C3 binding on PNH red cells from a representative PNH patient on eculizumab after complement activation in vitro by mild acidification in presence of eculizumab. This experiment shows the C3 binding (C3+) on PNH (CD59-negative) red cells before (*left panel*) and after 6 h (*right panel*) from complement activation in vitro. The number in the *upper right angle* of each diagram indicates the time in hours (h); 0 h indicates the sample analyzed before complement activation in vitro. **b** Kinetics of C3 binding on PNH red cells from a series of PNH patients on eculizumab after complement activation in vitro by mild acidification in presence of eculizumab. The line diagram shows the average (+SD) proportion of C3-negative PNH red cells that became bound with C3 in a series of eculizumab-treated patients at different time points (from 1 to 24 h) since complement activation in their own sera (*continuous line* without MgCl₂; *dotted line* with 1.25 mM MgCl₂). In this 7 patients, the percentage PNH red cells that in vivo were already bound with C3 was 30.3 ± 24.5%. *n*: number of different patients studied at the indicated time points. **c** Effect of in vitro complement activation on C3–PNH red cells from patients on eculizumab. RBCs without C3 binding have been sorted by FACS from a patient on eculizumab and subjected to complement activation in the presence of eculizumab. Analysis of RBC with anti-CD59 and anti-C3d: just after sample drawing (*left panel*); after FACS sorting of C3–RBC (*middle panel*); sorted C3–RBC after 2 h from complement activation in presence of eculizumab (*right panel*). HCl: serum acidification. ECU: eculizumab


than in non-reticulocytes (10.5% vs. 27.7%; $P < 0.0025$. Fig. 5a). Reticulocytes mature into RBCs within few hours also in vitro [50]; thus, we have studied C3 binding only after complement activation by serum acidification. In apparent contrast with what we have observed in vivo in patients on eculizumab, in untreated PNH patients ($n = 14$) a similar proportion of C3+ PNH RBCs was observed among both reticulocytes and non-reticulocytes (Fig. 5b) after in vitro incubation with acidified sera with eculizumab (Fig. 5c: 45.6 vs. 37.1%).

We have obtained direct confirmation of this scenario in 6 patients, which were tested in vitro before starting eculizumab (C3+ PNH reticulocytes and non-reticulocytes were $59 \pm 21\%$ and $53 \pm 25\%$, respectively), and re-tested in vivo after 4–8 months of eculizumab treatment (C3+ PNH reticulocytes and non-reticulocytes were $8 \pm 5\%$ and $23 \pm 15\%$, respectively). In all these experiments, C3 binding has never been observed on non-PNH (GPI-positive) reticulocytes (data not shown).

Discussion

The binding of C3 fragments to RBCs, characteristic of various autoimmune hemolytic anemias, had never been found in PNH and it has emerged as a novel phenomenon in PNH patients on eculizumab [23, 24, 51]. It is possible to visualize that in the absence of eculizumab no C3+ RBCs are seen because, once C3 is activated on the surface of PNH RBCs, they will promptly undergo hemolysis. In contrast, when C5 is blocked by eculizumab, PNH RBCs with bound C3 survive because they are no longer lysed by the terminal complement pathway; however, there will be now a “new” steady state in which part of these C3-opsonized RBCs are removed by macrophages, likely via interaction with the complement receptor 3 [52] producing the extravascular hemolysis [23].

This “new” steady state does not detract from the clinical benefit arising from the abrogation of intravascular hemolysis. However, this observation accounts for the fact that only a minority of PNH patients on eculizumab normalizes their hemoglobin level, and in some patients, residual anemia may even



require red blood cell transfusions [14, 15, 20, 23, 24]. For these patients there are not yet standard treatments: steroids have been proven to be ineffective [51]; splenectomy [53] or splenic artery embolization [54, 55] may be effective but raise concerns about infection and thrombosis susceptibility [15].

We have effectively reproduced in vitro the phenomenon of C3 binding: in fact, in vitro complement activation in the context of C5 blockade generates a distinct population of C3+ PNH RBCs which co-exists with C3- PNH RBCs. It is intriguing that spontaneous complement activation in vitro, a condition that may mimic the chronic low level of complement activation present in vivo, results in the same order of C3 binding we have previously observed in vivo in PNH patients ($n = 41$) on eculizumab: 29 vs. 27% [23].

In addition, this mild spontaneous complement activation is almost unable to lyse PNH RBCs (less than 10% after 5 days), whereas serum acidification results in a non-negligible hemolysis of PNH RBCs (~42% after 24 h) that is not prevented by an excess of eculizumab. This is because serum acidification forces a level of complement activation in excess of what happens in vivo in the steady state [37, 56] and that might be similar to the massive complement activation present during infection or inflammation. This may explain the hemolytic crisis observed in vivo in PNH patients on eculizumab in these specific clinical circumstances, providing evidence for a “pharmacodynamic breakthrough” [57]. Our observations [31, 58] are in keeping with the recent finding that strong complement activation overrides C5 inhibition by eculizumab possibly due to the generation of high density C3 products on the RBC surface [59].

In any event, the most singular feature of C3 binding in PNH patients on eculizumab is that, despite all PNH RBCs are uniformly lacking GPI-anchored proteins, there are always two distinct PNH RBC populations with or without C3 binding. This is different from other conditions, such as the paradigmatic cold agglutinin disease [60], in which C3 binding is present on all RBCs. Thus, C3 binding in PNH patients on eculizumab emerges as a unique phenomenon with surprising features.

In principle, these two distinct populations might arise from (a) their different ability to bind C3 or (b) from a stochastic effect. However, when C5 is blocked by eculizumab, there was the prompt appearance of a discrete population of C3+ PNH RBCs whose size increased not only with time but also with the level of complement activation. In fact, in vitro the proportion of C3+ PNH RBCs after mild spontaneous complement activation (Fig. 2b) was smaller than after more intense complement activation by acidification (Fig. 3b). Moreover, in vitro, at variance with in vivo observations, eventually all PNH RBCs become C3+. This holds true also for C3- PNH RBCs from patients on eculizumab that, in vitro, become all C3+, regardless of the size of C3+ PNH RBCs population in vivo (Fig. 4b). Finally, young RBCs are not selectively protected from C3 binding, in fact in vitro a very similar proportion of young (reticulocytes) and mature (non-reticulocytes) PNH RBCs become C3+ (Fig. 5b, c).

Altogether, these results indicate that the C3+ and the C3- PNH RBCs are not intrinsically different: any PNH RBCs, when exposed to complement activation in the context of C5 blockade, may bind C3 to the same extent. Thus, C3 binding is not a prerogative of a discrete subset of PNH RBCs. In addition, the prompt appearance, within 5 min from serum acidification (Fig. 3a), of two distinct populations of PNH RBCs (with and without C3) suggests that C3 binding is an *all-or-nothing* phenomenon in which a detectable level of C3 fragments stably bound to PNH

RBCs is generated only when complement activation exceed a minimal threshold.

These findings support a stochastic model in which the longer each individual RBC circulates, the higher the probability to be exposed in specific districts of the bloodstream to levels of complement activation that exceeds the threshold able to trigger C3 binding: for this reason in patients on eculizumab in vivo, the percentage of C3+ cells is much lower in reticulocytes (10.5%) than in mature RBCs (27.7%).

On the other hand, this stochastic model may also suggest that complement activation, rather than be always systemic and generalized, may also occur within localized spaces and within a limited time frame. Indeed, different organs might harbor specific mechanisms of activation and regulation of complement. This finely regulated homeostasis could explain why in most complement-mediated diseases the clinical presentations and complications are often organ specific (e.g., abdominal or central nervous vein thrombosis in PNH, renal or ocular involvement in genetically determined hemolytic-uremic syndrome and age-related macular degeneration) [61].

Conclusions

This in vitro study helps in understanding what happens in vivo in PNH patients on eculizumab and it paves the way for alternative strategies to overcome the clinical consequences of C3 binding in PNH patients who already benefit, completely or partially, from C5 blockade [34, 57]. For example, since the stochastic nature and the *all-or-nothing* characteristic of this process, it is possible to hypothesize that in the context of C5 blockade, the prevention of C3 binding would not require a complete inhibition/full blockade of C3 convertase activity. Indeed, even a partial inhibition of C3 convertase activity, just below the threshold that triggers irreversible C3 binding, might result in meaningful clinical benefit with less concerns about the possible infectious risk associated with a complete disabling of the complement pathway.

Additional file

Additional file 1: Figure S1. In vivo C3 binding on red cells of PNH patients on eculizumab. **Figure S2.** Effect of spontaneous complement activation on PNH red cells. (PDF 1927 kb)

Abbreviations

C3: Complement component 3; C5: Complement component 5; CAP: Complement alternative pathway; *CR1*: Complement receptor 1; GPI: Glycosylphosphatidylinositol; LDH: Lactate dehydrogenase; PNH: Paroxysmal nocturnal hemoglobinuria; RBC(s): Red blood cell(s)

Acknowledgements

We thank Alessandra Fanelli and Roberto Caporale for helpful technical advises. We thank Luca Boni for statistical analysis advises. We thank Lucio Luzzatto for much support, for the helpful suggestions, and the fruitful discussions. Last but

not least, we thank the patients, their families, and the "Associazione Italiana Emoglobinuria Parossistica Notturna" for their warm support.

Funding

This study has been funded in part by a start-up grant from the "Istituto Toscano Tumori" (RN).

Availability of data and materials

All data generated and analyzed during this study are included in this published article and its Additional file.

Authors' contributions

MS participated in designing and performing the research, in analyzing data and writing the paper; TR, PR, and MD participated in performing the research; AMR and RN designed the research, analyzed data and wrote the paper. All the authors critically revised the paper and contributed to the preparation of the final version. All authors read and approved the final manuscript.

Competing interests

AMR has served as consultant for and has received research funding and lecture fee from Alexion Pharmaceuticals; AMR has also received research funding from Alnylam, Rapharma, and Novartis. The remaining authors declare that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

The study was reviewed and approved by the Ethics Committees of AOU Careggi (Firenze, Italy) and of Federico II University (Naples, Italy). It has been performed according the Declaration of Helsinki on ethical principles for medical research involving human subjects. A written signed informed consent was obtained from all the subjects involved in the study.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Laboratory of Cancer Genetics and Gene Transfer, Core Research Laboratory – Istituto Toscano Tumori (CRL-ITT), AOU Careggi, viale Pieraccini 6, 50139 Florence, Italy. ²Hematology, Department of Clinical Medicine and Surgery, Federico II University, Naples, Italy. ³CCOM-CNR, Sesto Fiorentino, Florence, Italy.

Received: 10 April 2017 Accepted: 12 June 2017

Published online: 19 June 2017

References

1. Takeda J, Miyata T, Kawagoe K, Iida Y, Endo Y, Fujita T, Takahashi M, Kitani T, Kinoshita T. Deficiency of the GPI anchor caused by a somatic mutation of the PIG-A gene in paroxysmal nocturnal hemoglobinuria. *Cell*. 1993;73:703–11.
2. Bessler M, Mason PJ, Hillmen P, Miyata T, Yamada N, Takeda J, Luzzatto L, Kinoshita T. Paroxysmal nocturnal haemoglobinuria (PNH) is caused by somatic mutations in the PIG-A gene. *EMBO J*. 1994;13:110–7.
3. van der Schoot CE, Huizinga TW, van't Veer-Korthof ET, Wijmans R, Pinkster J, von dem Borne AE. Deficiency of glycosyl-phosphatidylinositol-linked membrane glycoproteins of leukocytes in paroxysmal nocturnal hemoglobinuria, description of a new diagnostic cytofluorometric assay. *Blood*. 1990;76:1853–9.
4. Luzzatto L, Notaro R. Paroxysmal nocturnal hemoglobinuria. In: Handin RL, Lux SE, Stossel TP, editors. *Blood, Principles and Practice of Hematology*. 2nd ed. Philadelphia: Lippincott Williams & Wilkins; 2003. p. 319–34.
5. Nicholson-Weller A, March JP, Rosenfeld SI, Austen KF. Affected erythrocytes of patients with paroxysmal nocturnal hemoglobinuria are deficient in the complement regulatory protein, decay accelerating factor. *Proc Natl Acad Sci U S A*. 1983;80:5066–70.
6. Pangburn MK, Schreiber RD, Muller-Eberhard HJ. Deficiency of an erythrocyte membrane protein with complement regulatory activity in paroxysmal nocturnal hemoglobinuria. *Proc Natl Acad Sci U S A*. 1983;80:5430–4.

7. Holguin MH, Fredrick LR, Bernshaw NJ, Wilcox LA, Parker CJ. Isolation and characterization of a membrane protein from normal human erythrocytes that inhibits reactive lysis of the erythrocytes of paroxysmal nocturnal hemoglobinuria. *J Clin Invest.* 1989;84:7–17.
8. Dacie JV. Paroxysmal nocturnal haemoglobinuria. In: Dacie JV, editor. *The haemolytic anaemias*, vol. 2. London: Churchill; 1967. p. 1128–260.
9. Rosse WF, Nishimura J. Clinical manifestations of paroxysmal nocturnal hemoglobinuria: present state and future problems. *Int J Hematol.* 2003;77: 113–20.
10. de Latour RP, Mary JY, Salanoubat C, Terriou L, Etienne G, Mohty M, Roth S, de Guibert S, Maury S, Cahn JY, Socie G. Paroxysmal nocturnal hemoglobinuria: natural history of disease subcategories. *Blood.* 2008;112:3099–106.
11. Rother RP, Rollins SA, Mojcik CF, Brodsky RA, Bell L. Discovery and development of the complement inhibitor eculizumab for the treatment of paroxysmal nocturnal hemoglobinuria. *Nat Biotechnol.* 2007;25:1256–64.
12. Hillmen P, Hall C, Marsh JC, Elebute M, Bombara MP, Petro BE, Cullen MJ, Richards SJ, Rollins SA, Mojcik CF, Rother RP. Effect of eculizumab on hemolysis and transfusion requirements in patients with paroxysmal nocturnal hemoglobinuria. *N Engl J Med.* 2004;350:552–9.
13. Hillmen P, Young NS, Schubert J, Brodsky RA, Socie G, Muus P, Roth A, Szer J, Elebute MO, Nakamura R, et al. The complement inhibitor eculizumab in paroxysmal nocturnal hemoglobinuria. *N Engl J Med.* 2006;355:1233–43.
14. Kelly RJ, Hill A, Arnold LM, Brooksbank GL, Richards SJ, Cullen M, Mitchell LD, Cohen DR, Gregory WM, Hillmen P. Long-term treatment with eculizumab in paroxysmal nocturnal hemoglobinuria: sustained efficacy and improved survival. *Blood.* 2011;117:6786–92. doi:10.1182/blood-2011-02-333997.
15. Luzzatto L, Gianfaldoni G, Notaro R. Management of paroxysmal nocturnal haemoglobinuria: a personal view. *Br J Haematol.* 2011;153:709–20. doi:10.1111/j.1365-2141.2011.08690.x.
16. Notaro R, Risitano AM. Extravascular Hemolysis After Eculizumab Treatment. In: *Paroxysmal Nocturnal Hemoglobinuria: From Bench to Bedside*, edited by Yuzuru Kanakura, Taroh Kinoshita, Jun ichi Nishimura. Springer, Japan KK, 2017. pp 283–295. doi:10.1007/978-4-431-56003-6_17.
17. Nishimura J, Yamamoto M, Hayashi S, Ohyashiki K, Ando K, Brodsky AL, Noji H, Kitamura K, Eto T, Takahashi T, et al. Genetic variants in C5 and poor response to eculizumab. *N Engl J Med.* 2014;370:632–9. doi:10.1056/NEJMoa1311084.
18. Hill A, Hillmen P, Richards SJ, Elebute D, Marsh JC, Chan J, Mojcik CF, Rother RP. Sustained response and long-term safety of eculizumab in paroxysmal nocturnal hemoglobinuria. *Blood.* 2005;106:2559–65. doi:10.1182/blood-2005-02-0564.
19. Roth A, Hock C, Konik A, Christoph S, Duhrsen U. Chronic treatment of paroxysmal nocturnal hemoglobinuria patients with eculizumab: safety, efficacy, and unexpected laboratory phenomena. *Int J Hematol.* 2011;93: 704–14. doi:10.1007/s12185-011-0867-y.
20. Olutogun T, Cutini I, Notaro R, Luzzatto L. Complement-mediated haemolysis and the role of blood transfusion in paroxysmal nocturnal haemoglobinuria. *Blood Transfus.* 2015;13:363–9. doi:10.2450/2015.0249-14.
21. Loschi M, Porcher R, Barraco F, Terriou L, Mohty M, de Guibert S, Mahe B, Lemal R, Dumas PY, Etienne G, et al. Impact of eculizumab treatment on paroxysmal nocturnal hemoglobinuria: a treatment versus no-treatment study. *Am J Hematol.* 2016;91:366–70. doi:10.1002/ajh.24278.
22. Hillmen P, Muus P, Duhrsen U, Risitano AM, Schubert J, Luzzatto L, Schrezenmeier H, Szer J, Brodsky RA, Hill A, et al. Effect of the complement inhibitor eculizumab on thromboembolism in patients with paroxysmal nocturnal hemoglobinuria. *Blood.* 2007;110:4123–8. doi:10.1182/blood-2007-06-095646.
23. Risitano AM, Notaro R, Marando L, Serio B, Ranaldi D, Seneca E, Ricci P, Alfinito F, Camera A, Gianfaldoni G, et al. Complement fraction 3 binding on erythrocytes as additional mechanism of disease in paroxysmal nocturnal hemoglobinuria patients treated by eculizumab. *Blood.* 2009;113:4094–100. doi:10.1182/blood-2008-11-189944.
24. Hill A, Rother RP, Arnold L, Kelly R, Cullen MJ, Richards SJ, Hillmen P. Eculizumab prevents intravascular hemolysis in patients with paroxysmal nocturnal hemoglobinuria and unmasks low-level extravascular hemolysis occurring through C3 opsonization. *Haematologica.* 2010;95:567–73. doi:10.3324/haematol.2009.007229.
25. Luzzatto L, Risitano AM, Notaro R. Paroxysmal nocturnal hemoglobinuria and eculizumab. *Haematologica.* 2010;95:523–6. doi:10.3324/haematol.2009.017848.
26. Rondelli T, Risitano AM, Peffault de Latour R, Sica M, Peruzzi B, Ricci P, Barcellini W, Iori AP, Boschetti C, Valle V, et al. Polymorphism of the complement receptor 1 gene correlates with the hematologic response to eculizumab in patients with paroxysmal nocturnal hemoglobinuria. *Haematologica.* 2014;99:262–6. doi:10.3324/haematol.2013.090001.
27. Ham TH. Chronic hemolytic anemia with paroxysmal nocturnal hemoglobinuria. A study of the mechanism of hemolysis in relation to acid-base equilibrium. *N Engl J Med.* 1937;217:915–8.
28. Dacie JV, Israels MCG, Wilkinson JF. Paroxysmal nocturnal haemoglobinuria of the Marchiafava type. *Lancet.* 1938;i:479–82.
29. Rosse WF. Dr Ham's test revisited. *Blood.* 1991;78:547–50.
30. Reiss UM, Schwartz J, Sakamoto KM, Puthenveetil G, Ogawa M, Bedrosian CL, Ware RE. Efficacy and safety of eculizumab in children and adolescents with paroxysmal nocturnal hemoglobinuria. *Pediatr Blood Cancer.* 2014;61: 1544–50. doi:10.1002/pbc.25068.
31. Sica M, Pascariello C, Rondelli T, Risitano AM, Notaro R. Kinetics of complement protein 3 (C3) binding to PNH (GPI-negative) erythrocytes under complement blockade by eculizumab. *Haematologica.* 2010;95:538.
32. Logue GL, Rosse WF, Adams JP. Mechanisms of immune lysis of red blood cells in vitro. I. Paroxysmal nocturnal hemoglobinuria cells. *J Clin Invest.* 1973;52:1129–37. doi:10.1172/JCI107279.
33. Holt DS, Botto M, Bygrave AE, Hanna SM, Walport MJ, Morgan BP. Targeted deletion of the CD59 gene causes spontaneous intravascular hemolysis and hemoglobinuria. *Blood.* 2001;98:442–9.
34. Risitano AM, Notaro R, Pascariello C, Sica M, del Vecchio L, Horvath CJ, Fridkis-Hareli M, Selleri C, Lindorfer MA, Taylor RP, et al. The complement receptor 2/factor H fusion protein TT30 protects paroxysmal nocturnal hemoglobinuria erythrocytes from complement-mediated hemolysis and C3 fragment. *Blood.* 2012;119:6307–16. doi:10.1182/blood-2011-12-398792.
35. Harboe M. A method for determination of hemoglobin in plasma by near-ultraviolet spectrophotometry. *Scand J Clin Lab Invest.* 1959;11:66–70. doi:10.3109/00365515909060410.
36. Ferreira VP, Pangburn MK. Factor H mediated cell surface protection from complement is critical for the survival of PNH erythrocytes. *Blood.* 2007;110: 2190–2.
37. Lindorfer MA, Pawluczko AW, Peek EM, Hickman K, Taylor RP, Parker CJ. A novel approach to preventing the hemolysis of paroxysmal nocturnal hemoglobinuria: both complement-mediated cytolysis and C3 deposition are blocked by a monoclonal antibody specific for the alternative pathway of complement. *Blood.* 2010;115:2283–91. doi:10.1182/blood-2009-09-244285.
38. Kono M, Kondo T, Takagi Y, Wada A, Fujimoto K. Morphological definition of CD71 positive reticulocytes by various staining techniques and electron microscopy compared to reticulocytes detected by an automated hematology analyzer. *Clin Chim Acta.* 2009;404:105–10. doi:10.1016/j.cca.2009.03.017.
39. Tamerius JD, Pangburn MK, Muller-Eberhard HJ. Detection of a neoantigen on human C3b and C3d by monoclonal antibody. *J Immunol.* 1985;135:2015–9.
40. Lachmann PJ, Halbwachs L. The influence of C3b inactivator (KAF) concentration on the ability of serum to support complement activation. *Clin Exp Immunol.* 1975;21:109–14.
41. Pangburn MK, Schreiber RD, Muller-Eberhard HJ. Formation of the initial C3 convertase of the alternative complement pathway. Acquisition of C3b-like activities by spontaneous hydrolysis of the putative thioester in native C3. *J Exp Med.* 1981;154:856–67.
42. Atkinson JP, Farries T. Separation of self from non-self in the complement system. *Immunol Today.* 1987;8:212–5. doi:10.1016/0167-5699(87)90167-8.
43. Lachmann PJ. The amplification loop of the complement pathways. *Adv Immunol.* 2009;104:115–49. doi:10.1016/S0065-2776(08)04004-2.
44. Nilsson B, Nilsson Ekdahl K. The tick-over theory revisited: is C3 a contact-activated protein? *Immunobiology.* 2012;217:1106–10. doi:10.1016/j.imbio.2012.07.008.
45. Marchiafava E. Anemia emolitica cronica con emosiderinuria perpetua. *Policlinico Med.* 1928;35:105–17.
46. May JE, Rosse W, Frank MM. Paroxysmal nocturnal hemoglobinuria. Alternate-complement-pathway-mediated lysis induced by magnesium. *N Engl J Med.* 1973;289:705–9. doi:10.1056/NEJM197310042891401.
47. Seaman GV, Knox RJ, Nordt FJ, Regan DH. Red cell aging. I. Surface charge density and sialic acid content of density-fractionated human erythrocytes. *Blood.* 1977;50:1001–11.
48. Sparrow RL, Healey G, Patton KA, Veale MF. Red blood cell age determines the impact of storage and leukocyte burden on cell adhesion molecules, glycophorin A and the release of annexin V. *Transfus Apher Sci.* 2006;34:15–23. doi:10.1016/j.transci.2005.09.006.

49. Lach-Trifilieff E, Marfurt J, Schwarz S, Sadallah S, Schifferli JA. Complement receptor 1 (CD35) on human reticulocytes: normal expression in systemic lupus erythematosus and HIV-infected patients. *J Immunol.* 1999;162:7549–54.
50. Gronowicz G, Swift H, Steck TL. Maturation of the reticulocyte in vitro. *J Cell Sci.* 1984;71:177–97.
51. Risitano AM, Notaro R, Luzzatto L, Hill A, Kelly R, Hillmen P. Paroxysmal nocturnal hemoglobinuria–hemolysis before and after eculizumab. *N Engl J Med.* 2010;363:2270–2. doi:10.1056/NEJMc1010351.
52. Lin Z, Schmidt CQ, Koutsogiannaki S, Ricci P, Risitano AM, Lambris JD, Ricklin D. Complement C3dg-mediated erythrophagocytosis: implications for paroxysmal nocturnal hemoglobinuria. *Blood.* 2015;126:891–4. doi:10.1182/blood-2015-02-625871.
53. Risitano AM, Marando L, Seneca E, Rotoli B. Hemoglobin normalization after splenectomy in a paroxysmal nocturnal hemoglobinuria patient treated by eculizumab. *Blood.* 2008;112:449–51. doi:10.1182/blood-2008-04-151613.
54. Krishnan SK, Hill A, Hillmen P, Arnold LM, Brooksbank GL, Wood A, Scarsbrook A, Davies MH, Kelly RJ. Improving cytopenia with splenic artery embolization in a patient with paroxysmal nocturnal hemoglobinuria on eculizumab. *Int J Hematol.* 2013;98:716–8. doi:10.1007/s12185-013-1454-1.
55. Araten DJ, Iori AP, Brown K, Torelli GF, Barberi W, Natalino F, De Propriis MS, Girmenia C, Salvatori FM, Zelig O, et al. Selective splenic artery embolization for the treatment of thrombocytopenia and hypersplenism in paroxysmal nocturnal hemoglobinuria. *J Hematol Oncol.* 2014;7:27. doi:10.1186/1756-8722-7-27.
56. Fishelson Z, Horstmann RD, Muller-Eberhard HJ. Regulation of the alternative pathway of complement by pH. *J Immunol.* 1987;138:3392–5.
57. Risitano AM, Ricklin D, Huang Y, Reis ES, Chen H, Ricci P, Lin Z, Pascariello C, Raia M, Sica M, et al. Peptide inhibitors of C3 activation as a novel strategy of complement inhibition for the treatment of paroxysmal nocturnal hemoglobinuria. *Blood.* 2014;123:2094–101. doi:10.1182/blood-2013-11-536573.
58. Sica M, Pascariello C, Rondelli T, Risitano AM, Notaro R. In vitro complement protein 5 (C5) blockade recapitulates the complement protein 3 (C3) binding to GPI-negative erythrocytes observed in paroxysmal nocturnal hemoglobinuria (PNH) patients on eculizumab. *Haematologica.* 2010;95:196.
59. Harder MJ, Kuhn N, Schrezenmeier H, Hochsmann B, von Zabern I, Weinstock C, Simmet T, Ricklin D, Lambris JD, Skerra A, et al. Incomplete inhibition by eculizumab: mechanistic evidence for residual C5 activity during strong complement activation. *Blood.* 2017;129:970–80. doi:10.1182/blood-2016-08-732800.
60. Berentsen S, Beiske K, Tjonnfjord GE. Primary chronic cold agglutinin disease: an update on pathogenesis, clinical features and therapy. *Hematology.* 2007;12:361–70. doi:10.1080/10245330701445392.
61. Holers VM. The spectrum of complement alternative pathway-mediated diseases. *Immunol Rev.* 2008;223:300–16. doi:10.1111/j.1600-065X.2008.00641.x.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

