

SCIENTIFIC REPORTS



OPEN

The multifunctional polydnavirus TnBVANK1 protein: impact on host apoptotic pathway

Rosanna Salvia¹, Gerarda Grossi¹, Angela Amoresano², Carmen Scieuzo¹, Marisa Nardiello¹, Chiara Giangrande², Ilaria Laurenzana³, Vitalba Ruggieri³, Sabino A. Bufo¹, S. Bradleigh Vinson⁴, Monica Carmosino¹, David Neunemann⁵, Heiko Vogel⁵, Pietro Pucci² & Patrizia Falabella¹

Toxoneuron nigriceps (Hymenoptera, Braconidae) is an endophagous parasitoid of the larval stages of the tobacco budworm, *Heliothis virescens* (Lepidoptera, Noctuidae). The bracovirus associated with this wasp (*TnBV*) is currently being studied. Several genes expressed in parasitised host larvae have been isolated and their possible roles partly elucidated. *TnBVank1* encodes an ankyrin motif protein similar to insect and mammalian I κ B, an inhibitor of the transcription nuclear factor κ B (NF- κ B). Here we show that, when *TnBVank1* was stably expressed in polyclonal *Drosophila* S2 cells, apoptosis is induced. Furthermore, we observed the same effects in haemocytes of *H. virescens* larvae, after *TnBVank1* *in vivo* transient transfection, and in haemocytes of parasitised larvae. Coimmunoprecipitation experiments showed that *TnBVANK1* binds to ALG-2 interacting protein X (Alix/AIP1), an interactor of apoptosis-linked gene protein 2 (ALG-2). Using double-immunofluorescence labeling, we observed the potential colocalization of *TnBVANK1* and Alix proteins in the cytoplasm of polyclonal S2 cells. When Alix was silenced by RNA interference, *TnBVANK1* was no longer able to cause apoptosis in both S2 cells and *H. virescens* haemocytes. Collectively, these results indicate that *TnBVANK1* induces apoptosis by interacting with Alix, suggesting a role of *TnBVANK1* in the suppression of host immune response observed after parasitisation by *T. nigriceps*.

Endophagous parasitoids belonging to the order of Hymenoptera develop during juvenile stages in the body of their hosts. During oviposition, the endoparasitoid female injects into the host several factors that alter its physiology, suppressing the host immune system, the endocrine balance, as well as reproductive activity or metabolism¹. Among parasitoid secretions, viruses belonging to the family of *Polydnaviridae* (PDVs) play a key role in the success of parasitism^{2,3}. PDVs are obligate symbionts of ichneumoid and braconid wasps attacking exclusively larval stages of their lepidopteran hosts⁴. Polydnavirus DNA is integrated into the genome of the parasitoid and therefore vertically transmitted through the germline⁴. The viral particles, containing circular dsDNA molecules of different sizes, replicate only in the epithelium of parasitoid female ovarian calyx, and are injected into the host, along with ovarian proteins, venom and the egg, during oviposition. PDVs express genes in several target tissues of the host and, together with other maternal or embryonic factors, lead to several functional alterations of their host, providing a suitable nutritional environment to the parasitoid larvae⁵. *Toxoneuron nigriceps* (Viereck) (Hymenoptera, Braconidae) is an endoparasitoid wasp belonging to the Braconidae family, parasitising the larval stages of tobacco budworm *Heliothis virescens* (Fabricius) (Lepidoptera, Noctuidae). *Toxoneuron nigriceps* Bracovirus (*TnBV*) encodes several gene families^{6,7} including ankyrin motif proteins (ANKs)⁸. The *TnBVANK* proteins show high levels of amino acid identity to the ankyrin motif of the *Drosophila* NF- κ B inhibitor (I κ B)-related protein cactus^{8,9}. Three ankyrin-like open reading frames (ORFs) were identified in the *TnBV* genome and were denoted *TnBVank1-3*⁸. *TnBVank* genes are characterised by ankyrin motif sequence repeats but

¹Department of Sciences, University of Basilicata, Via dell'Ateneo Lucano 10, 85100, Potenza, Italy. ²Department of Chemical Sciences, Federico II University, Via Cintia 6, 80126, Naples, Italy. ³Laboratory of Preclinical and Translational Research, IRCCS Referral Cancer Center of Basilicata (CROB), Via Padre Pio 1, 85028, Rionero in Vulture (PZ), Italy. ⁴Department of Entomology, Texas A&M University, 370 Olsen Blvd, College Station, TX, 77843-2475, USA. ⁵Department of Entomology, Max Planck Institute for Chemical Ecology, Hans-Knöll-Strasse 8, D-07745, Jena, Germany. Correspondence and requests for materials should be addressed to P.F. (email: patrizia.falabella@unibas.it)

the usually phosphorylatable serine residues at the N-terminus are absent⁸. Moreover, none of putative proteins encoded by the *TnBVank* genes has the C-terminal PEST domain present in the protein Cactus/I κ B, which is necessary for signal-mediated degradation^{10–12} and for the control of protein turnover¹³. The predicted structure of these proteins suggested that *TnBV* ankyrin proteins may bind NF- κ B/Rel transcription factors of the tumour necrosis factor (TNF)/Toll immune pathway altering the signal transduction cascade⁸. Indeed, it was demonstrated that in parasitised *H. virescens* larvae, after bacterial challenge, the nuclear import of NF- κ B was inhibited⁸. Moreover, transfection experiments in human HeLa cells demonstrated that a *TnBVank1* gene product reduced the efficiency of the TNF- α -induced expression of a reporter gene under NF- κ B transcriptional control, confirming that viral I κ B-like proteins are involved in the suppression of the insect immune response^{8,14,15}.

TnBVank1 and *TnBVank3* are expressed as early as three hours and six hours, respectively, after parasitisation in haemocytes of *H. virescens* last instar larvae, but the transcripts are present also several hours after parasitisation reducing over the time. Although expression of *TnBVank1* and *TnBVank3* is high in *H. virescens* haemocytes, a much more limited expression level can also be found in the fat body⁸. Moreover, it is reported that several hours after natural parasitism, the granulocytes, a subpopulation of haemocytes, showed a number of structural damages, as evidenced by actin cytoskeleton disruption and loss of adhesion properties, with general morphological changes which suggest the occurrence of apoptosis¹⁶. This further suggests a possible role of parasitism in early immune response suppression.

In insects the processes of cell death are involved in development, but also play an important role in allowing, maintaining and regulating many host-parasitoid interactions. Accordingly, it was shown that parasitoids are able to induce or regulate cell death phenomena in host tissues^{16–20}. Here, we report that *TnBVANK1* induces cell death in polyclonal *Drosophila* S2 cells that stably express *TnBVank1*. To better understand the role of *TnBVANK1*, coimmunoprecipitation and coimmunolocalisation experiments were carried out in polyclonal *Drosophila* S2 cells and the interaction between *TnBVANK1* and ALG-2 interacting protein X (Alix) was investigated. Alix is a multifunctional protein, it is ubiquitously expressed and concentrated in phagosomes and exosomes and it was first characterised as an interactor of apoptosis-linked gene protein 2 (ALG-2) that is a Ca²⁺-binding protein necessary for cell death²¹. Here, we show that upon Alix silencing by RNA interference (RNAi), *TnBVANK1* was no longer able to induce apoptosis in S2 cells.

These data were further confirmed by *in vivo* transient transfection of *TnBVank1* and by silencing of Alix transcript by RNAi in *H. virescens* larval haemocytes.

Results

Expression of *TnBVank1* in S2 cells induces apoptosis. We analysed the profile of polyclonal S2 cells stably expressing *TnBVank1* (S2[ank1]) by flow cytometry. We used S2 cells and polyclonal S2[ank1] cells 20 h after treatment with the cytotoxic alkaloid camptothecin as positive controls and untransfected S2 cells as negative control. All the cell samples were harvested at the same splitting passages and were doubly stained with Annexin V-FITC and PI (Propidium Iodide). Results revealed a significant increase of apoptotic cells in polyclonal S2[ank1] cells line compared to the negative control, indeed the percentage of apoptotic cells, amounted to 39.56%, was not significantly different from the positive controls (S2 + CAM and S2[ank1] + CAM) (Fig. 1a).

To further investigate the involvement of *TnBVank1* in cell death pathway, we analysed also the caspase-3 activity. Caspase activity detection is recognised as a standard method for assessing apoptosis²². Caspase-3 is activated early during apoptosis. We assayed its activity in polyclonal S2 cells stably expressing *TnBVank1* by using the fluorogenic effector caspase substrate Ac-DEVD-ACM. Cell lysates were harvested from S2 cells stably expressing *TnBVank1*, untransfected S2 cells (negative control), as well as both the positive controls. Caspase-3 activity was significantly increased in S2[ank1] cells compared to the negative control S2 but comparable to positive controls (S2 + CAM and S2[ank1] + CAM) (Fig. 1b). This result suggests that *TnBVank1* stimulates the activation of caspase-3 indicating a role of *TnBVANK1* in induction of apoptosis in S2 cells.

Moreover, the nuclear morphology, examined by staining with Hoechst 33258, showed intact nuclei appearing light blue in colour in the negative control (S2 cells) while polyclonal S2 cells exhibited bright blue indicating condensed or fragmented nuclei, typical morphological characteristics of apoptosis (see Supplementary Fig. S1).

Western blot time course. To test whether the production of *TnBVANK1* decreases over time in the stable polyclonal cell line, as a consequence of apoptotic cell death, a western blot time-course analysis was performed on cell lysates harvested at subsequent splitting passages (2, 4, 10, 20, 26, 39 passages). Western blot analysis showed an effective reduction of *TnBVank1* gene expression over time, indicating that the population of cells, stably expressing *TnBVank1* within the stable polyclonal cell line, progressively decreases with a concomitant increase of cell number not expressing *TnBVank1* (Fig. 2). This result strongly suggest that the expression of the viral gene *TnBVank1* induces cell death in S2 cells.

***In vivo* transient expression of *TnBVank1* efficiency in *H. virescens* haemocytes.** Haemocytes exhibited transient expression of *TnBVank1* when the recombinant vector pIZT/V5-His-ANK1, mixed with a transfection reagent, was injected into the haemocoel of fifth instar *H. virescens* larva (day one). After trying different transfection reagents, and different time after injection, we selected Lipofectin as the most efficient one and 56 h as the best time for *in vivo* transient expression, at least for our experimental conditions. The expression of *TnBVank1* transcripts in transfected haemocytes was detected by reverse transcriptase-polymerase chain reaction (RT-PCR) 56 h after injection of pIZT/V5-His-ANK1 using specific primers for *TnBVank1* (Fig. 3a-i). The expression of *TnBVANK1* protein was assessed by western blot analysis using the anti-V5 antibody (Fig. 3a-ii) and detecting the epifluorescence signal of GFP reporter and the *TnBVANK1* protein expression (red signal) by immunofluorescence (Fig. 3b). No signals were detected in RT-PCR, western blot and immunofluorescence experimental controls (haemocytes transfected with the empty pIZT/V5-His vector) except in this last case in




Figure 1. *TnBVank1* induces apoptosis and stimulates caspase-3 activity in S2 cells stably expressing *TnBVank1* gene (S2[ank1]). Apoptosis was induced in positive control S2 cells and S2 cells stably expressing *TnBVank1* through a treatment with 1 μ M camptothecin (S2 + CAM, S2[ank1] + CAM). (a) Cells were harvested and doubly labeled with Annexin V-FITC and PI before analysis by flow cytometry. Dot plots of single positive for Annexin V and double positive for Annexin V and PI cells were interpreted as signs of early and late phases of apoptosis respectively (lower and upper right quadrant respectively). Data are quantified in the bar chart on the right showing the percentage of apoptotic cells. (b) Caspase-3 activity was determined using Ac-DEVD-AMC as a fluorescent substrate. For the assay, 3×10^5 cells were used. Results are expressed as fluorescence units (F.U.); an excitation wavelength of 380 nm and an emission wavelength of 440 nm were utilised. Data represent the mean of 3 replicates \pm SD. Statistically significant differences are indicated with asterisks (* $P \leq 0.05$, ** $P \leq 0.01$, one way ANOVA and Tukey's Test $P < 0.05$).

which the epifluorescence signal of GFP reporter was detected. Different haemocyte types were successfully transfected and the *in vivo* transfection efficiency was calculated by counting fluorescent haemocytes in the control because of the signal of GFP protein, and in haemocytes *in vivo* transfected (see Supplementary Fig. S2). Approximately 80% of haemocytes expressed the GFP reporter (Hemo) or both the GFP reporter and the viral gene *TnBVank1* (Hemo[ank1]), respectively (Fig. 3c).

***In vivo* transient expression of *TnBVank1* and natural parasitism induce apoptosis in *H. virescens* haemocytes.** We investigated the effect of natural parasitism and of *in vivo* transfected haemocytes expressing *TnBVank1* on apoptosis by flow cytometric analysis. Cells undergoing early and late stages of apoptosis were single positive for Annexin V and double positive for Annexin V and PI respectively. Apoptotic cells were detected in haemocytes deriving from parasitised larvae (Hemo[para]) and in haemocytes from *in vivo*




Figure 2. Western blot time course of *TnBVANK1* expression. Cell lysates of S2 cells stably expressing *TnBVank1* were harvested at subsequent passages (2, 4, 10, 20, 26, 39). Detection of *TnBVANK1* was determined by using anti-V5 antibody; each lane was loaded with the same quantity of protein as revealed by the endogenous control, α -Actin. Original Western Blot is reported in Supplementary Information.




Figure 3. Transient expression of *TnBVank1* in *H. virescens* haemocytes. (a) RT-PCR (i) and western blot (ii) showed expression of *TnBVank1* (Hemo[ank1]) and *TnBVANK1* protein (Hemo[ANK1]) 56 h after the injection of pIZT/V5-His-ANK1 into haemocytes of early fifth instar *H. virescens* larvae. *H. virescens* haemocytes extracted from larvae transfected with the empty vector pIZT/V5-His were used as control (Hemo). Original Gel and Western Blot are reported in Supplementary Information. (b) Immunofluorescence of *in vivo* transfected haemocytes showed epifluorescent (for GFP) signal in both control (haemocytes transfected with the empty vector) and *TnBVANK1* (haemocytes transfected with *TnBVank1*) (green signal). Red signal of *TnBVANK1* was detected for haemocytes transfected with *TnBVank1* (ANK1). Nuclei were stained with DAPI (blue signal). Images were observed at immunofluorescence microscopy using Nikon Eclipse 80i equipped with a Nikon Plan Fluor 100x/0.5–1.3 Oil Iris objective, the images were recorded with Nikon Digital Sight DS-U1 camera and ImageJ software. (c) Transfection efficiency was measured by counting fluorescent haemocytes 56 h post injection. Each measurement used randomly chosen 100 haemocytes and it was independently replicated three times. Data represent the mean of 3 replicates \pm SD. Statistical analysis was performed by one way ANOVA and Tukey's Test $P < 0.05$.

transfected larvae (Hemo[ank1]), as well as in the positive control (haemocytes treated with camptothecin). As negative control haemocytes transfected with the empty pIZT/V5-His vector were used and as positive controls, camptothecin (1 μ M) was injected into haemocoel of the larvae with the vector pIZT/V5-His either with or without *TnBVank1* (Fig. 4a).

To further investigate apoptosis with same haemocytes samples, the caspase-3 activity was measured, using the fluorogenic effector caspase substrate Ac-DEVD-ACM. Haemocytes transfected with *TnBVank1* (Hemo[ank1]) and haemocytes deriving from parasitised larvae (Hemo[para]) showed a strong increase of caspase-3 activity compared to haemocytes transfected with the empty pIZT/V5-His vector (negative control) (Hemo) (Fig. 4b).




Figure 4. Parasitism and transient expression of *TnBVank1* in *H. virescens* haemocytes induces apoptosis. (a) Flow cytometry analysis on haemocytes collected from parasitised larvae (Hemo[para]) and from *TnBVank1* *in vivo* transfected haemocytes (Hemo[ank1]). As negative control haemocytes transfected with the empty pIZT/V5-His vector (Hemo) and as positive control haemocytes treated with camptothecin (Hemo + CAM) were used. Cells were doubly labeled with Annexin V-FITC and PI before analysis by flow cytometry. Dot plots of single positive for Annexin V and double positive for Annexin V and PI cells were interpreted as signs of early and late phases of apoptosis respectively (lower and upper right quadrant respectively). Data are quantified in the bar chart on the right showing the percentage of apoptotic cells. (b) Circulating haemocytes were transfected with *TnBVank1* gene (Hemo[ank1]) and their caspase-3 activity was compared to the haemocytes (Hemo), transfected with the empty pIZT/V5-His vector (negative control). Apoptosis was induced in positive controls by injecting into haemocoel of the larvae 1 μ M camptothecin with the vector pIZT/V5-His with or without *TnBVANK1* (Hemo[ank1] + CAM, Hemo + CAM). Caspase-3 activity was determined using Ac-DEVD-AMC as fluorescent substrate. For the assay, 3×10^5 cells were used. Results are expressed as fluorescence (F.U.); an excitation wavelength of 380 nm and an emission wavelength of 430 nm were utilised. Data represent the mean of 3 replicates \pm SD. Statistically significant differences between samples are indicated with asterisk (***) $P \leq 0.001$, one way ANOVA and Tukey's Test $P < 0.05$). (c) Actin filaments were detected with TRITC-conjugated phalloidin staining in haemocytes collected from parasitised larvae (Hemo[para]), from larvae after *TnBVank1* *in vivo* transfection (Hemo[ank1]) and from unparasitised larvae (Hemo). Images were observed at immunofluorescence microscopy using Nikon Eclipse 80i equipped with a Nikon Plan Fluor 100x/0.5–1.3 Oil Iris objective, the images were recorded with Nikon Digital Sight DS-U1 camera and ImageJ software.

Moreover, phalloidin staining showed actin filaments grouped in bundles under the membrane both in haemocytes from parasitised larvae and in *TnBVank1* transfected haemocytes (Fig. 4c). The percentage of apoptotic cells, the caspase-3 activity and the phalloidin staining were comparable in parasitised haemocytes and in *TnBVank1* transfected haemocytes. These results suggest that the apoptosis does not seem to be due to an overexpression of the gene *TnBVank1* after *in vivo* transfection respect to its expression in haemocytes after parasitisation. Indeed, when a qPCR on haemocytes collected after *in vivo* transfection with *TnBVank1* and after parasitisation was performed, results did not show significant differences of *TnBVank1* transcript expression level after parasitisation and after *in vivo* transfection (see Supplementary Fig. S3).

Also the nuclei staining with Hoechst 33342 revealed the occurring of apoptosis in *H. virescens* haemocytes transfected with *TnBVank1* (Hemo[ank1]). The cells in the haemocytes transfected with the empty pIZT/V5-His vector (negative control) (Hemo) showed roundshaped and homogeneously stained nuclei, whereas haemocytes transfected with *TnBVank1* (Hemo[ank1]) showed accumulation of fluorescent dye indicating obvious chromatin condensation and fragmentation (see Supplementary Fig. S1).

Functional proteomics experiments. Isolation of *TnBVANK1* complexes was performed on a total protein extract from S2 cells stably expressing a V5-tagged form of the protein. Following pre-cleaning treatment, the unbound fraction was immunoprecipitated with an anti-V5 specific antibody in order to capture *TnBVANK1* functional complexes. The occurrence of *TnBVANK1* protein within the immunoprecipitated material was detected by western blot using anti-V5 antibody (see Supplementary Fig. S4a). Immunoprecipitated complexes were fractionated by SDS-PAGE (see Supplementary Fig. S4b) and protein bands, excised from the gel, were identified by LC-MS/MS analysis. Protein bands from the control S2 cells were also analysed as control. Putative *TnBVANK1* protein interactors were obtained by differential analysis comparing the proteins identified in S2 cells stably expressing *TnBVank1* and those from the control S2 cells.

The results of this functional proteomic approach are summarised in Supplementary Table S1. As the experiment was performed on the *Drosophila* S2 cell line, putative interactors were annotated against the *Drosophila melanogaster* genome and then compared to *Homo sapiens* proteins. Among interactor proteins, NF- κ B p110 subunit was identified, confirming previous studies showing interaction between *TnBVANK1* and the transcription factor NF- κ B⁸.

Moreover, we found several proteins involved in the apoptotic pathway, as the transcription factor Stwl²³, the mitochondrial carrier homolog 1 (Mtch)²⁴, ALG-2 interacting protein X²¹ and the GTPase Rab1^{25,26}. Among these proteins, we focused our attention on the ALG-2 interacting protein X, also called Alix. This protein was chosen on the basis of its involvement in programmed cell death, for subsequent analyses.

***TnBVANK1* protein colocalises with Alix protein in polyclonal S2 cell line.** To further support the interaction between *TnBVANK1* and Alix, the localization of both these proteins was examined in polyclonal S2 cell line by immunofluorescence labeling using both anti-Alix and anti-*TnBVANK1* antibodies. Figure 5(ii,iii) shows that both *TnBVANK1* (green signal) and Alix (red signal) are widely distributed throughout the cytoplasm of polyclonal S2 cell line and the merged picture (Fig. 5-iv) indicates that the two proteins seem to colocalise, as there is a clear overlap of red and green signals of analysed proteins. This result further supports the potential to interact of the two proteins as demonstrated by functional proteomics experiments. In the control, represented by *Drosophila* S2 cells, as expected, no signal of *TnBVANK1* was detected while Alix signal is present in the cytoplasm as in polyclonal S2 cells. This finding provides a further element supporting the hypothesis of the interaction between these two proteins.

***In vitro* and *in vivo* silencing of Alix suppresses caspase-3 activity induced by *TnBVank1*.** In order to assess whether the apoptosis induced by *TnBVank1* is mediated by the interaction with Alix, we silenced the Alix gene transcript by RNAi both in the S2 cell line stably expressing *TnBVank1* and in *H. virescens* circulating haemocytes transiently expressing *TnBVank1*.

The success of RNAi was verified by quantitative Real time PCR (qPCR) and western blot performed on polyclonal S2 cells stably expressing *TnBVank1*, harvested at different time points (0, 6, and 10h) (Fig. 6a and b) and on *H. virescens* haemocytes transiently expressing *TnBVank1* (Fig. 7a and b) both treated with Alix dsRNA. The cells in which Alix was silenced (S2[ank1] Alix Si and Hemo[ank1] Alix Si) showed that the available transcript of Alix gene was significantly reduced compared to its expression in S2 cell line, in S2 cell line stably expressing *TnBVank1* (S2[ank1]) and in S2 cell line stably expressing *TnBVank1* treated with the ds*Apis*OBP3 (S2[ank1] OBP3 Si) as well as in circulating haemocytes transiently transfected with the empty pIZT/V5-His vector (Hemo), in haemocytes transiently expressing *TnBVank1* (Hemo[ank1]) and in haemocytes transiently expressing *TnBVank1* treated with the ds*Apis*OBP3 (Hemo[ank1] OBP3 Si) (Fig. 7a).

These results were confirmed by western blot analysis showing that Alix protein expression was reduced in S2 cells and haemocytes after RNAi experiments (Figs 6b and 7b).

Same samples were tested by qPCR for their expression of *TnBVank1* transcript demonstrating its expression in all described conditions except in the case of S2 cell line and in circulating haemocytes transiently transfected with the empty pIZT/V5-His vector (Figs 6a and 7b).

Caspase-3 assay performed on same samples analysed by qPCR as listed above indicated that in S2 cell line stably expressing *TnBVank1* and silenced for Alix transcript, caspase-3 activity was highly reduced compared to the S2 cell line stably expressing *TnBVank1* (Fig. 6c). Similar results were obtained *in vivo*, since caspase-3 activity measured in *H. virescens* haemocytes cotransfected with Alix dsRNA and *TnBVank1* was significantly reduced compared to haemocytes transfected with *TnBVank1* alone (Fig. 7c). Taken together, the *in vivo* and *in vitro* experiments suggest that Alix/*TnBVANK1* interaction strongly contributes in inducing apoptosis both in S2 cell line and in circulating haemocytes.




Figure 5. *TnBVANK1* protein colocalises with Alix in polyclonal S2 cell line. i-iv) Polyclonal S2 cell line coexpressing both *TnBVANK1* and Alix proteins stained with anti-*TnBVANK1* and anti-Alix antibodies. v-viii) Staining of control S2 cells expressing only Alix protein. The localisation of *TnBVANK1* and Alix proteins was determined by FITC (green signal) and TRITC (red signal) fluorescence, respectively and nucleus was stained with DAPI (blue signal). In merged images there is a clear overlap of red and green signals showing a colocalisation (yellow signal) of analysed proteins. Images were observed at immunofluorescence microscopy using Nikon Eclipse 80i equipped with a Nikon Plan Fluor 100x/0.5–1.3 Oil Iris objective, the images were recorded with Nikon Digital Sight DS-U1 camera and ImageJ software.

Discussion

The bracovirus associated with the wasp *Toxoneuron nigriceps* (*TnBV*) is one of parasitic regulatory factors of maternal origin involved in immunosuppression and alteration of endocrine balance of the host *Heliothis virescens*^{1,4,12}. A large number of studies have been addressed to the elucidation of parasitised host developmental syndromes^{6,7,18,27}. Briefly, in this experimental model, parasitised last instar larvae fail to pupate, allowing a suitable nutritional substrate for parasitoid larvae over an extended time period²⁸. Here, we report experimental evidence demonstrating the role of *TnBVank1*, a viral gene member of the *TnBV* ankyrin (*ank*) gene family⁸, in inducing apoptosis processes in a stably *TnBVank1* transfected polyclonal insect cell line (*Drosophila* Schneider's S2 cells) and in circulating haemocytes of *H. virescens* larvae transiently transfected with *TnBVank1* gene, thus reproducing the apoptotic effects observed in haemocytes of parasitised *H. virescens* larvae. *TnBVank1*, together with *TnBVank2* and *TnBVank3*, belongs to the *TnBVank* family⁸ and displays significant sequence similarity with members of the I κ B family, that are proteins generally involved in the control of NF- κ B signaling pathways both in insects and in vertebrates²⁹.

I κ B members are characterised by repeated sequences of ankyrins that bind to transcription factors NF- κ B/Rel, masking their nuclear localization signal. Upstream sequences of ankyrin serine residues can be phosphorylated by the I κ B kinase (I κ K) in response to a number of extracellular signals²⁹. Once phosphorylated, I κ B can be released and degraded via the proteasome, allowing translocation of the transcription factor NF- κ B/Rel in the nucleus and subsequent activation of target genes²⁹. In *Drosophila*, the I κ B-like protein regulates various cellular responses triggered by nuclear transfer protein NF- κ B/Rel, such as the embryonic development and the release of defense-related antimicrobial peptides²⁹.

The gene family of viral ankyrins (also called *vankyrin*), is widespread among both Bracovirus and Ichnovirus, and each genome has several copies of these genes^{8,15,30–32}. All known polydnavirus (PDV) ankyrin genes lack the




Figure 6. Alix and *TnBVank1* transcript levels in S2 cells and post-Alix silencing caspase-3 assay. **(a)** Expression levels of Alix and *TnBVank1* at different hours (T₀, T₆, T₁₀) after Alix silencing in S2 cells stably expressing *TnBVank1* (S2[ank1] Alix Si), normalised to the endogenous control Rpl32⁵⁴. As control ds*ApisOBP3* was used (S2[ank1] OBP3 Si). No expression of *TnBVank1* was observed in the control S2 cells. **(b)** Western blot of Alix expression at different hours (T₀, T₆, T₁₀) after Alix silencing in S2 cells stably expressing *TnBVank1*. Original Western Blot is reported in Supplementary Information. **(c)** Post-Alix silencing caspase-3 activity in S2 cells, S2 cells stably expressing *TnBVank1* (S2[ank1]), S2 cells stably expressing *TnBVank1* with silenced Alix transcript (S2[ank1] Alix Si) and S2 cells stably expressing *TnBVank1* transfected with dsRNA of *ApisOBP3* as control (S2[ank1] OBP3 Si), harvested at different times (0, 6 and 10 h) after the replacement of the medium containing dsRNA. Data represent the mean of 3 replicates \pm SD. Statistically significant differences between samples are indicated with asterisk (***) $P \leq 0.001$, one way ANOVA and Tukey's Test $P < 0.05$.

regulatory sites, serine residues at the N-terminus and PEST domain at the C-terminal end^{13,15,33–36}. These peculiar features strongly suggest that viral ankyrins may act as inhibitors by irreversibly binding and retaining NF- κ B in the cytoplasm thus affecting a variety of cellular responses in parasitised hosts, including alteration of innate immune defenses^{8,14,15}. This mechanism, proposed to explain the alteration of the host immune response induced by the parasitoid, is very similar to the strategy used by ASFV (African swine fever virus) to inhibit the activation of NF- κ B transcription factors in mammals. The ASFV I κ B-like protein irreversibly binds NF- κ B, preventing its entry into the nucleus³⁷, similarly to the effect observed for *TnBVank1*⁸.

The genome of *Campoletis sonorensis* Ichnovirus (CsIV) contains seven copies of ankyrin genes named vankyrin³³. Among these vankyrins, P-vank-1 is an interesting viral factor, as it, when is stably expressed in eukaryotic Sf9 insect cells, seems to have a role in protecting cells from apoptosis³⁴. The role of vankyrin P-vank-1, which belongs to the same family as *TnBVank1*, led us to hypothesise a possible involvement of *TnBVank1* in the regulation of programmed cell death. Here we have shown that, unlike vankyrin P-vank1, the viral gene *TnBVank1* induces apoptosis. The different biological roles of these proteins are surprising but not unusual; it is reported that the vankyrin I²-vank-3 of *Campoletis sonorensis* Ichnovirus (CsIV), although highly similar to P-vank-1 (with 83% amino acid sequence identity), plays a different role³⁶. This could be also explained by their differential expression and localisation in the host tissues.

In the polyclonal S2 cell line stably expressing *TnBVank1*, a significant increase of apoptotic cell number, compared to untreated S2 cells (negative control), was observed by flow cytometry when doubly stained with Annexin V-FITC and PI. Moreover, compared to negative control, in the polyclonal S2 cell line stably expressing *TnBVank1* a significantly higher caspase-3 activity and apoptotic nuclei were detected. A previous work¹⁶ revealed typical morphological modifications of apoptosis in parasitised *H. virescens* granulocytes (a subpopulation of haemocytes). Here we confirmed that haemocytes undergo apoptosis after natural parasitism, by flow cytometry analysis, caspase-3 assay and phalloidin staining. This result led us to deepen the role of *TnBVank1*




Figure 7. Alix and *TnBVank1* transcript levels in *H. virescens* haemocytes and post-Alix silencing caspase-3 assay. **(a)** Expression level of Alix and *TnBVank1* transcripts into *H. virescens* haemocytes 56 h after Alix silencing (Hemo[ank1] Alix Si), normalised to the endogenous controls (EF1 α and RP13⁴⁹). No expression of *TnBVank1* was observed in the control haemocytes transfected with the empty pIZT/V5-His vector (Hemo). **(b)** Western blot of Alix expression in haemocytes 56 h after Alix silencing and transfected with *TnBVank1* (Hemo[ank1] Alix Si); as control haemocytes were transfected with the empty vector (Hemo). Original Western Blot is reported in Supplementary Information. **(c)** Post-Alix silencing caspase-3 activity in haemocytes, transfected with *TnBVank1* gene (Hemo[ank1]), cotransfected with Alix dsRNA and *TnBVank1* (Hemo[ank1] Alix Si). As control haemocytes were transfected with the empty vector (Hemo) and cotransfected with *ApisOBP3* dsRNA and *TnBVank1* (Hemo[ank1] OBP3 Si). Data represent the mean of 3 replicates \pm SD. Statistically significant differences between samples are indicated with asterisk (***) $P \leq 0.001$, one way ANOVA and Tukey's Test $P < 0.05$).

in apoptotic pathway in haemocytes of non parasitised larvae by *in vivo* transient transfection that is an useful method to determine the biological function of a foreign gene. We observed an apoptotic profile by flow cytometry, a significantly increased caspase-3 activity and nuclear morphological changes in *H. virescens* haemocytes after *TnBVank1* *in vivo* transfection. These results demonstrate that *TnBVank1* is one of the Polydnavirus genes responsible for apoptosis observed in *H. virescens* haemocytes after parasitisation¹⁶. A qPCR performed on both haemocytes derived from parasitised host and *TnBVank1* *in vivo* transient transfected haemocytes evidenced that the expression level of *TnBVank1* after transient expression was comparable to its expression after natural parasitism, thus demonstrating that the observed apoptotic effect is not due to the result of an unnatural overexpression of the protein.

TnBVank1 would be the first viral ankyrin that, besides preventing the nuclear translocation of NF- κ B⁸ thus disrupting cell pathways under its control^{15,29,38}, is also able to directly induce apoptosis. *TnBVank1* is not the first *TnBV* gene that induces apoptosis. Lapointe and colleagues¹⁸ showed that the gene *TnBV1* induces apoptosis when expressed in cell lines derived from *Spodoptera frugiperda* (SF21) and *Tricoplusia ni* (High Five). However, when the authors tried the *in vivo* expression of *TnBV1*-recombinant baculovirus, no apoptotic effect was detected, leaving the role of *TnBV1* expression in parasitised *H. virescens* larvae elusive. *TnBV1* protein may interact with other *TnBV* proteins during parasitism to exert its function in the apoptotic pathway¹⁸. In contrast,

our results indicate that a single viral gene, *TnBVank1*, is able to induce apoptosis, *in vitro* and *in vivo*, without the requirement of other virus-encoded genes as interaction partners. The induction of apoptosis is one of the most common strategies adopted by various PDVs to suppress the immune response³⁹. However, the parasitism do not affect all the cells of the immune system¹⁶ probably as a strategy to allow the host to respond to a possible further pathogen invasions ensuring the development of wasp embryo^{39,40}.

The molecular mechanisms of apoptosis induction exerted by *TnBVank1* were more closely investigated by functional proteomic approaches. Multiprotein complexes comprising *TnBVANK1* were isolated from polyclonal S2 stably expressing *TnBVank1* by immunoprecipitation and the individual protein components were identified by mass spectrometry. Among the various *TnBVANK1* protein partners, we focused our attention on Alix (ALG-2 interacting protein X) as this multifunctional protein is involved in several protein interaction networks, including programmed cell death, endocytic membrane trafficking and cytoskeletal remodeling^{41,42}. To confirm the possible interaction between *TnBVANK1* and Alix proteins, polyclonal S2 cell line was examined by co-immunolocalisation experiments, using anti-*TnBVANK1* and anti-Alix antibodies. Observations by immunofluorescence microscopy revealed that *TnBVANK1* are both present in the cytoplasm and seems to colocalises with Alix protein.

The role of *TnBVANK1*-Alix interaction in apoptosis was further investigated by silencing experiments both *in vitro* and *in vivo*. Silencing of Alix expression in *TnBVank1* stably transfected S2 cells resulted in a significant reduction of apoptosis assessed by caspase-3 activity. Similarly, silencing Alix in circulating haemocytes of *H. virescens* larvae transiently transfected with *TnBVank1* drastically reduced caspase-3 activity. In both cases, the *TnBVank1* transcript expression level evaluated by qPCR remained constant.

These *in vitro* and *in vivo* results suggest that *TnBVank1* induces apoptosis through interaction with Alix.

The interaction between *TnBVANK1* and Alix was also reported by a recent study showing their colocalisation in prothoracic glands of a *Drosophila melanogaster* mutated strain expressing *TnBVank1*. It is reported that *TnBVANK1* impairs the ecdysone biosynthesis and causes the developmental arrest by altering the vesicular traffic of ecdysteroid precursors in the prothoracic gland steroidogenic cells of *Drosophila melanogaster*. The authors showed that the *TnBVank1* affects the sterol delivery interacting with Alix that is also involved in controlling the cholesterol export from endosomes^{16,43}.

Although the molecular mechanisms are unclear, Alix cooperating with ALG-2 is involved in the induction of apoptosis through both intrinsic and extrinsic apoptotic pathways⁴¹. A possible way of induction of apoptosis by the protein Alix is via TNF-R1 (TNF receptor-1), which contains a DD (Death Domain) capable of activating caspase 8, an initiator of apoptotic process. The interaction TNF-R1/Caspase 8 initiates the apoptotic cascade. This mechanism requires that Alix forms a complex with TNF-R1⁴⁴.

Our data show that *TnBVANK1* strongly induces apoptosis interacting with Alix, via the mitochondrial apoptotic pathway. Moreover, the previously demonstrated role of *TnBVank1* in the control of the NF- κ B signaling⁸ preventing its nuclear translocation, could also further increase the sensitivity of the cell to the cell death signals³⁸.

This seems like an effective strategy of the virus to disable the host immune system and facilitate successful development of its associated wasp.

Materials and Methods

Insect cells culture. *Drosophila* Schneider's S2 cells (Life Technologies, Carlsbad, CA, USA) were cultured in Schneider's *Drosophila* medium (WVR, Radnor, PA, USA) supplemented with 10% heat inactivated fetal bovine serum (Life Technologies, Carlsbad, CA, USA) and were maintained in 75 cm² tissue culture flasks (Corning, NY, USA) at 27°C.

Stable transfection of *TnBVank1* in S2 cells. The *TnBVank1* cDNA, obtained as previously described by Falabella and colleagues⁸, was amplified with specific primers containing XbaI and SacII enzyme restriction sites (underlined):

Ank1F 5'-CTAGTCTAGAATGGAAACTCATTACTCATTG-3'
Ank1R 5'-TCCCCGCGGATTATCATCACACTTAGCGCC-3'

The obtained amplicon was cloned into TOPO vector (Life Technologies, Carlsbad, CA, USA) and sequenced. The insert was then released by XbaI and SacII restriction enzyme digestion (underlined) and cloned into the expression vector pIZT/V5-His (Life Technologies, Carlsbad, CA, USA). This vector uses the OpIE2 promoter from the *Orygia pseudotsugata* baculovirus for constitutive expression of the gene of interest and encodes a zeocin-green fluorescent protein (GFP) gene fusion under control of the OpIE1 promoter. The 3' ends of the Ank1 fragments were cloned in frame with a "V5 epitope" and with a tail of six histidines. The obtained construct was sequenced, verified and transiently expressed in S2 cells by cationic lipid-mediated transfection. 24 h prior transfection S2 cells were seeded at 70 to 80% confluency (0.5×10^6) in 12-well culture plates (Corning, NY, USA) and subsequently transfected using 8 μ l of Cellfectin (Life Technologies, Carlsbad, CA, USA) and 1 μ g of DNA per 1 ml of medium and incubated at 27°C for 6 h. After incubation, the medium was removed and cells were re-incubated with 1 ml of Schneider's *Drosophila* complete medium containing 10% FBS at 27°C for 48 h. Stably transformed cells were selected by adding 400 μ g/ml of zeocin (Life Technologies, Carlsbad, CA, USA). Populations of antibiotic-resistant cells were amplified to obtain polyclonal S2 cell lines stably expressing *TnBVank1*.

Insect rearing. *Heliothis virescens* larvae were maintained on an artificial diet as previously described²⁸. *Toxoneuron nigriceps* was reared in the laboratory, on larval stages of its host, *H. virescens*, according to Vinson *et al.*⁴⁵. Rearing temperature was 29 ± 1 °C for the host, parasitised host larvae and cocoons. *T. nigriceps* adults were kept

at 25 ± 1 °C and fed with water and honey. A 16 h light photoperiod was adopted and the relative humidity was $70 \pm 5\%$. *H. virescens* last instar larvae were staged according to Webb & Dahlman⁴⁶ and synchronised as reported by Pennacchio *et al.*⁴⁷.

For experimental procedures, described below, *H. virescens* larvae were parasitised at the day one of the last larval instar.

***In vivo* transient expression of *TnBVank1* in *H. virescens* haemocytes.** For *in vivo* transfection of pIZT/V5-His-ANK1, Lipofectin reagent (Life Technologies, Carlsbad, CA, USA) was used. pIZT/V5-His-ANK1 (0.5 µg) was mixed with 4 µl of transfection reagent and incubated for 20 min at room temperature for the formation of DNA-lipid complex before injection into the haemocoel of *H. virescens* fifth instar larvae (day one) using a Hamilton syringe (Sigma-Aldrich St Louis, MO, USA). Before injection, larvae were anaesthetised by immersion in sterile ice water. Haemocytes were collected as described by Ferrarese and colleagues¹⁶ 56 h post-transfection to perform all the biological assays. This time was selected as the most successful for *TnBVANK1* expression, tested after a time course at 24 h, 48 h and 56 h post injection (data not shown).

Success of the haemocytes transfection was analysed by reverse transcriptase-polymerase chain reaction, and the protein expression was detected by western blot analysis using anti-V5 antibody (Life Technologies, Carlsbad, CA, USA) and by immunofluorescence experiments. To assess whether *TnBVANK1* induces apoptosis *in vivo*, the caspase-3 assay was performed on haemocytes as describe below.

Reverse transcriptase-polymerase chain reaction. Total RNA was extracted from *H. virescens* haemocytes, previously transfected with the vector pIZT/V5-His (as negative control) or recombinant vector expressing *TnBVank1*, by using TRI Reagent (Sigma-Aldrich St Louis, MO, USA). Concentration and purity of the total RNA samples were measured using the NanoDrop ND-1000 Spectrophotometer (ND-1000, Thermo Scientific, Waltham, MA, USA). To remove DNA contamination, the samples were treated with 1U of DNase I (Life Technologies, Carlsbad, CA, USA) per microgram of RNA, for 15 min at room temperature. The reaction was stopped by adding 1 µl of 25 mmol/L EDTA and incubating at 65 °C for 10 min. The complementary DNA was synthesised by SuperScript III Reverse Transcriptase (Life Technologies, Carlsbad, CA, USA) according to the manufacturer's protocol. The cDNA was used as a template for PCR amplification using the following primers:

Ank1F 5'-ATGGAAAACCTCATTACTCATTG-3'
Ank1R 5'-ATTATCATCACACTTAGCGCC-3'

To confirm no vector DNA contamination was present after RNA extraction, an aliquot of the isolated RNA was directly used as a template for a test PCR (i.e. without reverse transcription), which showed no PCR product.

***In vivo* transient transfection efficiency.** *Heliothis virescens* haemocytes were isolated 56 h after injection of the vector pIZT/V5-His with or without *TnBVank1* as described above. Haemocytes were fixed for 10 min at room temperature in 4% formaldehyde in 1X phosphate buffer saline (PBS) pH 7.5 and then permeabilised for 30 min with PBS containing 0.1% Triton X-100. The cells were blocked with 1% bovine serum albumin (BSA) for 30 min and, subsequently, only the haemocytes transfected with *TnBVank1* were incubated with the primary antibody, anti-*TnBVANK1*⁴⁸ diluted 1:200 in BSA 1%, in a humidified chamber at 4 °C overnight. After three washes in PBS 1X, the cells were incubated with the secondary antibody TRITC-conjugated (Sigma-Aldrich St Louis, MO, USA), 1:1000 in 1% BSA for 1 h at room temperature in dark condition. After washing in PBS 1X three times, the slides were mounted with FluoroshieldTM with DAPI, histology mounting medium (Sigma-Aldrich St Louis, MO, USA).

To measure the total cell fluorescence, the slides were observed microscopically with Nikon Eclipse 80i equipped with a Nikon Plan Fluor 100x/0.5–1.3 Oil Iris objective and the images were recorded with Nikon Digital Sight DS-U1 camera. The intensity of the cell fluorescence was measured using the ImageJ software (<http://rsbweb.nih.gov/ij/>).

The GFP protein expression was analysed directly by counting fluorescent haemocytes compared to total haemocytes. A hundred haemocytes from a randomly selected field of view at 40x magnification were analysed for each measurement. Each treatment was independently replicated three times.

***TnBVank1* expression in haemocytes after *in vivo* transfection and after parasitism.** To evaluate *TnBVank1* expression in haemocytes both *in vivo* transfected and after parasitism, quantitative RT-PCR (qPCR) experiments were carried out. RNA extraction and cDNA synthesis were conducted as previously described. PCR amplification was performed using GoTaq qPCR Master Mix (Promega, Fitchburg, WI, USA) on an ABI PRISM[®] 7500 Fast Real-Time PCR System Thermal Cycler (Applied Biosystems, USA) and in a 20 µl final volume. Sequence specific primers, each in a concentration of 0.3 µM, were:

Ank1F 5'-ATCATGAGGGGCATGCATG-3'
Ank1R 5'-AGCACCAGCGCTGTGAAGT-3'

As reference genes we used the *H. virescens* genes coding for the Elongation Factor 1-alpha (EF1α) and 60S Ribosomal Protein 13 (RP13)⁴⁹ (see Supplementary Fig S5). Primers were:

EF1αF 5'-TTGAAGCCTGGTACCATCGT-3'
EF1αR 5'-GTTGGTGGGTTGTTCTTGG-3'

RP13F 5'-TCGTGGTAAGGTGAAGGCAT-3'
 RP13R 5'-AGTCACAGCCTCAACGATCT-3'

All qPCR analyses were performed for a set of 3 biological replicates. Quantification analysis of amplification was done using the comparative C_T (ΔC_T) method. The efficiencies of the three amplicons were approximately equal (Ank1 = 0.892159; EF1 α = 0.914590; Rp13 = 0.896174).

In vitro and in vivo flow cytometric analysis. Apoptosis was evaluated by flow cytometric analysis of Annexin V and Propidium Iodide (PI)-stained cells using fluorescein isothiocyanate (FITC) Annexin V Apoptosis Detection kit I (Becton Dickinson, Biosciences, San Jose, CA, USA). The assay was carried out on S2 cells (negative control), on S2 cells stably expressing *TnBVank1* and on *H. virescens* haemocytes after *TnBVank1* *in vivo* transfection and after parasitism. S2 cells and polyclonal S2 cells were used in passages 4–10. For the positive control, apoptosis was induced by treating the cells with 1 μ M camptothecin, a cytotoxic alkaloid inhibitor of Topoisomerase I, known to induce apoptosis^{50,51} (Sigma-Aldrich St Louis, MO, USA), or injecting into haemocoel of larvae directly 1 μ M camptothecin for 20 h³⁴. As negative control for the *in vivo* assay, *H. virescens* haemocytes after injection of the vector pIZT/V5-His without *TnBVank1* were used. Cell samples were washed twice and incubated with Annexin V-FITC/PI for 15 min as manufacturer's protocol. Stained samples were acquired using Navios flow cytometer and analysed by Kaluza Analysis 1.3 software (Beckman Coulter). Single positive for Annexin V and double positive for Annexin V and PI cells were interpreted as signs of early and late phases of apoptosis respectively.

In vitro and in vivo Caspase-3 activity assay. The caspase-3 fluorimetric assay was carried out on S2 cells (negative control), on S2 cells stably expressing *TnBVank1* and on *H. virescens* haemocytes after *TnBVank1* *in vivo* transfection. S2 cells and polyclonal S2 cells were used in passages 4–10. For the positive control, apoptosis was induced by treating the cells with 1 μ M camptothecin^{50,51} (Sigma-Aldrich St Louis, MO, USA) or injecting into haemocoel of larvae directly 1 μ M camptothecin for 20 h³⁴. As negative control for the *in vivo* assay, *H. virescens* haemocytes after injection of the vector pIZT/V5-His without *TnBVank1* were used. As a substrate of caspase-3 the Ac-DEVD-AMC caspase-3 fluorogenic substrate (BD Biosciences Pharmingen, San Diego, CA, USA) was used. The assay protocol requires primarily lysis of cells. Cells were centrifuged at 1000 g for 10 min and then resuspended with Cell Lysis Buffer (10 mM Tris-HCl, 10 mM NaH₂PO₄/NaHPO₄, pH 7.5, 130 mM NaCl, 1% Triton-X-100 and 10 mM NaPPI) in suitable amounts such as to obtain 3×10^5 cells/100 μ l. The peptide Ac-DEVD-AMC (20 μ M) and 1 ml of Protease Assay Buffer (20 mM HEPES pH 7.5, 10% glycerol, 2 mM DTT) were added to each reaction. The reaction was then incubated at 37 °C for 1 h. The amount of AMC released was measured by the spectrofluorometer (Cary Eclipse Fluorescence Spectrophotometer, Agilent Technologies, Walnut Creek, CA, USA) exciting at 380 nm and detecting in a range between 430–460 nm.

Phalloidin staining. Haemocytes were isolated 56 h after injection of the vector pIZT/V5-His with or without *TnBVank1* as described above and from parasitised larvae. Cells were fixed, permeabilised and blocked as previously described. Subsequently, haemocytes were incubated with TRITC-conjugated phalloidin (Sigma-Aldrich St Louis, MO, USA), diluted 50 μ g/ml in 1% BSA for 2 h at room temperature in dark condition. After washing in PBS 1X three times, the slides were mounted with glycerol (Sigma-Aldrich St Louis, MO, USA).

The slides were observed microscopically with Nikon Eclipse 80i equipped with a Nikon Plan Fluor 100x/0.5–1.3 Oil Iris objective and the images were recorded with Nikon Digital Sight DS-U1 camera.

Hoechst staining. S2 cells (negative control), S2 cells stably expressing *TnBVank1* and *H. virescens* haemocytes isolated 56 h after injection of the vector pIZT/V5-His with or without *TnBVank1* were stained with Hoechst 33258 (Sigma-Aldrich, St Louis, MO, USA) to detect nuclear morphology. Cells were fixed with 4% paraformaldehyde for 10 min, washed with PBS, and stained with 10 μ g/mL Hoechst 33258 at room temperature for 10 min in the dark. The cells were washed with PBS 1X three times and the slides were mounted with glycerol for morphologic observation by fluorescence microscopy (NIKON Eclipse 80i) equipped with a Nikon Plan Fluor 40x and 100x/0.5–1.3 Oil Iris objective and the images were recorded with Nikon Digital Sight DS-U1 camera.

Western blot time course. The *TnBVANK1* protein production over time in the polyclonal S2 cell line that stably expresses *TnBVank1* was evaluated by western blot analysis. For proteins extraction, S2 cell line expressing *TnBVank1*, harvested at different successive splitting (2, 4, 10, 20, 26, 39 passages), was resuspended, transferred to sterile 1.5 ml tubes (Eppendorf, Hamburg, Germany) and centrifuged at 1000 g for 5 min at room temperature. The cell pellet was then resuspended in lysis buffer (50 mM Tris-HCl, pH 7.8, 150 mM NaCl, 1% Nonidet P-40) to which 1% protease inhibitor cocktail (Sigma-Aldrich, St Louis, MO, USA) was added. After incubation at 37 °C for 10 min, the lysed cells were centrifuged at 16000 g for 10 min at room temperature and the supernatant containing soluble proteins was transferred to a new 1.5 ml tube (Eppendorf, Hamburg, Germany). The protein concentration was determined by the Bradford method⁵², using BSA as standard and 100 μ g of protein lysate were used for each SDS PAGE gel lane. The viral protein *TnBVANK1* was detected using the murine anti-V5 antibody (Life Technologies, Carlsbad, CA, USA) as primary antibody (diluted 1:5000 in 5% milk), and anti-mouse conjugated to horseradish peroxidase (Life Technologies, Carlsbad, CA, USA) (diluted 1:5000 in PBS), as secondary antibody. α -actin was chosen as endogenous control, using the anti- α -actin antibody (Sigma-Aldrich, St Louis, MO, USA) (diluted 1:5000 in 5% milk) and the secondary antibody anti-rabbit conjugated to horseradish peroxidase (Life Technologies, Carlsbad, CA, USA, USA) (diluted 1:15000 in PBS). Finally, the Western blot Chemiluminescent HRP Substrate (LiteAblo, Euroclone, Milan, Italy) to detect target proteins was used.

Functional proteomics. Functional proteomic experiments were performed on protein lysates of S2 cells stably expressing V-5-tagged *TnBVANK1*. A total protein extract was subjected to pre-cleaning treatment on underivatized agarose beads and the unbound fraction was immunoprecipitated by incubation onto anti-V5 agarose-conjugated antibody overnight under gentle stirring. An aliquot of the non specific bound material was eluted and used as control.

Beads were collected by centrifugation (3,000 rpm for 5 min) and extensively washed with lysis buffer supplemented with 150 mmol/l NaCl. Elution was performed by competition with V-5 peptide in elution buffer. The presence of *TnBVANK1* in the eluted fraction was assessed by Western-blot using anti-V5 antibody.

The eluted proteins were precipitated in methanol/chloroform and then loaded onto a 10% SDS-PAGE. Protein bands were excised, reduced with 10 mM DTT and carboxyamidomethylated with 55 mM iodoacetamide. Tryptic digestion was carried out with 12.5 ng/μl at 4 °C for 2 h in 10 mM NH₄HCO₃ buffer, pH 7.8 and then for 16 h at 37 °C. Peptides were then extracted with 10 mM NH₄HCO₃ and 1% formic acid in 50% acetonitrile at room temperature. Gel slices from the negative control S2 cells were also analysed.

Peptide mixtures were analysed by LC-MS/MS using a CHIP MS 6520 QTOF equipped with a capillary 1200 HPLC system and a chip cube (Agilent Technologies, Palo Alto, CA, Italy). Peptide analysis was performed using data-dependent acquisition. Mass spectral data were used to search for the UniProt/SwissProt protein database using an in house version of MASCOT (Matrix Science, Boston, USA).

Coimmunofluorescence analysis. For the immunofluorescent labeling, S2 cells were fixed for 10 min at room temperature in 4% formaldehyde in 1X phosphate buffer saline (PBS) pH 7.5. After permeabilisation for 30 min in 0.1% Triton X-100 solution and washing in PBS 1X, the cells were incubated in 1% BSA for 30 min and, subsequently, with the mixture of two primary antibodies, anti-*TnBVANK1*⁴⁵ and anti-Alix⁵³, 1:200 in BSA 1%, in a humidified chamber at 4 °C overnight. After three washes in PBS 1X, the cells were incubated with the mixture of two secondary antibodies, which were raised in different species (with two different fluorochromes, i.e. TRITC-conjugated against mouse and FITC-conjugated against rabbit) (Sigma-Aldrich St Louis, MO, USA), 1:1000 in 1% BSA for 1 h at room temperature in dark condition. After washing in PBS 1X, the slides were mounted with Fluoroshield™ with DAPI, histology mounting medium (Sigma-Aldrich St Louis, MO, USA).

To measure the total cell fluorescence, the slides were observed microscopically with Nikon Eclipse 80i equipped with a Nikon Plan Fluor 100x/0.5–1.3 Oil Iris objective and the images were recorded with Nikon Digital Sight DS-U1 camera. The intensity of the cell fluorescence was measured using the ImageJ software (<http://rsbweb.nih.gov/ij/>).

In vivo and in vitro Alix silencing by RNA interference. RNA interference (RNAi) experiments both *in vivo* and *in vitro* were performed using double-stranded (ds)RNA of the Alix gene (Accession Number Q9VB05) (see Supplementary Fig. S6). As negative control dsRNA of the *ApisOBP3* gene (Accession Number 001160057), which is not expressed in S2 cells and in *H. virescens* haemocytes, was used.

Part of the coding sequences for Alix and *ApisOBP3* proteins were amplified by PCR with specific primers containing the T7 binding site at their 5' ends (underlined):

S2AlixF 5'-TAATACGACTCACTATAGGGAGAATTGGCGAGGAGATTGCT-3'
 S2AlixR 5'-TAATACGACTCACTATAGGGAGAATCGCGAAGTCATCACCAAT-3'
*H.vir*AlixF 5'-TAATACGACTCACTATAGGGAGATGAAAGGTATGGAGACTGA-3'
*H.vir*AlixR 5'-TAATACGACTCACTATAGGGAGAATCGAAGGTGTAGCAGCTG-3'
 OBP3F 5'-TAATACGACTCACTATAGGGGTAATAAAACAAGGCGCACAG-3'
 OBP3R 5'-TAATACGACTCACTATAGGGATCGTCGTCGGATCAAGGAA-3'

PCR products were gel purified (Quantum Prep™ Freeze'N Squeeze DNA Gel Extraction Spin Columns, Biorad, Hercules, CA, USA) and Alix dsRNA and *ApisOBP3* dsRNA were prepared using the RNAi MEGAscript® kit (Ambion, Austin, TX, USA) following the manufacturer's instructions. The quality of dsRNA obtained was observed on an agarose gel and the quantity was measured using a NanoDrop (ND-1000, Thermo Scientific, Waltham, MA, USA).

For *in vitro* RNAi, dsRNAs (16 μg for each well) were directly added to the culture medium. A number of cells equal to 4 × 10⁶ cells in 2 ml were plated 24 h before silencing in a 6-well plate. Wells contained S2 cells and S2 cells stably expressing *TnBVank1*, respectively. The same procedure was performed in control cells, but no dsRNA was added. After 2 hours the medium containing the dsRNA was replaced with fresh medium without dsRNA and at different time points (T0, T6 and T12 h) 2 × 10⁶ cells were harvested for RNA extraction and 3.5 × 10⁵ cells for the post silencing caspase-3 assay.

For *in vivo* RNAi, 0.5 μg of both dsRNAs were mixed with 0.5 μg of pIZT/V5-His-ANK1 and 3 μl of transfection reagent (Lipofectin) and incubated for 20 min at room temperature before injection into the haemocoel of fifth instar *H. virescens* larvae (day one). Treated larvae were processed for qPCR analysis 56 h after the dsRNA administration.

Post silencing Alix and *TnBVank1* expression in S2 cells. In order to verify the Alix silencing, its transcript expression level was detected by qPCR and, on the same samples, the *TnBVank1* transcript level expression was determined. RNA was extracted at different time points from cells (T0, T6 and T10 h after adding dsRNA) using TRI Reagent (Sigma-Aldrich St Louis, MO, USA) and RNA quality was evaluated on an agarose gel. For qPCR, 1 μg of total RNA was treated with DNase (Deoxyribonuclease I Amplification Grade, Life Technologies, Carlsbad, CA, USA) and then it was reverse transcribed using Super Script™ III First-Strand Synthesis System

for RT-PCR (Life Technologies, Carlsbad, CA, USA, USA), with oligo-dT primers, following the manufacturer's protocol. qPCR was performed using GoTaq[®] qPCR Master Mix (Promega, Fitchburg, WI, USA) on an ABI Prism[®] 7500 Fast Real-Time PCR System Thermal Cycler (Applied Biosystem, USA). Sequence specific primers, each in a concentration of 0.3 μ M were:

AlixF 5'-CGCCCTGCAGAGCAACA-3'
 AlixR 5'-AGGGCACTGCCGCTGG-3'
 Ank1F 5'-ATCATGAGGGGCATGCATG-3'
 Ank1R 5'-AGCACCAGCGCTGTGAAGT-3'

As a reference gene, the gene coding for the *Drosophila* ribosomal protein L32 (RPL32, Accession Number P04359)⁵⁴ was used. Primers were:

Rpl32f 5'-TGCTAAGCTGTCGCACAAAT-3'
 Rpl32r 5'-GTTTCGATCCGTAACCGATG-3'

All qPCR analyses were carried out in triplicate. Quantification analysis of amplification was done using the comparative C_T (ΔC_T) method. The efficiencies of the two amplicons were approximately equal (Alix = 0.744509; Ank1 = 0.902369; Rpl32r = 0.841033).

Post silencing Alix and *TnBVank1* expression in haemocytes. In order to verify the Alix silencing, its transcript expression level was detected by qPCR also in haemocytes and on the same samples the *TnBVank1* transcript level expression was determined. RNA was extracted at 56 h after injection using TRI Reagent (Sigma-Aldrich St Louis, MO, USA), the quality and quantity were determined using NanoDrop spectrophotometer (ND-1000, Thermo Scientific, Waltham, MA, USA) and potential DNA contamination was eliminated with DNase (Deoxyribonuclease I Amplification Grade, Life Technologies, Carlsbad, CA, USA). cDNA was obtained as described above. qPCR was performed using GoTaq[®] qPCR Master Mix (Promega, Fitchburg, WI, USA) in a ABI Prism[®] 7500 Fast Real-Time PCR System Thermal Cycler (Applied Biosystem, USA). Sequence specific primers, each in a concentration of 0.3 μ M, were:

AlixF 5'-GTTGAACATCTTGGCACGGT-3'
 AlixR 5'-CAAAGCATGGACTCACGAGC-3'

TnBVank1 and reference gene primer sequences are reported above. All qPCR analyses were carried out in triplicate. Quantification analysis of amplification was done using the comparative C_T (ΔC_T) method. The efficiencies of the three amplicons were approximately equal (Alix = 0.931310; Ank1 = 0.898875; EF1 α = 0.874059; Rpl13 = 0.894415).

Statistical analysis. All data were presented as mean \pm SD of three independent biological replicates and were compared with analysis of variance (ANOVA) and Tukey's test using GraphPad Prism 6 software, La Jolla, California, USA.

References

- Pennacchio, F. & Strand, M. R. Evolution of developmental strategies in parasitic hymenoptera. *Annu. Rev. Entomol.* **51**, 233–258 (2006).
- Turnbull, M. & Webb, B. Perspectives on polydnavirus origins and evolution. *Adv. Virus Res.* **58**, 203–254 (2002).
- Strand, M. R. & Burke, G. R. Polydnaviruses: From discovery to current insights. *Virology*. **479–480**, 393–402 (2015).
- Webb, B. A. *et al.* Family Polydnaviridae in *Virus Taxonomy: Seventh Report of the International Committee on Taxonomy of Viruses* (ed. Van Regenmortel, M. H. V., Fauquet, C. M., Bishop, D. H. L.) 253–260 (Academic Press, 2000).
- Strand, M. R. & Burke, G. R. Polydnaviruses: nature's genetic engineers. *Annu. Rev. Virol.* **1**, 333–354 (2014).
- Falabella, P. *et al.* *Toxoneuron nigriceps* polydnavirus encodes a putative aspartyl protease highly expressed in parasitized host larvae. *Insect Mol. Biol.* **12**, 9–17 (2003).
- Falabella, P., Caccialupi, P., Varricchio, P., Malva, C. & Pennacchio, F. Protein tyrosine phosphatases of *Toxoneuron nigriceps* bracovirus as potential disrupters of host prothoracic gland function. *Arch. Insect Biochem. Physiol.* **61**(3), 157–169 (2006).
- Falabella, P. *et al.* Characterization of the κ B-like gene family in polydnaviruses associated with wasps belonging to different Braconid subfamilies. *J. Gen. Virol.* **88**, 92–104 (2007).
- Meng, X., Khanuja, B. S. & Ip, Y. T. Toll receptor-mediated *Drosophila* immune response requires Dif, an NF- κ B factor. *Genes Dev.* **13**, 792–797 (1999).
- Ghosh, S., May, M. J., Kopp, E. B. & NF- κ B, B. and Rel proteins: evolutionarily conserved mediators of immune responses. *Annu. Rev. Immunol.* **16**, 225–260 (1998).
- Ghosh, S. & Karin, M. Missing pieces in the NF- κ B puzzle. *Cell* **109**, 81–96 (2002).
- Webb, B.A., Strand, M.R. The biology and genomics of polydnaviruses. in *Comprehensive Molecular Insect Science* (ed Gilbert, L. I., Iatrou, K., Gill S. S.) **6**, 260–323 (Elsevier, 2005).
- Rogers, S., Wells, R. & Rechsteiner, M. Amino acid sequences common to rapidly degraded proteins: the PEST hypothesis. *Science* **234**, 364–368 (1986).
- Bitra, K., Suderman, R. J. & Strand, M. R. Polydnavirus Ank proteins bind NF- κ B homodimers and inhibit processing of Relish. *PLoS Pathog.* **8**(5), e1002722 (2012).
- Thoetkiattikul, H., Beck, M. H. & Strand, M. R. Inhibitor κ B-like proteins from a polydnavirus inhibit NF- κ B activation and suppress the insect immune response. *Proc. Nat. Acad. Sci. USA* **102**(32), 11426–11431 (2005).
- Ferrarese, R. *et al.* Early suppression of immune response in *Heliothis virescens* larvae by the endoparasitoid *Toxoneuron nigriceps*. *Inver. Surv. J.* **2**, 60–68 (2005).
- Beckage, N. E. & Gelman, D. B. Wasp parasitoid disruption of host development: implications for new biologically based strategies for insect control. *Annu. Rev. Entomol.* **49**, 299–330 (2004).

18. Lapointe, R. *et al.* Expression of a *Toxoneuron nigriceps* polydnavirus-encoded protein causes apoptosis-like programmed cell death in lepidopteran insect cells. *J. Gen. Virol.* **86**, 963–971 (2005).
19. Anitha, J., Pradeep, A. R. & Sivaprasad, V. Upregulation of Atg5 and AIF gene expression in synchronization with programmed cellular death events in integumental epithelium of *Bombyx mori* induced by a dipteran parasitoid infection. *Bull. Entomol. Res.* **104**(6), 794–800 (2014).
20. Li, M. *et al.* A Transcriptome Analysis Suggests Apoptosis-Related Signaling Pathways in Haemocytes of *Spodoptera Litura* After Parasitization by *Microplitis Bicoloratus*. *PLoS ONE*. **9**(10), e110967 (2014).
21. Missotten, M., Nichols, A., Rieger, K. & Sadoul, R. Alix, a novel mouse protein undergoing calcium-dependent interaction with the apoptosis-linked-gene 2 (ALG-2) protein. *Cell Death Differ.* **6**, 124–129 (1999).
22. Crawford, E. D. & Wells, J. A. Caspase substrates and cellular remodeling. *Annu. Rev. Biochem.* **80**, 1055–1087 (2011).
23. Brun, S. *et al.* The myb-related gene stonewall induces both hyperplasia and cell death in *Drosophila*: rescue of fly lethality by coexpression of apoptosis inducers. *Cell Death Differ.* **10**, 1752–62 (2006).
24. Nelo-Bazán, M. A. *et al.* Early growth response 1 (EGR-1) is a transcriptional regulator of mitochondrial carrier homolog 1 (MTC1)/presenilin 1-associated protein (PSAP). *Gene*. **578**(1), 52–62 (2016).
25. Yang, Y. *et al.* miR-15b-5p induces endoplasmic reticulum stress and apoptosis in human hepatocellular carcinoma, both *in vitro* and *in vivo*, by suppressing Rab1A. *Oncotarget*. **6**(18), 16227–38 (2015).
26. Dong, Z. MiR-223 modulates hepatocellular carcinoma cell proliferation through promoting apoptosis via the Rab1-mediated mTOR activation. *Biochem. Biophys. Res. Commun.* **483**(1), 630–637 (2017).
27. Pennacchio, F., Malva, C., Vinson, S.B. Regulation of host endocrine system by the endoparasitoid braconid *Cardiochiles nigriceps* and its polydnavirus. in *Endocrine Interactions of Insect Parasites and Pathogens* (ed. Edward, J. P., Weaver, R. J.) 123–132 (Oxford: BIOS, 2001).
28. Pennacchio, F., Falabella, P. & Vinson, S. B. Regulation of *Heliothis virescens* prothoracic glands by *Cardiochiles nigriceps* polydnavirus. *Arch. Insect. Biochem. Physiol.* **38**, 1–10 (1998).
29. Silverman, N. & Maniatis, T. NF- κ B signaling pathways in mammalian and insect innate immunity. *Genes Dev.* **15**, 2321–2342 (2001).
30. Espagne, E. *et al.* Genome sequence of a polydnavirus: insights into symbiotic virus evolution. *Science* **306**, 286–289 (2004).
31. Shi, M. *et al.* Characterization of a novel gene encoding ankyrin repeat domain from *Cotesia vestalis* polydnavirus (CvBV). *Virology* **375**, 374–382 (2008).
32. Strand, M. R. & Burke, G. R. Polydnaviruses as symbionts and gene delivery systems. *PLoS Pathog.* **8**(7), e1002757 (2012).
33. Gitatua, G. W., Gundersen-Rindal, D., Pedronib, M., Mbugic, P. J. & Dupas, S. Differential expression of the CrV1 haemocyte inactivation-associated polydnavirus gene in the African maize stem borer *Busseola fusca* (Fuller) parasitized by two biotypes of the endoparasitoid *Cotesia sesamiae* (Cameron). *J. Insect. Physiol.* **53**, 676–684 (2007).
34. Fath-Goodin, A., Kroemer, J. A. & Webb, B. A. The *Campoletis sonorensis* ichnovirus vankyrin protein P-vank-1 inhibits apoptosis in insect Sf9 cells. *Insect. Mol. Biol.* **18**, 497–506 (2009).
35. Magkrioti, C., Iatrou, K. & Labropoulou, V. Differential inhibition of BmRelish1-dependent transcription in lepidopteran cells by bracovirus ankyrin-repeat proteins. *Insect Biochem. Mol. Biol.* **41**, 993–1002 (2011).
36. Gueguen, G., Kalamarz, M., Ramroop, J., Uribe, J. & Govind, S. Polydnviral Ankyrin Proteins Aid Parasitic Wasp Survival by Coordinate and Selective Inhibition of Hematopoietic and Immune NF- κ B Signaling in Insect Hosts. *PLoS Pathog.* **9**(8), e1003580 (2013).
37. Revilla, Y. *et al.* Inhibition of Nuclear Factor B Activation by a Virus-encoded I κ B like Protein. *J. Biol. Chem.* **273**, 5405–5411 (1998).
38. Reuther, J. Y. & Baldwin, A. S. Jr. Apoptosis Promotes a Caspase-induced Amino-terminal Truncation of I κ B α That Functions as a Stable Inhibitor of NF- κ B. *J. Biol. Chem.* **274**, 20664–20670 (1999).
39. Suzuki, M. & Tanaka, T. Virus-like particles in venom of *Meteorus pulchricornis* induce host hemocyte apoptosis. *J. Insect Physiol.* **52**, 602–613 (2006).
40. Strand, M. R. & Pech, L. L. Microplitis demolitor polydnavirus induces apoptosis of a specific haemocyte morphotype in *Pseudaletia includens*. *J. Gen. Virol.* **76**, 283–291 (1995).
41. Odorizzi, G. The multiple personalities of Alix. *J. Cell. Sci.* **119**, 3025–3032 (2006).
42. Bongiovanni, A. *et al.* Alix protein is substrate of Ozz-E3 ligase and modulates actin remodeling in skeletal muscle. *J. Biol. Chem.* **287**(15), 12159–12171 (2012).
43. Valzania, L. *et al.* A Polydnavirus ANK Protein Acts as Virulence Factor by Disrupting the Function of Prothoracic Gland Steroidogenic Cells. *PLoS ONE* **9**(4), e95104 (2014).
44. Mahul-Mellier, A. L. *et al.* Alix and ALG-2 are involved in tumor necrosis factor receptor 1-induced cell death. *J. Biol. Chem.* **283**(50), 34954–34965 (2008).
45. Vinson, S. B., Guillot, F. S. & Hays, D. B. Rearing of *Cardiochiles nigriceps* in the laboratory, with *Heliothis virescens* as hosts. *Anti. Entomol. Soc. Am.* **66**, 1170–1172 (1973).
46. Webb, B. A. & Dahlman, D. L. Developmental pathology of *Heliothis virescens* larvae parasitized by *Microplitis croceipes*: Parasite-mediated host developmental arrest. *Arch. Insect Biochem. Phys.* **2**(2), 131–143 (1985).
47. Pennacchio, F., Vinson, S. B. & Tremblay, E. Host regulation effects of *Heliothis virescens* (F.) larvae induced by teratocytes of *Cardiochiles nigriceps* Viereck (Lepidoptera, Noctuidae Hymenoptera, Braconidae). *Arch. Insect. Biochem. Physiol.* **19**, 177–192 (1992).
48. Duchi, S. *et al.* The impact on microtubule network of a bracovirus I κ B-like protein. *Cell. Mol. Life Sci.* **67**, 1699–1712 (2010).
49. Bustin, S. A. *et al.* The MIQE guidelines: minimum information for publication of quantitative real-time PCR Experiments. *Clin. Chem.* **55**, 611–622 (2009).
50. Sordet, O., Khan, Q. A., Kohn, K. W. & Pommier, Y. Apoptosis induced by topoisomerase inhibitors. *Curr. Med. Chem. Anti-Canc. Agents.* **3**, 271–90 (2003).
51. Pommier, Y. *et al.* Repair of and checkpoint response to topoisomerase I-mediated DNA damage. *Mutat. Res.* **532**, 173–203 (2003).
52. Bradford, M. M. A rapid and sensitive for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248–254 (1976).
53. Tsuda, M., Seong, K. H. & Aigaki, T. POSH, a scaffold protein for JNK signaling, binds to ALG-2 and ALIX in *Drosophila*. *FEBS Lett.* **580**, 3296–3300 (2006).
54. Ponton, F., Chapuis, M. P., Pernice, M., Sword, G. A. & Simpson, S. J. Evaluation of potential reference genes for reverse transcription-qPCR studies of physiological responses in *Drosophila melanogaster*. *J. Insect Physiol.* **57**, 840–850 (2011).

Acknowledgements

This work was supported by University of Basilicata (RILfunds). We thank Tab Consulting s.r.l. for financing a Ph.D grant. We also thank Dr. Toshiro Aigaki for kindly giving us anti-Alix antibody, Dr. Mario Lippolis, Dr. Luciana Possidente, Giovanni Petruolo and Francesco Petruolo for their technical support in immunofluorescence experiments and Dr. Giuseppe Cugno and Dr. Pellegrino Musto. This manuscript is in memory of Dr. Franco Graziani of the Institute of Genetics and Biophysics - CNR (Naples, Italy).

Author Contributions

P.F. designed the experiments, wrote and critically revised the paper; M.C., H.V., P.P., S.B.V., D.N. and S.A.B. contributed to the data interpretation and critically revised the paper; R.S. carried out experimental work and analysed data. A.A. and C.G. performed functional proteomic experiments. C.S. collected RNA samples. M.N. and R.S. performed coimmunolocalisation experiments. G.G. and R.S. performed qPCR. R.S., I.L. and V.R. performed flow cytometry analysis. All authors read and approved the final manuscript.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-017-11939-x>.

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017