



**Syddansk Universitet**

## **Long-term safety of icatibant treatment of patients with angioedema in real-world clinical practice**

Zanichelli, A; Maurer, Marcus; Aberer, W; Caballero, T; Longhurst, H J; Bouillet, L; Fabien, V; Andresen, I; IOS Study Group; Bygum, Anette

*Published in:*  
Allergy

*DOI:*  
[10.1111/all.13103](https://doi.org/10.1111/all.13103)

*Publication date:*  
2017

*Document version*  
Publisher's PDF, also known as Version of record

*Document license*  
CC BY-NC

*Citation for published version (APA):*  
Zanichelli, A., Maurer, M., Aberer, W., Caballero, T., Longhurst, H. J., Bouillet, L., ... Bygum, A. (2017). Long-term safety of icatibant treatment of patients with angioedema in real-world clinical practice. *Allergy*, 72(6), 994-998. DOI: 10.1111/all.13103

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## BRIEF COMMUNICATION

# Long-term safety of icatibant treatment of patients with angioedema in real-world clinical practice

A. Zanichelli<sup>1</sup>, M. Maurer<sup>2</sup>, W. Aberer<sup>3</sup>, T. Caballero<sup>4</sup>, H. J. Longhurst<sup>5</sup>, L. Bouillet<sup>6</sup>, V. Fabien<sup>7</sup> and I. Andresen<sup>8</sup> for the IOS Study Group

<sup>1</sup>Department of Biomedical and Clinical Sciences Luigi Sacco, University of Milan, ASST Fatebenefratelli Sacco, Milan, Italy; <sup>2</sup>Department of Dermatology and Allergy, Charité–Universitätsmedizin Berlin, Berlin, Germany; <sup>3</sup>Department of Dermatology and Venerology, Medical University of Graz, Graz, Austria; <sup>4</sup>Department of Allergy, Hospital La Paz Institute for Health Research (IdiPaz), Biomedical Research Network on Rare Diseases (CIBERER, U754), Madrid, Spain; <sup>5</sup>Department of Immunology, Barts Health NHS Trust, London, UK; <sup>6</sup>National Reference Centre for Angioedema, Internal Medicine Department, Grenoble University Hospital, Grenoble, France; <sup>7</sup>Shire, Zug, Switzerland at time of data analysis; now with Vifor Pharma, Glattbrugg, Switzerland; <sup>8</sup>Shire, Zug, Switzerland

**To cite this article:** Zanichelli A, Maurer M, Aberer W, Caballero T, Longhurst HJ, Bouillet L, Fabien V, Andresen I. Long-term safety of icatibant treatment of patients with angioedema in real-world clinical practice. *Allergy* 2017; **72**: 994–998.

## Keywords

angioedema; icatibant; real-world; safety.

## Correspondence

Dr. Andrea Zanichelli, Department of Biomedical and Clinical Sciences Luigi Sacco, University of Milan, Luigi Sacco Hospital Milan, Via G.B. Grassi 74, 20157 Milan, Italy.  
Tel.: +39 0239042324  
Fax: +39 0250319828  
E-mail: andrea.zanichelli@unimi.it

IOS Study Group members are in Acknowledgments.

Accepted for publication 30 November 2016

DOI:10.1111/all.13103

Edited by: Stephan Weidinger

Icatibant has demonstrated tolerability and efficacy in patients with hereditary angioedema with C1 inhibitor deficiency (C1-INH-HAE) type I/II (1–5). Icatibant phase 3 studies included around 200 patients (228 attacks) and follow-up periods  $\leq 24$  weeks (1, 2). No icatibant-related serious adverse events (SAEs) were reported (1, 2). Nevertheless, considering the limited number of patients enrolled and the short follow-up, additional data are desirable.

Patient registries provide useful sources of additional data on rare conditions. The Icatibant Outcome Survey (IOS; NCT01034969) is an ongoing, international, prospective, observational registry monitoring efficacy and safety of icatibant in the real-world setting. Preclinical studies have

## Abstract

The Icatibant Outcome Survey (IOS) is an observational study monitoring safety and effectiveness of icatibant in the real-world setting. We analyzed safety data from 3025 icatibant-treated attacks in 557 patients (enrolled between July 2009 and February 2015). Icatibant was generally well tolerated. Excluding off-label use and pregnancy, 438 patients (78.6%) did not report adverse events (AEs). The remaining 119 (21.4%) patients reported 341 AEs, primarily gastrointestinal disorders (19.6%). Of these, 43 AEs in 17 patients (3.1%) were related to icatibant. Serious AEs (SAEs) occurred infrequently. A total of 143 SAEs occurred in 59 (10.6%) patients; only three events (drug inefficacy, gastritis, and reflux esophagitis) in two patients were considered related to icatibant. Notably, no SAEs related to icatibant occurred in patients with cardiovascular disease, nor in those using icatibant at a frequency above label guidelines. Additionally, no major differences were noted in AEs occurring in on-label vs off-label icatibant users.

suggested that icatibant-associated adverse events (AEs) may relate to its mechanism of action (6, 7). The current analysis evaluated incidence of cardiovascular-related AEs, as well as other common AEs in icatibant-treated patients. The objective was to assess safety data from the IOS database.

## Patients and methods

Eligible patients were currently receiving or were candidates for icatibant use. Data were collected by physicians completing electronic forms with information from patients' follow-up visits; the protocol recommended every 6-month visits, but visit frequency was not mandated. IOS is conducted per

local ethics committees and/or health authorities at participating sites, the Declaration of Helsinki and the International Conference on Harmonisation Good Clinical Practice guidelines, and all patients provided written informed consent prior to participation. Additional details regarding the IOS registry have been previously described (8).

Safety data from the IOS reported herein were collected between July 2009 and February 2015 from 47 participating centers in 11 countries. Adverse events were categorized per Medical Dictionary for Regulatory Activities (MedDRA) system organ classification and analyzed by number/percentage of patients reporting a safety event and number/percentage of the occurrence of any safety event. The MedDRA term 'general disorders and administration site conditions' encompassed the following: application site pain, chest discomfort, chills, drug ineffective, drug intolerance, edema, fatigue, feeling hot, gait disturbance, hyperhidrosis, hyperplasia, influenza-like illness, infusion site pain, injection site erythema, injection site pain, injection site urticaria, localized edema, local swelling, malaise, noncardiac chest pain, pain, peripheral edema, pyrexia, and therapeutic product ineffective. Although pregnancy and off-label use (defined as patients with angioedema other than HAE type I/II or occurrence of AEs in patients <18 years) were documented as AEs in the IOS registry from a regulatory standpoint, these events were not considered clinically meaningful AEs and therefore were analyzed separately from our safety analysis.

## Results

### Patients

As of February 2015,  $\geq 1$  dose of icatibant was used to treat 3025 attacks in 557 patients. Mean age [standard deviation (SD)] at enrollment was 41.6 (15.2) years, with a female/male gender distribution of 63.6/36.4%.

Of patients with documented ethnicity ( $n = 540$ ), most were white (93.3%). A total of 378 (69.9%) and 23 (4.3%) patients had C1-INH-HAE type I/II, respectively; the remaining patients ( $n = 140$ , 25.9%) had HAE with normal C1-INH, angiotensin converting enzyme (ACE) inhibitor-induced angioedema, idiopathic angioedema, acquired angioedema (CI-INH-AAE), or other (missing diagnoses,  $n = 16$ ). Demographic parameters varied across HAE types. For instance, mean (SD) age at diagnosis was lower for patients with C1-INH-HAE type I/II [24.0 (16.0) years] and HAE with normal C1-INH [36.0 (14.3) years] vs those with idiopathic angioedema [44.7 (13.7) years], CI-INH-AAE [59.2 (9.7) years], and ACE inhibitor-induced angioedema [73.2 (6.1) years]. However, patients with CI-INH-AAE and ACE inhibitor-induced angioedema had a shorter mean (SD) delay between symptom onset and diagnosis [1.0 (1.2) and 1.7 (2.6) years, respectively] vs those with idiopathic angioedema [8.4 (10.8) years], C1-INH-HAE type I/II [10.1 (13.5) years], or HAE with normal C1-INH [11.1 (11.3) years].

Since entry into the IOS, the overall patient population was followed for a mean (SD) of 2.97 (1.42) years (<1 year,

10.2%; 1–2 years, 19.4%; 2–3 years, 17.2%; 3–4 years, 24.1%; 4–5 years, 26%, and  $\geq 5$  years, 3.1%). Follow-up times were similar across HAE subtypes. In total, patients were followed for 1655.2 icatibant-treated patient years.

### AEs overall

Most icatibant-treated patients ( $n = 438$ , 78.6%) did not report an AE; the remaining 119 (21.4%) reported 341 AEs. The most commonly reported events were gastrointestinal (GI) disorders (19.6%), general disorders and administration site conditions (13.8%), infections and infestations (11.1%), respiratory, thoracic, and mediastinal disorders (8.8%), musculoskeletal and connective tissue disorders (6.5%), and skin and subcutaneous tissue disorders (6.5%). All other AEs represented <5% of total events. No major differences were noted in AEs occurring in on-label vs off-label icatibant users.

### AEs related to icatibant

A total of 17 patients (3.1%) reported 43 treatment-related events (Table 1). Of these, 19 events (44.2%) in five patients were considered possibly related, and 24 events (55.8%) in 12 patients probably related. The most common treatment-related AEs were general disorders and administration site conditions (53.5%), GI disorders (11.6%), investigations (weight or blood pressure decreases, 11.6%), vascular disorders (hyperemia, 9.3%), and nervous system disorders (7.0%).

### Serious adverse events

A total of 59 patients (10.6%) reported 143 SAEs (Table 2). The most common events were GI disorders (28%), respiratory, thoracic, and mediastinal disorders (12.6%), neoplasms (benign, malignant, or unspecified, 8.4%), general disorders and administration site conditions (7.7%), and infections and infestations (7.7%). Three SAEs (drug inefficacy, gastritis, reflux oesophagitis) in two patients were considered possibly or probably related to icatibant.

### AEs in patients with cardiovascular disease

Cardiovascular disease (CVD) was reported in 95 icatibant-treated patients. At time of enrollment, patients reported: hypertension ( $n = 80$ ), transient ischemic attack (TIA) or stroke ( $n = 5$ ), angina ( $n = 4$ ), and ischemic heart disease ( $n = 3$ ). At follow-up, the following new onset CVDs were reported: hypertension ( $n = 5$ ), angina ( $n = 1$ ), ischemic heart disease ( $n = 1$ ), and TIA or stroke ( $n = 1$ ). Additionally at follow-up, hypertension was reported in three patients whose CVD history was missing at baseline.

A total of 22 (23.2%) patients with CVD reported 49 AEs, most commonly general disorders and administration site conditions (eight events, 16.3%), respiratory, thoracic, and mediastinal disorders (seven events, 14.3%), and GI disorders (five events, 10.2%). Of AEs occurring in patients with CVD, 13 events in three patients were considered possibly or

**Table 1** AEs (excluding off-label use\* and pregnancy) considered by the investigator to be possibly or probably related to icodebant

	Icodebant-treated patients (n = 557)	
	Number of events related to icodebant use (%)	Number of patients experiencing icodebant-related events (%)
Any event	43 (100.0)	17 (3.1)
Drug ineffective	6 (14.0)	5 (0.9)
Injection site erythema	6 (14.0)	1 (0.2)
Blood pressure decreased	4 (9.3)	1 (0.2)
Hyperemia	4 (9.3)	3 (0.5)
Pain	3 (7.0)	2 (0.4)
Gastritis	2 (4.7)	1 (0.2)
Application site pain	1 (2.3)	1 (0.2)
Chest discomfort	1 (2.3)	1 (0.2)
Cholelithiasis	1 (2.3)	1 (0.2)
Depression	1 (2.3)	1 (0.2)
Dizziness	1 (2.3)	1 (0.2)
Epigastric discomfort	1 (2.3)	1 (0.2)
Feeling hot	1 (2.3)	1 (0.2)
Headache	1 (2.3)	1 (0.2)
Herpes zoster	1 (2.3)	1 (0.2)
Infusion site pain	1 (2.3)	1 (0.2)
Injection site pain	1 (2.3)	1 (0.2)
Injection site urticaria	1 (2.3)	1 (0.2)
Nausea	1 (2.3)	1 (0.2)
Noncardiac chest pain	1 (2.3)	1 (0.2)
Therapeutic product ineffective	1 (2.3)	1 (0.2)
Postherpetic neuralgia	1 (2.3)	1 (0.2)
Reflux oesophagitis	1 (2.3)	1 (0.2)
Weight decreased	1 (2.3)	1 (0.2)

AEs, adverse events.

\*Off-label use refers to patients with angioedema other than HAE type I or II or those in whom AEs occurred prior to 18 years of age.

probably related to icodebant, with the majority being general disorders and administration site conditions (eight events, 61.5%). Sixteen patients reported 24 SAEs, none of which were considered related to icodebant.

#### AEs in patients who used icodebant at a frequency above label guidelines

Per the icodebant summary of product characteristics,  $\leq 8$  icodebant injections per month have been administered in clinical trials (9). In the IOS registry, of 10 patients who received  $\geq 9$  icodebant injections in 1 month, drug ineffectiveness was reported in two patients. The icodebant package information recommends an interval of  $\geq 6$  h between doses and no more than three doses in 24 h (10). In the IOS registry, no AEs were reported by 13 patients whose intervals between icodebant doses were  $< 6$  h, nor in four patients who received icodebant more frequently than three times in 24 h.

**Table 2** SAEs (excluding off-label\* use and pregnancy) occurring with a frequency of  $> 1\%$  of total events

	Icodebant-treated patients (n = 557)	
	Number of events (%)	Number of patients (%)
Any event	143 (100.0)	59 (10.6)
Abdominal pain	8 (5.6)	7 (1.3)
Angioedema	4 (2.8)	4 (0.7)
Diarrhea	4 (2.8)	2 (0.4)
Peripheral edema	4 (2.8)	2 (0.4)
Abdominal distension	3 (2.1)	3 (0.5)
Face swelling	3 (2.1)	3 (0.5)
Laryngeal edema	3 (2.1)	3 (0.5)
Abdominal hernia	2 (1.4)	2 (0.4)
Abdominal pain (upper)	2 (1.4)	2 (0.4)
Colonic polyp	2 (1.4)	1 (0.2)
Cough	2 (1.4)	2 (0.4)
Dyspnoea	2 (1.4)	2 (0.4)
Hematochezia	2 (1.4)	1 (0.2)
Hereditary angioedema	2 (1.4)	2 (0.4)
Legionella infection	2 (1.4)	1 (0.2)
Local administration site swelling	2 (1.4)	2 (0.4)
Myelodysplastic syndrome	2 (1.4)	1 (0.2)
Pyrexia	2 (1.4)	2 (0.4)
Respiratory failure	2 (1.4)	1 (0.2)
Suicide attempt	2 (1.4)	2 (0.4)

SAEs, serious adverse events.

\*Off-label use refers to patients with angioedema other than HAE type I or II or those in whom adverse events occurred prior to 18 years of age.

#### Discussion

Overall, no unexpected safety signals emerged in this large database, and AEs were similar in off-label vs on-label users. General disorders and administration site conditions occurred in 13.8% of icodebant-treated patients, including but not limited to local injection site reactions. Occurrence of these reactions was notably lower than their 97% incidence in icodebant clinical trials (10), especially since the 13.8% incidence may be an overestimation (given that these reactions were recorded as part of a larger bucket of events). This discrepancy may reflect multiple factors, including differences in the processes followed for reporting and documenting such reactions in a real-world setting vs a controlled environment. Whereas in a clinical trial setting investigators proactively document occurrence of all (including very minor and transient) injection site reactions in a detailed manner upon their immediate occurrence, in a drug registry setting, such reactions are documented retrospectively, possibly leading to recall bias. Patients may fail to report a well-tolerated local reaction, or the investigators may choose not to document this expected, minor reaction as an AE. The extended time lapse between occurrences of angioedema attacks and when patients report AEs during follow-up visits may further

contribute to underreporting. Nonetheless, our findings present a real-world perspective with regard to icatibant tolerability, which may indeed differ somewhat from findings in clinical trials.

Serious adverse events occurred infrequently, and none were cardiovascular-related. Studies have shown that bradykinin B2 receptor blockade might impair endothelial repair after an acute cardiovascular event (6, 7). However, icatibant has a short half-life (9) and the effect of intermittent treatments is likely transient.

Potential limitations of the IOS study were the uncontrolled clinical environment inherent in an observational study design, as well as possible inconsistencies in patient and physician-based reporting of AEs across registry study sites. Nonetheless, observational registries offer valuable insights into the safety and efficacy of treatments in real-world settings.

Our results show that icatibant was generally well tolerated; incidence and severity of AEs was not greater than was expected in this patient population. No major differences were noted in on-label *vs* off-label icatibant users, and no SAEs occurred in patients with CVD using icatibant.

### Acknowledgments

The authors would like to thank the IOS investigators for their contributions:

*Austria:* W. Aberer; *Brazil:* A. S. Grumach; *Denmark:* A. Bygum; *France:* C. Blanchard Delaunay, I. Boccon-Gibod, L. Bouillet, B. Coppere, A. Du Thanh, C. Dzviga, O. Fain, B. Goichot, A. Gompel, S. Guez, P. Jeandel, G. Kanny, D. Launay, H. Maillard, L. Martin, A. Masseur, Y. Ollivier, A. Sobel; *Germany:* E. Aygören-Pürsün, M. Baş, M. Bauer, K. Bork, J. Greve, M. Magerl, I. Martinez Sauer, M. Maurer; U. Strassen; *Greece:* E. Papadopoulou-Alataki, F. Psarros; *Israel:* Y. Graif, S. Kivity, A. Reshef, E. Toubi; *Italy:* F. Arcoletto, M. Bova, M. Cicardi, P. Manconi, G. Marone, V. Montinaro, M. Triggiani, A. Zanichelli; *Spain:* M. L. Baeza, T. Caballero, R. Cabañas, G. Gala Ortiz, M. Guilarte, D. Hernandez, C. Hernando de Larramendi, R. Leonart, T. Lobera, L. Marques, B. Saenz de San Pedro; *Sweden:* J. Björkander; and *United Kingdom:* C. Bethune, T. Garcez, H. J. Longhurst.

### Author contributions

Andrea Zanichelli contributed to study conception and design, data acquisition, analysis and interpretation, drafting the manuscript and critical content revisions, as well as final content approval for submission; Marcus Maurer contributed to study conception and design, data acquisition, analysis and interpretation, drafting the manuscript and critical content revisions, as well as final content approval for submission; Werner Aberer contributed to data acquisition, analysis and interpretation, drafting the manuscript and critical content revisions, as well as final content approval for submission; Teresa Caballero contributed to study conception and design, data acquisition, analysis and interpretation, drafting

the manuscript and critical content revisions, as well as final content approval for submission; Hilary J. Longhurst contributed to data acquisition, analysis and interpretation, drafting the manuscript and critical content revisions, as well as final content approval for submission; Laurence Bouillet contributed to data acquisition, drafting the manuscript and critical content revisions, as well as final content approval for submission; Vincent Fabien contributed to study conception and design, analysis and interpretation of data, drafting the manuscript and critical content revisions, as well as final content approval for submission; and Irmgard Andresen contributed to study conception and design, analysis and interpretation of data, drafting the manuscript and critical content revisions, as well as final content approval for submission.

### Funding

Research was funded by Shire Development, LLC. Under the direction of the authors, Linda Wagner, PharmD, and Sophia Shumyatsky, PharmD, employees of Excel Scientific Solutions, provided writing assistance for this publication. Editorial assistance in formatting, proofreading, copyediting, and fact checking also was provided by Excel Scientific Solutions. Shire Development, LLC, provided funding to Excel Scientific Solutions for support in writing and editing this manuscript. The interpretation of the data was made by the authors independently.

### Conflicts of interest

Dr Andrea Zanichelli has received speaker fees from CSL Behring, Shire, and Sobi; consultancy fees from CSL Behring and Shire; and has acted on the medical/advisory boards for CSL Behring and Shire. Prof Marcus Maurer has received research grant support and/or speaker/consultancy fees from BioCryst, CSL Behring, Dyax, and Shire/Jerini AG. Prof Werner Aberer has acted as a medical advisor and speaker for BioCryst, CSL Behring, Pharming, and Shire, and has received funding to attend conferences/educational events, and donations to his departmental fund from and participated in clinical trials for Shire. Dr. Teresa Caballero has received speaker fees from CSL Behring, GlaxoSmithKline, MSD, Novartis, and Shire; consultancy fees from BioCryst, CSL Behring, Novartis, Shire, and Sobi; funding for travel and meeting attendance from CSL Behring, Novartis, and Shire, and has participated in clinical trials/registries for CSL Behring, Dyax, Novartis, Pharming, and Shire. She is a researcher from the IdiPaz Program for promoting research activities. Dr Hilary J. Longhurst has received research grant support and/or speaker/consultancy fees from BioCryst, CSL Behring, Dyax, Shire, and SOBI-Biovitrum. Prof Laurence Bouillet has received honoraria from BioCryst, CSL Behring, Novartis, Pharming, and Shire, and her institute has received research funding from CSL Behring, GlaxoSmithKline, Novartis, Roche, and Shire. Dr Irmgard Andresen is a full-time employee of Shire, Zug, Switzerland. Vincent Fabien was a Shire employee at time of analysis.

## References

1. Cicardi M, Banerji A, Bracho F, Malbrán A, Rosenkranz B, Riedl M et al. Icatibant, a new bradykinin-receptor antagonist, in hereditary angioedema. *N Engl J Med* 2010;**363**:532–541.
2. Lumry WR, Li HH, Levy RJ, Potter PC, Farkas H, Moldovan D et al. Randomized placebo-controlled trial of the bradykinin B2 receptor antagonist icatibant for the treatment of acute attacks of hereditary angioedema: the FAST-3 trial. *Ann Allergy Asthma Immunol* 2011;**107**:529–537.
3. Malbrán A, Riedl M, Ritchie B, Smith WB, Yang W, Banerji A et al. Repeat treatment of acute hereditary angioedema attacks with open-label icatibant in the FAST-1 trial. *Clin Exp Immunol* 2014;**177**:544–553.
4. Aberer W, Maurer M, Reshef A, Longhurst H, Kivity S, Bygum A et al. Open-label, multicenter study of self-administered icatibant for attacks of hereditary angioedema. *Allergy* 2014;**69**:305–314.
5. Bas M, Greve J, Hoffmann TK, Reshef A, Aberer W, Maurer M et al. Repeat treatment with icatibant for multiple hereditary angioedema attacks: FAST-2 open-label study. *Allergy* 2013;**68**:1452–1459.
6. Kränkel N, Kuschnerus K, Müller M, Speer T, Mocharla P, Madeddu P et al. Novel insights into the critical role of bradykinin and the kinin B2 receptor for vascular recruitment of circulating endothelial repair-promoting mononuclear cell subsets: alterations in patients with coronary disease. *Circulation* 2013;**127**:594–603.
7. Yao Y, Sheng Z, Li Y, Yan F, Fu C, Ma G et al. Tissue kallikrein promotes cardiac neovascularization by enhancing endothelial progenitor cell functional capacity. *Hum Gene Ther* 2012;**23**:859–870.
8. Maurer M, Aberer W, Bouillet L, Caballero T, Fabien V, Kanny G et al. Hereditary angioedema attacks resolve faster and are shorter after early icatibant treatment. *PLoS One* 2013;**8**:e53773.
9. Firazyri<sup>®</sup> (icatibant) summary of product characteristics. Available at: [http://www.eur.europa.eu/docs/en\\_GB/document\\_library/EPAR\\_-\\_Product\\_Information/human/000899/WC500022966.pdf](http://www.eur.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/000899/WC500022966.pdf) (accessed 9 September 2016).
10. Firazyri<sup>®</sup> (icatibant) [package insert]. Shire Orphan Therapies, LLC: Lexington, MA; 2015.