

Existence of Equilibrium for Integer Allocation Problems

Somdeb Lahiri

Institute for Financial Management and Research

24, Kothari Road, Nungambakkam,

Chennai- 600 034,

Tamil Nadu, India.

Email: lahiri@ifmr.ac.in

First version: December 24, 2005

This version: January 3, 2005.

Abstract

In this paper we show that if all agents are equipped with discrete concave production functions, then a feasible price allocation pair is a market equilibrium if and only if it solves a linear programming problem, similar to, but perhaps simpler than the one invoked in Yang (2001).

Using this result, but assuming discrete concave production functions for the agents once again, we are able to show that the necessary and sufficient condition for the existence of market equilibrium available in Sun and Yang (2004), which involved obtaining a price vector that satisfied infinitely many inequalities, can be reduced to one where such a price vector satisfies finitely many inequalities. A necessary and sufficient condition for the existence of a market equilibrium when the maximum value function is Weakly Monotonic at the initial endowment that follows from our results is that the maximum value function is partially concave at the initial endowment.

1. Introduction: The equilibrium existence problem with indivisibilities has been investigated by Yang (2001) and more recently by Sun and Yang (2004). The model they consider is the model of a market game due to Shapley and Shubik (1969, 1976), with the goods being available for redistribution among the agents being available in integer units only. Yang (2004), shows that a constrained market equilibrium (i.e. a market clearing allocation, where each agent is constrained to maximize profit subject to its consumption not exceeding the initial endowment of the goods), exists if and only if there is a feasible price allocation pair that solves a certain linear programming problem. The result is obtained without any concavity assumption being imposed on the production function of the agents.

In this paper we show that if all agents are equipped with discrete concave production function, then a feasible price allocation pair is a market equilibrium (i.e. where agents solve an unconstrained profit maximization problem at given prices to obtain the resulting market clearing allocation) if and only if it solves a linear programming problem, similar to, but perhaps simpler than the one invoked in Yang (2001).

Using this result, but assuming discrete concave production function for the agents once again, we are able to show that the necessary and sufficient condition for the existence of market equilibrium available in Sun and Yang (2004), which involved obtaining a price

vector that satisfied infinitely many inequalities, can be reduced to one where such a price vector satisfies finitely many such inequalities.

These results provide the necessary computational device for obtaining a market equilibrium for integer allocation problems.

2. The Model: We now develop the general equilibrium model for the case where the inputs are available in integer amount only.

Let $N = \mathbb{N} \cup \{0\}$, where \mathbb{N} denotes the set of natural numbers. Let there be $H > 0$ agents and $L+1 > 1$ commodities. The first L commodities are used as inputs to produce the $L+1^{\text{th}}$ commodity, which is a numeraire consumption good. Let $w \in N^L$ denote the aggregate initial endowment of the inputs which is available for distribution among the agents.

For $j = 1, \dots, L$, let w_j denote the aggregate amount of commodity j that is initially available in the economy.

A function $f: N^L \rightarrow \mathcal{R}$ is said to be discrete concave if there exists a continuous concave function $g: \mathcal{R}_+^L \rightarrow \mathcal{R}$ such that the restriction of g to N^L coincides with f .

Given functions $f: N^L \rightarrow \mathcal{R}$ and $g: \mathcal{R}_+^L \rightarrow \mathcal{R}$, let $\text{graph}(f) \equiv \{(x, \alpha) \in N^L \times \mathcal{R} / \alpha \leq f(x)\}$ and $\text{graph}(g) \equiv \{(x, \alpha) \in \mathcal{R}_+^L \times \mathcal{R} / \alpha \leq g(x)\}$.

Given a function $f: N^L \rightarrow \mathcal{R}$ its canonical extension is the function $g^f: \mathcal{R}_+^L \rightarrow \mathcal{R}$ such that the $\text{graph}(g^f) = \text{convex hull of graph}(f)$. Clearly g^f is continuous and concave.

Let e denote the vector in \mathcal{R}^L all whose coordinates are equal to one and for $j = 1, \dots, L$, let e^j denote the vector in \mathcal{R}^L whose j^{th} coordinate is equal to one and all other coordinates are equal to zero.

If f is discrete concave, then the restriction of its canonical extension g^f to N^L coincides with f .

Note: For $x \in N^L$ and $x' \in \mathcal{R}^L$ with $x \leq x' \leq x + e$, $g^f(x') \in \text{convex hull of } \{f(x + \sum_{j=1}^L k^j e^j) / k^j \in \{0, 1\} \text{ for } j = 1, \dots, L\}$.

In fact, if we let $N(x')$ denote $\{y \in N^L / x \leq y \leq x + e\}$, and Δ' denote the $|N(x')| - 1$ simplex in $\mathcal{R}^{N(x')}$, then $g^f(x') = \text{Max} \{ \sum_{k=1}^K t^k f(y^k) / K$ is a positive integer, $y^k \in N(x')$ for $k = 1, \dots, K$,

$x' = \sum_{k=1}^K t^k y^k$, and the array $t = \langle t^k / k = 1, \dots, K \rangle \in \Delta' \}$.

Each agent i has preferences defined over N^L which is represented by a discrete concave production function f^i .

The pair $\langle \{f^i / i = 1, \dots, H\}, w \rangle$ is called an integer allocation problem and is assumed to be a given for the rest of the analysis.

An input consumption vector of agent i is denoted by a vector $X^i \in N^L$.

A price vector p is an element of $\mathcal{R}_+^L \setminus \{0\}$, where for $j = 1, \dots, L$, p_j denotes the price of input j .

At a price vector p , the objective of agent i is to maximize profits:

Maximize $[f^i(X^i) - p^T X^i]$

An allocation is an array $X = \langle X^i / i = 1, \dots, H \rangle$ such that $X^i \in N^L$ for all $i = 1, \dots, H$.

Given $x \in \mathbb{N}^L$, let $F(x) = \{ X = \langle X^i / i = 1, \dots, H \rangle / X \text{ is an allocation satisfying } \sum_{i=1}^H X^i = x \}$.

An allocation X is said to be feasible if $X \in F(w)$.

A market equilibrium is a pair $\langle p^*, X^* \rangle$ where p^* is a price vector, X^* is a feasible allocation and for all $i = 1, \dots, H$, X^{*i} maximizes profits for agent i .

The function $V: \mathbb{N}^L \rightarrow \mathbb{R}_+$ such that for all $x \in \mathbb{N}^L$: $V(x) = \text{Max} \{ \sum_{i=1}^H f^i(X^i) / X = \langle X^i / i = 1, \dots, H \rangle \in F(x) \}$, is called the maximum value function.

A feasible allocation $X^* = \langle X^{*i} / i = 1, \dots, H \rangle$ is said to be efficient if $\sum_{i=1}^H f^i(X^{*i}) = V(w)$.

The following result is available in Lahiri (2005) and will be used in the sequel.

Proposition 1: Let $\langle p^*, X^* \rangle$ be a market equilibrium. Then X^* is an efficient allocation.

A constrained market equilibrium is a pair $\langle p^*, X^* \rangle$ where p^* is a price vector, X^* is a feasible allocation such that for all $i = 1, \dots, H$, X^{*i} solves:

$$\text{Maximize } [f^i(X^i) - p^T X^i]$$

Subject to $X^i \leq w$.

Clearly a market equilibrium is a constrained market equilibrium. The following example shows that a constrained market equilibrium need not be a market equilibrium.

Example 1: Let $H = 1$, $L = 1$, $f^1(x) = x$ for all $x \in \mathbb{N}$ and $w = 1$. Let $X^{*1} = 1$ and $X^* = \langle X^{*1} \rangle$. For all $p \in (0, 1]$, $\langle p, X^* \rangle$ is a constrained market equilibrium, though for $p \in (0, 1)$, $\langle p, X^* \rangle$ is never a market equilibrium. $\langle 1, X^* \rangle$ is the unique market equilibrium for this integer allocation problem.

3. Existence of Market Equilibrium: This section is devoted to obtaining results pertaining to the existence of market equilibrium for the given integer allocation problem. We begin with a lemma.

Lemma 1: Let $f: \mathbb{N}^L \rightarrow \mathbb{R}$ be a discrete concave function with g^f being its canonical extension and let $p \in \mathbb{R}^L$. Let $x \in \mathbb{N}^L$ and $x' \in \mathbb{R}_+^L$ with $x' \leq x$. Then $g^f(x') + p^T x' \leq \text{Max} \{ f(y') + p^T y' / y' \in \mathbb{N}^L \text{ and } y' \leq x \}$.

Proof: Since $x' \leq x$, there exists $y \in \mathbb{N}^L$ such that $y \leq x \leq y + e$. Thus, $g^f(x') \in \text{convex hull of } \{ f(y + \sum_{j=1}^L k^j e^j) / k^j \in \{0, 1\} \text{ for } j = 1, \dots, L \}$. This implies that $g^f(x') + p^T x' \in \text{convex hull of } \{ f(y + \sum_{j=1}^L k^j e^j) + p^T (y + \sum_{j=1}^L k^j e^j) / k^j \in \{0, 1\} \text{ for } j = 1, \dots, L \}$. Hence, $g^f(x') + p^T x' \leq$

Max $\{ f(y + \sum_{j=1}^L k^j e^j) + p^T (y + \sum_{j=1}^L k^j e^j) / k^j \in \{0,1\} \text{ for } j = 1, \dots, L \}$. Thus, $g^f(x') \leq$

Max $\{ f(y') + p^T y' / y' \in N^L \text{ and } y' \leq x \}$. Q.E.D.

We now state and prove the main theorem of this paper.

Theorem 1: Let X^* be a feasible allocation and p^* a price vector. $\langle p^*, X^* \rangle$ is a market equilibrium if and only if the pair $\langle p^*, m^* \rangle$ solves:

Minimize $\sum_{i=1}^H m(i) + p^T w$

Subject to $\sum_{i=1}^H m(i) + p^T x \geq V(x)$ for all $x \in N^L$ with $x \leq w + e$, $p \in \mathcal{R}_+^L$

where $m^* \in \mathcal{R}^H$ with $m^* = \langle m^*(i) / i = 1, \dots, H \rangle$ satisfies $m^*(i) = f^i(X^{*i}) - p^{*T} X^{*i}$ for $i = 1, \dots, H$.

Proof: Suppose $\langle p^*, X^* \rangle$ is a market equilibrium. Let $x \in N^L$ and $V(x) = \sum_{i=1}^H f^i(X^i)$ where

$X = \langle X^i / i = 1, \dots, H \rangle \in F(x)$.

Thus, for all $i = 1, \dots, H$: $f^i(X^{*i}) - p^{*T} X^{*i} \geq f^i(X^i) - p^{*T} X^i$.

Summing over i we get: $\sum_{i=1}^H m^*(i) + p^{*T} x \geq V(x)$ for all $x \in N^L$.

Thus, $\langle p^*, m^* \rangle$ satisfies the constraints.

Now, let $\langle p, m \rangle$ satisfy the constraints. Thus, $\sum_{i=1}^H m(i) + p^T w \geq V(w)$.

However, by Proposition 1, X^* is efficient and hence $\sum_{i=1}^H m^*(i) + p^{*T} w = V(w)$.

Thus, $\sum_{i=1}^H m(i) + p^T w \geq \sum_{i=1}^H m^*(i) + p^{*T} w$.

Hence, $\langle p^*, m^* \rangle$ solves the minimization problem.

Now, suppose $\langle p^*, m^* \rangle$ solves the given minimization problem. Towards a contradiction suppose $\langle p^*, X^* \rangle$ is not a market equilibrium. Thus, there exists $i \in H$ and $x \in N^L$, such that $f^i(x) - p^{*T} x > f^i(X^{*i}) - p^{*T} X^{*i}$.

Suppose $x \leq X^{*i} + e$.

Since $X^* \in F(w)$, $w^* = x + \sum_{k \neq i} X^{*k} \leq X^{*i} + e + \sum_{k \neq i} X^{*k} = w + e$.

Thus, $V(w^*) \geq f^i(x) + \sum_{k \neq i} f^k(X^{*k})$

$$\begin{aligned} &> f^i(X^{*i}) - p^{*T} X^{*i} + p^{*T} x + \sum_{k \neq i} f^k(X^{*k}) - p^{*T} \sum_{k \neq i} X^{*k} + p^{*T} \sum_{k \neq i} X^{*k} \\ &= f^i(X^{*i}) - p^{*T} X^{*i} + \sum_{k \neq i} f^k(X^{*k}) - p^{*T} \sum_{k \neq i} X^{*k} + p^{*T} w^* \end{aligned}$$

$$= \sum_{i=1}^H m^*(i) + p^{*\top} w^*,$$

leading to a contradiction.

Thus it is not the case that $x \leq X^{*i} + e$.

In fact we have shown that for $i \in H$ and $x \in N^L$, with $x \leq X^{*i} + e$, it is the case that $f^i(x) - p^{*\top} x \leq f^i(X^{*i}) - p^{*\top} X^{*i}$.

Since $X^{*i} \ll X^{*i} + e$, for $t \in (0,1)$, t sufficiently small, the real vector $X^{*i} + t(x - X^{*i}) \ll X^{*i} + e$.

Let $g^i = g^{f^i}$ denote the canonical extension of f^i .

Thus, $g^i(x) - p^{*\top} x > g^i(X^{*i}) - p^{*\top} X^{*i}$.

Since g^i is concave $g^i(X^{*i} + t(x - X^{*i})) \geq g^i(X^{*i}) + t(g^i(x) - g^i(X^{*i}))$.

Hence, $g^i(X^{*i} + t(x - X^{*i})) - p^{*\top}(X^{*i} + t(x - X^{*i})) \geq g^i(X^{*i}) + t(g^i(x) - g^i(X^{*i})) - p^{*\top}(X^{*i} + t(x - X^{*i}))$.

Now, $g^i(X^{*i} + t(x - X^{*i})) - p^{*\top}(X^{*i} + t(x - X^{*i})) \geq g^i(X^{*i}) - p^{*\top} X^{*i} + t([g^i(x) - p^{*\top} x] - [g^i(X^{*i}) - p^{*\top} X^{*i}]) > g^i(X^{*i}) - p^{*\top} X^{*i}$.

However, $X^{*i} + t(x - X^{*i}) \ll X^{*i} + e$.

Thus by Lemma 1, $g^i(X^{*i} + t(x - X^{*i})) - p^{*\top}(X^{*i} + t(x - X^{*i})) \leq \text{Max} \{f^i(y) - p^{*\top} y / y \in N^L \text{ such that } y \leq X^{*i} + e\} = f^i(X^{*i}) - p^{*\top} X^{*i}$, leading to a contradiction.

This establishes the theorem. Q.E.D.

Note: In the statement of Theorem 1 and in its proof, the constraint $p \in \mathfrak{R}_+^L$, which appears for the linear, programming (minimization) problem, could be easily dispensed with without diluting the result in any way. The fact that the theorem concerns a price vector would then imply our version of Theorem 1.

The main result (Theorem 4.1) in Yang (2001) can be strengthened without requiring the production functions to be discrete concave, as follows:

Theorem 2: Let X^* be a feasible allocation and p^* a price vector. $\langle p^*, X^* \rangle$ is a constrained market equilibrium if and only if the pair $\langle p^*, m^* \rangle$ solves:

$$\text{Minimize } \sum_{i=1}^H m(i) + p^{\top} w$$

$$\text{Subject to } \sum_{i=1}^H m(i) + p^{\top} x \geq V(x) \text{ for all } x \in N^L \text{ with } x \leq w, p \in \mathfrak{R}_+^L,$$

$$m(i) + p^{\top} x \geq f^i(x), \text{ for all } x \in N^L \text{ with } x \leq w \text{ and } i = 1, \dots, H.$$

where $m^* \in \mathfrak{R}^H$ with $m^* = \langle m^*(i) / i = 1, \dots, H \rangle$ satisfies $m^*(i) = f^i(X^{*i}) - p^{*\top} X^{*i}$ for $i = 1, \dots, H$.

Proof: The proof is similar to the proof of Theorem 1, but is being provided here for completeness.

Suppose $\langle p^*, X^* \rangle$ is a constrained market equilibrium. Let $x \in N^L$ with $x \leq w$ and $V(x) = \sum_{i=1}^H f^i(X^i)$ where $X = \langle X^i / i = 1, \dots, H \rangle \in F(x)$.

Clearly, $X^i \leq x \leq w$ for all $i = 1, \dots, H$.

Thus, for all $i = 1, \dots, H$: $f^i(X^{*i}) - p^{*T}X^{*i} \geq f^i(X^i) - p^{*T}X^i$.

Summing over i we get: $\sum_{i=1}^H m^*(i) + p^{*T}x \geq V(x)$ for all $x \in N^L$.

Thus, $\langle p^*, m^* \rangle$ satisfies the constraints.

Now, let $\langle p, m \rangle$ satisfy the constraints. Thus, $\sum_{i=1}^H m(i) + p^T w \geq V(w)$.

However, by Proposition 1, X^* is efficient and hence $\sum_{i=1}^H m^*(i) + p^{*T}w = V(w)$.

Thus, $\sum_{i=1}^H m(i) + p^T w \geq \sum_{i=1}^H m^*(i) + p^{*T}w$.

Thus, $\langle p^*, m^* \rangle$ solves the minimization problem.

Now, suppose $\langle p^*, m^* \rangle$ solves the given minimization problem. Towards a contradiction suppose $\langle p^*, X^* \rangle$ is not a constrained market equilibrium. Thus, there exists $i \in H$ and $x \in N^L$ with $x \leq w$, such that $f^i(x) - p^{*T}x > f^i(X^{*i}) - p^{*T}X^{*i}$.

Thus, $f^i(x) > m^*(i) + p^{*T}x$, which leads to a violation of a constraint of the minimization problem and consequently a contradiction.

Thus, $\langle p^*, X^* \rangle$ is a constrained market equilibrium. Q.E.D.

Note: Sun and Yang (2004) establish the existence of market equilibrium allocations without assuming that the production functions are discrete concave. They show that a market equilibrium exists if and only if **there exists a price vector p^* such that $V(w) - p^{*T}w \geq V(x) - p^{*T}x$ for all $x \in N^L$.**

However, we are able to show that under the assumption of concave production functions the following is true:

Theorem 3: There exists a market equilibrium if and only if there exists a price vector p^* such that $V(w) - p^{*T}w \geq V(x) - p^{*T}x$ for all $x \in N^L$ with $x \leq w + e$.

Proof: Let $\langle p^*, X^* \rangle$ be a market equilibrium. By Proposition 1, $\sum_{i=1}^H f^i(X^{*i}) = V(w)$.

Let $x \in N^L$ with $x \leq w + e$. By Theorem 1, $V(w) - p^{*T}w \geq V(x) - p^{*T}x$, since $\sum_{i=1}^H X^{*i} = w$.

Now suppose there exists a price vector p^* such that $V(w) - p^{*T}w \geq V(x) - p^{*T}x$ for all $x \in N^L$ with $x \leq w + e$. Let X^* be an efficient allocation. Thus, $\sum_{i=1}^H f^i(X^{*i}) = V(w)$. Let,

$m^* \in \mathcal{R}^H$ with $m^* = \langle m^*(i) / i = 1, \dots, H \rangle$ satisfying $m^*(i) = f^i(X^{*i}) - p^{*T}X^{*i}$ for $i = 1, \dots, H$.

Since, $\sum_{i=1}^H m^*(i) + p^{*T}x = V(w) - p^{*T}w + p^{*T}x \geq V(x)$ for all $x \in N^L$ with $x \leq w + e$,

$\langle p^*, m^* \rangle$ satisfies the constraints of the linear programming problem in Theorem 1.

Let $m \in \mathfrak{R}^H$ with $m = \langle m(i) / i = 1, \dots, H \rangle$ satisfy $\sum_{i=1}^H m(i) + p^T x \geq V(x)$ for all $x \in N^L$ with $x \leq w + e$.

Thus, $\sum_{i=1}^H m(i) + p^T w \geq V(w) = \sum_{i=1}^H m^*(i) + p^{*T} w$.

Thus, $\langle p^*, m^* \rangle$ solves the linear programming problem in Theorem 1. By Theorem 1, $\langle p^*, X^* \rangle$ is a market equilibrium. Q.E.D.

4. Properties of the maximum value function for existence of equilibrium: We now investigate properties, which when satisfied by the maximum value function, guarantees the existence of a market equilibrium. In this section we assume that $w \in N^L \cap \mathfrak{R}_{++}^L$.

Let $C(w) = \{x \in N^L / x \leq w + e\}$ and let $C^*(w)$ denote the convex-hull of $C(w)$.

The cardinality of $C(w) = \prod_{j=1}^L (w^j + 2)$. Let M denote the integer $\prod_{j=1}^L (w^j + 2) - 1$.

A function $f: N^L \rightarrow \mathfrak{R}$ is said to be partially concave at w if for any positive integer K and arrays $\langle x^k / k = 1, \dots, K \rangle$, $\langle t^k / k = 1, \dots, K \rangle$ with $x^k \in C(w)$ and $t^k \geq 0$ for $k = 1, \dots, K$: $[w = \sum_{k=1}^K t^k x^k, \sum_{k=1}^K t^k = 1]$ implies $[f(w) \geq \sum_{k=1}^K t^k f(x^k)]$.

Lemma 2: A function $f: N^L \rightarrow \mathfrak{R}$ is partially concave at w if and only if there exists $p \in \mathfrak{R}^L$ such that $f(w) - p^T w \geq f(x) - p^T x$ for all $x \in C(w)$.

Proof: Let $C(w) \setminus \{w\}$ be equal to the set $\{x^k / k = 1, \dots, M\}$.

Suppose f is partially concave at w . Towards a contradiction suppose there does not exist $p \in \mathfrak{R}^L$ such that $f(w) - p^T w \geq f(x) - p^T x$ for all $x \in C(w)$.

Hence, there does not exist $\alpha, \beta, \gamma \in \mathfrak{R}_+^L$: $\alpha^T (x^k - w) - \beta^T (x^k - w) + \gamma^k = f(x^k) - f(w)$ for all $k = 1, \dots, M$.

By Farkas' Theorem there exists $t \in \mathfrak{R}_+^M$ such that $\sum_{k=1}^M t^k (x^k - w) \leq 0$, $-\sum_{k=1}^M t^k (x^k - w) \leq 0$

and $\sum_{k=1}^M t^k [f(x^k) - f(w)] > 0$.

Thus, $\sum_{k=1}^M t^k > 0$.

Dividing the three inequalities above by $\sum_{k=1}^M t^k$, we get there exists $s \in \mathfrak{R}_+^M$ such that

$\sum_{k=1}^M s^k x^k = w$, $\sum_{k=1}^M s^k = 1$ and $\sum_{k=1}^M s^k f(x^k) > f(w)$, contradicting that f is partially concave at w .

Hence, there exists $p \in \mathfrak{R}^L$ such that $f(w) - p^T w \geq f(x) - p^T x$ for all $x \in C(w)$.

Now suppose that there exists $p \in \mathfrak{R}^L$ such that $f(w) - p^T w \geq f(x) - p^T x$ for all $x \in C(w)$.

Hence there exists $\alpha, \beta, \gamma \in \mathfrak{R}_+^L$: $\alpha^T (x^k - w) - \beta^T (x^k - w) + \gamma^k = f(x^k) - f(w)$ for all $k = 1, \dots, M$.

By Farkas' Theorem there does not exist $t \in \mathfrak{R}_+^M$ such that $\sum_{k=1}^M t^k (x^k - w) = 0$ and

$$\sum_{k=1}^M t^k [f(x^k) - f(w)] > 0.$$

Thus, $[t \in \mathfrak{R}_+^M, t^k \geq 0 \text{ for } k = 1, \dots, M, w = \sum_{k=1}^M t^k x^k, \sum_{k=1}^M t^k = 1]$ implies $[f(w) \geq$

$$\sum_{k=1}^M t^k f(x^k)].$$

Thus, f is partially concave at w . Q.E.D.

A function $f: N^L \rightarrow \mathfrak{R}$ is said to be Weakly Monotonic at w if:

- (1) For all $j = 1, \dots, L$: $f(w + e^j) \geq f(w)$;
- (2) $f(w + e) > f(w)$.

It is easy to see that if for some i , f^i is non-decreasing (i.e. for all $x, y \in N^L$: $[x \geq y]$ implies $[f^i(x) \geq f^i(y)]$) and weakly increasing (i.e. for all $x, y \in N^L$: $[x \gg y]$ implies $[f^i(x) > f^i(y)]$), then V is Weakly Monotonic at w .

Lemma 3: Suppose $f: N^L \rightarrow \mathfrak{R}$ is Weakly Monotonic at w . Then, f is partially concave at w if and only if there exists $p \in \mathfrak{R}_+^L \setminus \{0\}$ such that $f(w) - p^T w \geq f(x) - p^T x$ for all $x \in C(w)$.

Proof: By Lemma 2, f is partially concave at w if and only if there exists $p \in \mathfrak{R}^L$ such that $f(w) - p^T w \geq f(x) - p^T x$ for all $x \in C(w)$.

Suppose towards a contradiction f is partially concave but $p_j < 0$, for some j . Then, $0 \leq f(w + e^j) - f(w)$ by Weak Monotonicity of f at w and $f(w + e^j) - f(w) \leq p^T (w + e^j - w) = p_j < 0$, leads to a contradiction.

Thus $p \in \mathfrak{R}_+^L$.

If $p = 0$, then $0 < f(w + e) - f(w)$ by Weak Monotonicity and $f(w + e) - f(w) \leq p^T e = 0$, again leads to a contradiction.

Thus, $p \in \mathfrak{R}_+^L \setminus \{0\}$. Q.E.D.

In view of Lemmas 2 and 3 and Theorem 3, we can state the following result:

Theorem 4: Suppose V is Weakly Monotonic at w . A market equilibrium exists if and only if V is partially concave at w .

5. An Illustrative Example: Consider the following two agent ($H = 2$), three input ($L = 3$) integer allocation problem with $w = e$. Let $f: \mathfrak{R}_+^L \rightarrow \mathfrak{R}_+$ be defined as follows: $f(0) = 0 = f(e^j)$ for $j = 1, 2, 3$; $f(e^j + e^k) = 3$ for $j, k \in \{1, 2, 3\}$ with $j \neq k$; $f(e) = 4$; for all $x \in N^3 \setminus \{y \in N^3 / y \leq e\}$, let $f(x) = f(\sum_{\{j / x_j > 0\}} e^j)$. f is discrete concave. Let $f^i = f$ for $i = 1, 2$. The integer

allocation problem $\langle \{f^1, f^2\}, e \rangle$ is an example of a bundle auction.

For this problem, if X is an efficient allocation then either $X^1 = e$ and $X^2 = 0$ or $X^1 = 0$ and $X^2 = e$.

Suppose $\langle p, X \rangle$ is a market equilibrium. Then by Proposition 1, X is efficient. Without loss of generality, suppose $X^1 = e$ and $X^2 = 0$. In order that X^2 maximize profits for agent 2 at price vector p , it must be that $p_j + p_k \geq 3$ for all $j, k \in \{1, 2, 3\}$ with $j \neq k$. Thus, $2(p_1 + p_2 + p_3) \geq 9$ or $p_1 + p_2 + p_3 \geq 4.5 > 4 = f(e) = f^1(e)$. Thus, X^1 does not maximize profits for agent 1 at price vector p .

Thus, this integer allocation problem does not have a market equilibrium.

It is easy to verify that the maximum value function V is not partially concave at $w = e$.

Note that $V(2e) = 8$, $V(e) = 4$, $V(e^j + e^k) = 3$ for $j, k \in \{1, 2, 3\}$ with $j \neq k$. Now $e = \frac{1}{4}(2e) +$

$$\frac{1}{4}(e^1 + e^2) + \frac{1}{4}(e^1 + e^3) + \frac{1}{4}(e^2 + e^3).$$

$$\frac{1}{4}V(2e) + \frac{1}{4}V(e^1 + e^2) + \frac{1}{4}V(e^1 + e^3) + \frac{1}{4}V(e^2 + e^3) = 2 + \frac{9}{4} = 4.25 > 4 = V(e).$$

Thus, V is not partially concave at w .

References

1. Lahiri, S. (2005): "Manipulation via Endowments in a market with profit maximizing agents", (mimeo) IFMR.
2. Shapley, L. and M. Shubik (1969): "On Market Games", *Journal of Economic Theory*, 1: 9-25.
3. Shapley, L. and M. Shubik (1976): "Competitive Outcomes in the Cores of Market Games", *International Journal of Game Theory*, 4: 229-237.
4. Sun, N. and Z. Yang (2004): "The Max-Convolution Approach to Equilibrium Models with Indivisibilities", (mimeo).
5. Yang, Z. (2001): "A Practical Competitive Market Model for Indivisible Commodities", Cowles Foundation Discussion Paper No. 1317, Cowles Foundation For Research in Economics, Yale University, New Haven, Connecticut.