



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in:
Current Biology

Cronfa URL for this paper:
<http://cronfa.swan.ac.uk/Record/cronfa34936>

Paper:

Unsworth, R. & Cullen-Unsworth, L. (2017). Seagrass meadows. *Current Biology*, 27(11), R443-R445.
<http://dx.doi.org/10.1016/j.cub.2017.01.021>

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.

Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.

<http://www.swansea.ac.uk/library/researchsupport/ris-support/>

Quick guide

Seagrass meadows

Richard K.F. Unsworth^{1,2}
and Leanne C. Cullen-Unsworth^{1,2}

What are seagrasses? Seagrasses are flowering plants adapted to life in the oceans. They are the only angiosperms able to withstand a saline existence, with all but one genus capable of completing their lifecycle in a fully submerged marine environment. Seagrasses comprise a relatively small group of plants, with 72 recognised species from four families of the plant order Alismatales. The comparatively low diversity of seagrass plants is counterbalanced by the high abundance and diversity of other organisms occupying the highly productive meadow ecosystems that these plants create. Seagrass meadows range in size and density from small 1m² patches to extensive continuous meadows covering tens of thousands of hectares. Meadows can be monospecific or comprise multispecies communities with up to 12 seagrass species present.

Just like terrestrial angiosperms, seagrasses flower, fruit and seed, with sexual reproduction facilitated (enabling pollination and movement of seeds and propagules) by animals as well as water movement (currents). Although sexual reproduction is key to maintaining diversity and adaptation, seagrasses also utilise asexual reproduction, which is important for meadow growth.

Where do seagrasses live?

Seagrasses are present in shallow seas on the continental shelf of all continents except Antarctica (Figure 1). Documented distributions include 125,000 km² globally, but estimates suggest seagrass meadows cover more than 600,000km² of the coastal ocean.

Seagrass meadows are largely soft sediment intertidal to subtidal benthic habitats that thrive in sheltered environments — such as shallow embayments, lagoons and estuaries — where wave attenuation is limited and light and nutrient levels are sufficient. Depth distribution is largely controlled by the availability of light with tidal



Figure 1. Seagrass meadows

Left, seagrasses such as this germinating *Zostera marina* seed are monocotyledons. Right, seagrass meadows are one of the worlds most productive marine habitats such as this *Zostera marina* meadow in North Wales (UK).

exposure, wave action, associated turbidity, and low salinity — all of which determine seagrass survival at their shallow edge. Intertidal seagrass is prevalent where there is sufficient shelter from wave action or where there is entrapment of water at low tide, providing protection from exposure.

Why are seagrasses defined as ecological engineers? Seagrass plants are termed ‘ecological engineers’, as they significantly modify their environment — by attenuating currents and waves, they reduce nutrient levels, decrease turbidity and thereby enhance conditions for optimal growth. The more extensive the area and denser the seagrass meadow, the greater the capacity to facilitate this positive feedback ensuring affable conditions.

Seagrasses enhance their own survival in sulphide-rich and otherwise toxic sediments by exuding oxygen from their roots into what would be anoxic mud. This action is supported by a three-way symbiosis between the seagrass, lucinid bivalves, and their sulphide-oxidizing gill bacteria. The symbiosis results in reduced sulphide stress for seagrasses, with provision of organic matter and radial oxygen release from the seagrass roots for the bivalves and their endosymbionts.

Why are seagrass meadows important?

The benefits that humans derive from seagrass meadows are termed ‘ecosystem services’. Some of these services span the global seagrass range — such as water filtration, habitat provision, fish nursery grounds and biodiversity support. However, local variation occurs in the services exploited and goods extracted.

Their provision of fish feeding and nursery habitat is often cited as rationale for seagrass protection. Indeed, seagrasses provide shelter from predators, and provide an abundant food supply, such as small crustaceans, supporting fish stocks [6]. There is growing recognition for the value of seagrass in supporting diverse and productive fauna that creates rich fishing grounds, providing a level of food security. For example, at low tide, exposed seagrass meadows in the tropics provide an accessible human hunting ground for small fish and edible invertebrates. At other times, traps and nets are used to catch predatory fish that migrate in and out of seagrass (Figure 2). The Atlantic cod (*Gadus morhua*) is the world’s third most landed fish species. This historically important fishery species favours seagrass as a nursery ground, with evidence suggesting that juvenile



Figure 2. Seagrass meadows support global food security.

Fishing activity in seagrass meadows is globally extensive, such as these women collecting intertidal invertebrates in Mozambique (left; source: Benjamin Jones) and the widespread use of nets to catch abundant fish in Indonesia (right).

cod grow faster and have improved chances of reaching maturity when living in seagrass meadows

Seagrass meadows also act as an important filter of coastal water pollution, maintaining the health of fisheries and associated coral reef habitats. Recent evidence has discovered that in the absence of coastal seagrass meadows, coral reef disease and bacterial contamination of fish species is increased due to reduced water filtration of wastewater.

In addition, seagrass meadows play a significant role in climate change mitigation through carbon storage within their sediments. Estimates suggest that seagrass meadows bury carbon 40 times faster than tropical forests, and provide one of the greatest contributions to the total carbon buried in ocean sediments. Because seagrasses are net photosynthesisers and are highly productive, they also rapidly utilise carbon from the water column. This has resulted in considerable attention being given to the role seagrass plays in altering seawater chemistry and potentially buffering small-scale impacts of ocean acidification on calcifying fauna, such as corals and molluscs.

What is the future outlook for seagrass meadows? The ecological value of seagrass meadows is irrefutable, and yet they remain in decline and largely marginalised on conservation agendas. Action is required to stem the loss of the world's seagrass meadows and to prioritise protection based on their array of ecosystem services.

As the human population grows, our impacts on the environment increase. Poor water quality (particularly elevated nutrients in waste water) as a result of poor catchment management together, with a diverse range of pollution sources is the biggest threat to seagrasses globally. Water quality problems are particularly acute in developing countries with rapidly growing economies, where municipal infrastructure and environmental legislation are lacking. Coastal development is a threat with boating, tourism, aquaculture, ports, energy projects and housing developments all placing pressure on seagrass systems. Local and regional threats to seagrass exacerbate the impacts of climate change, ocean acidification and sea level rise. Minimising local stress on seagrass will support their capacity to remain resilient to the impacts of larger scale, longer term stressors. This

means improving local water quality, preventing damage from destructive fishing practices, including seagrasses in Marine Protected Areas, preventing overexploitation of seagrass fisheries and mitigating stress from coastal development. Bold steps are required to ensure the recovery and restoration of these habitats, but the cumulative effect of multiple small-scale actions can work towards a brighter future for these underappreciated undersea gardens.

Where can I find out more?

- Cullen-Unsworth, L.C., and Unsworth, R.K.F. (2016). Strategies to enhance the resilience of the world's seagrass meadows. *J. Appl. Ecol.* 53, 967–972.
- Fourqurean, J.W., Duarte, C.M., Kennedy, H., Marba, N., Holmer, M., Mateo, M.A., Apostolaki, E.T., Kendrick, G.A., Krause-Jensen, D., McGlathery, K.J., *et al.* (2012). Seagrass ecosystems as a globally significant carbon stock. *Nat. Geosci.* 5, 505–509.
- Hemminga, M.A., and Duarte, C.M. (2000). *Seagrass Ecology*, First Edition, (Cambridge University Press).
- Lamb, J. B., J. A. J. M. van de Water, D. G. Bourne, C. Altier, M. Y. Hein, E. A. Fiorenza, N. Abu, J. Jompa, and C. D. Harvell. (2017) Seagrass ecosystems reduce exposure to bacterial pathogens of humans, fishes, and invertebrates. *Science* 355:731–733.
- Maxwell, P.S., Eklöf, J.S., van Katwijk, M.M., O'Brien, K.R., de la Torre-Castro, M., Boström, C., Bouma, T.J., Krause-Jensen, D., Unsworth, R.K.F., van Tussenbroek, B.I., *et al.* (2016). The fundamental role of ecological feedback mechanisms for the adaptive management of seagrass ecosystems – a review. *Biologi. Rev.*, DOI: 10.1111/brv.12294.
- Mtwana Nordlund, L., Koch, E.W., Barbier, E.B., and Creed, J.C. (2016). Seagrass ecosystem services and their variability across genera and geographical regions. *PLoS ONE* 11, e0163091.
- Unsworth, R.K.F., Hinder, S.L., Bodger, O.G., and Cullen-Unsworth, L.C. (2014). Food supply depends on seagrass meadows in the coral triangle. *Environ. Res. Lett.* 9, 094005.
- van der Heide, T., Govers, L.L., de Fouw, J., Olf, H., van der Geest, M., van Katwijk, M.M., Piersma, T., van de Koppel, J., Silliman, B.R., Smolders, A.J.P., *et al.* (2012). A three-stage symbiosis forms the foundation of seagrass ecosystems. *Science* 336, 1432–1434.
- van Tussenbroek, B.I., Villamil, N., Márquez-Guzmán, J., Wong, R., Monroy-Velázquez, L.V., and Solís-Weiss, V. (2016). Experimental evidence of pollination in marine flowers by invertebrate fauna. *Nat. Commun.* 7, 12980.
- Waycott, M., Duarte, C.M., Carruthers, T.J.B., Orth, R.J., Dennison, W.C., Olyarnik, S., Calladine, A., Fourqurean, J.W., Heck, K.L., Hughes, A.R., *et al.* (2009). Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proceed. Nat. Acad. Sci. USA* 106, 12377–12381.

¹Department of Biosciences, Wallace Building, Swansea University, Swansea, SA2 8PP, UK;
²Sustainable Places Research Institute, Cardiff University, UK.

E-mail: r.k.f.unsworth@swansea.ac.uk

