

Research

Open Access

## Patients' satisfaction and quality of life in coronary artery disease

Mohsen Asadi-Lari<sup>1</sup>, Chris Packham<sup>2</sup> and David Gray\*<sup>1</sup>

Address: <sup>1</sup>Division of Cardiovascular Medicine, University Hospital, Nottingham, NG7 2UH, UK and <sup>2</sup>Division of Epidemiology & Public Health, University of Nottingham, UK

Email: Mohsen Asadi-Lari - msxma@nottingham.ac.uk; Chris Packham - Chris.Packham@nottingham.ac.uk; David Gray\* - d.gray@nottingham.ac.uk

\* Corresponding author

Published: 22 October 2003

Received: 09 July 2003

*Health and Quality of Life Outcomes* 2003, **1**:57

Accepted: 22 October 2003

This article is available from: <http://www.hqlo.com/content/1/1/57>

© 2003 Asadi-Lari et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

### Abstract

**Objectives:** To assess satisfaction of survivors of coronary artery diseases (CAD) with healthcare services and to determine whether specific components of standard health-related quality of life (HRQL) assessment tools might identify areas of satisfaction and dissatisfaction.

**Method:** A specific tool developed to provide a comprehensive assessment of healthcare needs was administered concomitantly with generic and specific HRQL instruments, on 242 patients with CAD, admitted to an acute coronary unit during a single year.

**Results:** 92.5% of patients confirmed their trust in and satisfaction with the care given by their General Practitioner; even so, one third experienced difficulty getting an appointment and a quarter wanted more time for each consultation or prompt referral to a specialist when needed. Around a third expressed dissatisfaction with advice from the practice nurse or hospital consultant. Overall 54% were highly satisfied with services, 33% moderately satisfied and 13% dissatisfied.

Cronbach's alpha was 0.87; the corrected total-item correlation ranged between 0.55–0.75, with trivial 'floor' score and low 'ceiling' effect. Several domains in all three HRQL tools correlated with items relating to satisfaction. The Seattle Angina Questionnaire Treatment Score correlated significantly with all satisfaction items and with the global satisfaction score.

**Conclusion:** Cardiac patients' demanded better services and advice from, and more time with, health professionals and easier surgery access. The satisfaction tool showed acceptable psychometric properties. In this patient group, disease-specific HRQL tools seem more appropriate than generic tools for surveys of patient satisfaction.

### Introduction

The modern approach to healthcare seeks to engage the attention of both patients and public in developing healthcare services and equity of access, but this is not easy to achieve, requiring time, commitment, political support and cultural change to overcome barriers to

change. [1] Improvement in selected aspects of health care delivery through quality assurance and outcome assessment has been driven by political expediency. While this is important, a 'bottom up' assessment of patient satisfaction seems preferable if service improvement is to be translated into improved quality of life. [2,3]

Patients' satisfaction is related to the extent to which general health care needs and condition-specific needs are met. Evaluating to what extent patients are satisfied with health services is clinically relevant, as satisfied patients are more likely to comply with treatment [4,5] and take an active role in their own care. [6] In addition, health professionals may benefit from satisfaction surveys that identify potential areas for service improvement and health expenditure may be optimised through patient-guided planning and evaluation. [7]

Critics draw attention to the lack of a standard approach to measuring satisfaction and of comparative studies [3] and so the significance of the results of surveys is often ignored. There is less controversy with respect to clinical outcome measures, as health-related quality of life (HRQL) is not only widely regarded as a robust measure of outcome but also is extensively used in several clinical areas. [8,9]

We conducted a cross-sectional survey to establish, from a patient perspective, how satisfied survivors of coronary artery diseases (CAD) were with healthcare services. This might indicate areas that warranted review, evaluation and change where appropriate.

We also wanted to determine whether specific components of standard health-related quality of life (HRQL) assessment tools might identify areas of satisfaction and dissatisfaction. Identifying variables that link patient satisfaction and accepted measures of outcome might result in identifying areas for quality improvement.

## Method

The health care needs of all patients with coronary artery disease (CAD) admitted to an coronary unit during a single year were established. Their consent to participate in the study was obtained. Diagnosis was based on clinical and angiographic features according to the Nottingham Heart Attack Register (NHAR), which categorised patients in five diagnostic groups: definite myocardial infarction, suspected MI, ischaemic heart disease, chest pain and symptoms not related to heart disease. [10,11] Patients were sent a detailed questionnaire three to six months after discharge from hospital.

The methodology has been described in detail elsewhere, [12] but briefly a needs assessment questionnaire was developed following a rigorous review of expert opinion and the literature, discussions with medical staff and information compiled during face-to-face interviews with patients with CAD. After a pilot study, [13] a healthcare needs assessment (HNA) tool emerged, consisting of 48 questions in 5-score Likert scale (1 indicates *more needs* versus 5 with *no needs*) in five domains of 'physical needs,

'satisfaction', 'informational needs', 'social needs, and 'concerns'. Two generic (Short Form-12 and EuroQoL-5D) quality of life questionnaires were chosen to cover the limitations of each tools.; in addition one specific HRQL instrument (Seattle Angina Questionnaire) was administered concomitantly. The satisfaction component, which was included in the main health needs questionnaire, consisted of 11 questions mainly related to hospital-based services and health centres or family doctor surgeries (table 2). A global score in each domain was calculated to permit comparison with other variables. We compared satisfaction data with health-related quality of life information obtained from the same patients, which could be assumed as a 'gold standard'.

### Seattle Angina Questionnaire (SAQ)

This has well-established psychometric properties, measures broader aspects of the effects of coronary disease than other disease-specific tools and can detect physical limitations due to coronary disease. It is particularly useful in the presence of co-morbidity, [14] corresponding well with the Canadian Cardiovascular Society Classification. [15] It consists of 19 items grouped in five components: *physical functioning* (SAQ Phys), *angina stability* (AS), *angina frequency* (AF), *treatment satisfaction* (TS), and *QOL perception* (SAQ QOL).

### Short Form 12 (SF-12)

This is an abridged form of the better-known Short Form 36 (SF-36) [16] which has produced consistent results in several European countries and in a diverse range of conditions. It contains 12 questions from which are derived physical and mental component scores (PCS & MCS); these are as precise as the SF-36 [17,18].

### EuroQoL (EQ-5D)

The EuroQoL questionnaire defines health in terms of five dimensions: *mobility, self-care, usual activities, pain or discomfort, and anxiety or depression* in a three degree format that is *no problem, moderate* and *severe*. The greater the score, the worse the quality of life. Another question deals with overall health in a 0–100 scaling format. The validity and reliability of the EQ-5D questionnaire have been tested in a range of patient groups. [19,20] There is a strong correlation between the EQ-5D and the SF-12 in adults, [21] but to our knowledge, there has been no published study of the EQ-5D in myocardial infarction.

### Statistical analysis

SPSS version 11.0 was used for statistical purposes, using descriptive and correlation analysis, comparison of means, reliability and non-parametric (Mann Whitney-U) tests where indicated. Contingency tables were formed to look for any correlation and chi-square test was used to measure the association between variables. Psychometric

**Table 1: Demographic data**

Variable		N	Percent
Age group	= <54	29	12
	55-64	47	20
	65-74	73	30
	75-84	67	28
	> = 85	23	10
Gender	Female	99	41
	Male	143	59
Education level	School certificate	23	10
	Left at age <16	146	66
	"O" or "A" level	14	6
	College or higher	39	18
Occupation	Retired	194	80.5
	Unemployed	15	6
	Full-time	24	10
	Part-time	8	3
Diagnosis	Definite MI	27	11
	Possible MI	34	14
	IHD	152	64
	Chest pain	26	11
Social class	I	3	1
	II	35	16
	III non-manual	31	14
	III manual	62	28
	IV	71	32
First call in emergencies	V	21	9
	GP	134	56
	999	81	34
	Others	24	10

analysis demonstrated acceptable properties, described elsewhere [12]; internal consistency in the five domains was quite high, ranging from 0.83 to 0.89 and specifically 0.87 in satisfaction domain. The corrected total-item correlation, as an indicator of item specificity, ranged between 0.55-0.75, with a trivial floor score of 0.4% and a ceiling effect of 8.7%. To compare satisfaction variables with HRQL and other HNA scores, correlation coefficients were computed, which was considered significant at  $p < 0.05$ . The regional ethical committee approved this survey.

## Results

### Demographic data

242 patients (59% male) returned the completed questionnaires; response rate was 93%. Ages ranged between 31-93 years (mean = 69.7, 95% CI: 68.2,71.2). Seventy-one percent (=169) left school at age 16 or less and 21% (=53) completed higher education. Table 1 describes the major demographic characteristics of all but 3 patients. Social class, derived from the last occupation, was determined in 223 patients. [22] 69 (31%) had non-manual and 154 (69%) manual jobs (table 1). Health-related

quality of life were generally lower in women, especially in SAQ-AS and PCS ( $t = -2.04$  and  $t = -1.99$  respectively,  $P < 0.05$ ) and in several domains of the three QOL tools in those over 65 years of age. [12]

### Health needs

Elderly affected physical ( $p < 0.01$ ) and social needs ( $p < 0.05$ ) and women were more dissatisfied in general ( $p < 0.05$ ). Lower social class was concomitant with more needs in all health needs domains ( $p < 0.01$ ) although not statistically significant in satisfaction score. A unique score was calculated in each domain to facilitate comparisons among various variables. The mean score (mean 4, SD 0.76) was highly skewed towards higher level of satisfaction (measure of skewness: -4.53). [23]

### Patients' satisfaction

Assessment of satisfaction was obtained from 11 scores contained within several HNA domains (table 2) dealing mainly with patients' perceptions of members of the health professions and the information and treatment provided. Scores less than 3 were categorised as 'poor' and

**Table 2: Descriptive distribution of satisfaction variables**

Variable	Mean	SD	Percentage of dissatisfaction*
1. Do you feel satisfied with ability to get an appointment with your GP?	3.77	1.27	33.6
2. Do you feel satisfied with having enough time with your GP at each visit?	4.18	1.06	23.1
3. Do you feel satisfied with ability to be referred to a consultant?	4.13	1.06	27.1
4. Do you feel that your GP does his/her best for you?	4.56	0.68	7.5
5. Do you feel happy with your current treatment?	4.0	0.99	30.1
6. Was the advice from your GP helpful?	3.73	1.13	32.9
7. Were you satisfied with nurse services	3.56	1.25	37
8. Were you satisfied with rehabilitation services?	3.65	1.50	33.7
9. Were you satisfied with hospital consultant services?	3.78	1.27	31.3
10. Do you think assistance with easier access to GP might help you cope with your illness better?	4.02	1.49	30.6
11. Does your GP fully understand your health needs?	4.30	0.98	17.6
HNA global satisfaction score	4.00	0.76	

\* The first 3 scores (Likert scale) in each group assumed as dissatisfied and more than 4 as satisfied.

a score of 4 or higher as 'reasonable' satisfaction with health services.

One-third reported difficulty getting an appointment with their GPs. About one-fourth were dissatisfied with the amount of time devoted to each visit and with their ability to see a consultant when needed. One-third expressed dissatisfaction with advice received from their GP, although more than 92% believed that the GP provided a good quality service (table 2).

Women ( $P < 0.05$ ) and younger patients ( $< 75$ ) were more likely to be dissatisfied with their ability to get to see their GP ( $P = 0.01$ ) or happy with their relationship with their GP ( $p < 0.05$ ). 18 patients (8%) used the private sector as their main health provider; there were no significant differences in levels of satisfaction among those utilising public rather than private healthcare services. Patients who reported that they often had to ask for a home visit (12%;  $n = 28$ ) also reported lower levels of satisfaction with the appointment process compared with patients who were able enough to get to the Health Centre ( $p < 0.05$ ). 57% ( $n = 138$ ) were quite confident that their GPs fully appreciated their health needs, but the remainder doubted their needs were appropriately recognised. Women in particular expected their GP to fully understand their needs ( $p = 0.001$ ).

#### Mean satisfaction score

Overall women were less satisfied with healthcare services ( $P < 0.05$ ), as were the under 75 s (mean rank 114.25 *v* 124.52), but this was not statistically significant. Social class did not appear to influence satisfaction.

Patients with co-morbid illness were significantly less satisfied than those suffering from coronary disease alone ( $P$

$< 0.01$ ). Mean satisfaction score was highly correlated with other HNA components (correlation coefficient ranged 0.26–0.70,  $p < 0.001$ ). Comparison of score means indicated that patients with other physical, informational, social needs and medical concerns were more dissatisfied ( $p < 0.01$ ). Overall, components of the HNA tool were indicative of overall levels of satisfaction with healthcare.

#### HRQL scores

The SAQ-TS correlated significantly with all satisfaction items, while several domains in all three HRQL tools correlated with satisfaction scores (tables 3,4). Satisfied patients valued their health status higher in visual analogue score by 9 scores; that was 56 in dissatisfied versus 65 in satisfied patients ( $P < 0.001$ ). Mean satisfaction score correlated moderately with EQ anxiety/depression ( $r = 0.40$ – $0.59$ ) and SAQ treatment satisfaction domains and correlated weakly with SF-12 domains, EQ self care and visual analogue scale and SAQ-QOL ( $r = 0.20$ – $0.39$ ). The remaining HRQL domains did not correlate with the mean satisfaction score.

#### Discussion

Patient satisfaction is considered by some to be of dubious benefit in facilitating the process of clinical care, as patients have no specific clinical expertise and are readily influenced by non-medical factors; in addition, there are few reports on the reliability of satisfaction surveys [3,24,25]. Nevertheless, satisfied patients are more likely to comply with medical treatment and therefore ought to have a better outcome. [3]

In the absence of an available instrument for assessing our patients' healthcare needs, we developed a healthcare needs assessment questionnaire. This included a series of

**Table 3: Mean differences in HRQL scores within various items of satisfaction**

	EQ-Mob	EQ-SC	EQ-UA	EQ-PD	EQ-AD	EQ-VAS	SAQ phys	SAQ AS	SAQ AF	SAQ TS	SAQ QOL	PCS	MCS
Getting an appointment	-.28	-.67	-.51	-.88	<b>-4.53</b>	-2.23*	-.21	-1.39	-.92	<b>-3.85</b>	<b>-2.85</b>	-.21	<b>-2.77</b>
Enough time in GP visits	-.47	-.14	-.95	-.48	<b>-3.17</b>	-1.54	-.66	-.38	-.59	<b>-3.58</b>	-1.96*	-.13	<b>-2.56</b>
Satisfaction with referral	-.47	<b>-2.63</b>	-.17	-.27	<b>-3.11</b>	-2.13*	-1.09	-.53	-.96	<b>-4.70</b>	<b>-2.89</b>	-1.75	<b>-2.95</b>
GP service satisfaction	-.28	-1.27	-.92	-.53	-2.34*	-.47	-.84	-1.64	-1.08	<b>-4.86</b>	-1.02	-.02	-2.24*
Satisfied with current treatment	<b>-3.16</b>	<b>-2.70</b>	-1.71	<b>-3.54</b>	<b>-5.23</b>	<b>-4.2</b>	<b>-2.93</b>	<b>-2.88</b>	<b>-2.40*</b>	<b>-7.71</b>	<b>-5.36</b>	<b>-2.90</b>	<b>-3.85</b>
Satisfied with GP advice	-1.74	-1.57	-1.82	-.33	<b>-3.36</b>	-1.16	-.6	-.03	-1.16	<b>-5.38</b>	-2.43*	-1.14	-2.34*
Satisfied with Nurse services	-1.40	-.74	-1.59	-.54	<b>-2.61</b>	-2.24	-1.24	-.52	-.71	<b>-5.41</b>	<b>-3.59</b>	-1.13	<b>-2.77</b>
Satisfied with Rehabilitation services	<b>-2.85</b>	<b>-2.82</b>	-2.53*	-1.35	<b>-3.30</b>	-2.35*	<b>-3.21</b>	-.39	-.68	<b>-3.56</b>	<b>-3.50</b>	<b>-2.57</b>	<b>-4.06</b>
Satisfied with consultant services	-1.15	-.52	-1.15	-1.45	-1.91	-1.62	-1.71	-.90	-.17	<b>-5.42</b>	<b>-2.93</b>	-1.33	-2.34*
Need easier access to health services	-.38	-1.9	-1.62	-1.32	<b>-4.5</b>	<b>-3.0</b>	-2.2*	-.85	-1.11	<b>-3.89</b>	<b>-2.74</b>	-1.6	<b>-3.25</b>
GP understands needs?	-2.34*	-.56	-1.87	-.39	-2.21*	-.95	-.92	-.10	-.98	<b>-6.01</b>	<b>-3.26</b>	-1.2	<b>-2.62</b>

Figures show the z value of mean differences within QOL scores. \* indicates  $p < 0.05$  and bold if  $p < 0.01$ . Abbreviations.: EQ-Mob = EQ-5D mobility, EQ-SC = EQ-5D self care, EQ-UA = EQ-5D usual activities, EQ-PD = EQ-5D pain/discomfort, EQ-AD = EQ-5D anxiety/depression, EQ-VAS = EQ-5D visual analogue scale. Seattle Angina Questionnaire components: SAQ Phys = physical activities; AS = angina frequency; AF = angina stability; TS = treatment satisfaction; QOL = quality of life perception. Short Form-12: PCS = physical component score; MCS = mental component score.

questions relating to some aspect of satisfaction with clinical care. While its internal consistency (and therefore its reliability) was satisfactory, further studies are warranted to investigate as Guldvog suggests [6] whether it will retain its reliability over time and its validity when compared with other related tools. Health needs and satisfaction with clinical care are related to some extent-meeting other health needs increases the likelihood of overall satisfaction with healthcare services generally.

Patients did think highly of their General Practitioners (more so than other members of the medical and nursing staff) and were generally satisfied with the care they delivered. This confirms the General Practitioner's central role in delivering healthcare to cardiac patients and highlights the importance that patients place on the availability of first line health provision. Nevertheless, from a patient perspective, four main areas were identified where clinical care might be improved.

Patients wanted first an appointment system to improve the accessibility of the family doctor. Optimal medication is no guarantee that intermittent exacerbation of symptoms of coronary heart disease will be prevented and cardiac patients recognise that they require easy and prompt access to their principal primary carer, the GP. Second, patients have issues regarding immediate and future treatment they wish to discuss and expect more time for each consultation. Third, dissatisfied patients have greater need for information which is consonant with other results. [26] Fourth, according to our routine clinical experience, patients read in the media about many of the technical aspects of modern cardiological practice available in secondary care and expect referral to hospital for a specialist

opinion. These study findings provide a snapshot of areas where patients consider services might be improved quite easily. Qualitative and quantitative studies are planned to investigate these further and to determine how satisfaction surveys can be introduced into routine clinical practice.

Our results are consistent with previous reports in terms of the probability of dissatisfaction with healthcare services among younger patients [3] but contradict findings of even greater dissatisfaction among female patients [27]. Gender differences in HRQL perception have been reported before, [28,29] which implies that measuring satisfaction in our study involved more than simply questioning patients about healthcare facilities. [27]

We had considered that readily available HRQL tools might be of value as surrogates for more detailed surveys of patient satisfaction. In our patients with coronary disease, satisfaction with the care process was closely correlated with some aspects of health-related quality of life, more so with the disease-specific than the generic tool-existing generic instruments are of limited application in the assessment of satisfaction with treatment.

There is at least a theoretical association between satisfaction and quality of life [3] and our study did find that the *global satisfaction* score showed moderate convergence with the *treatment satisfaction* domain in the SAQ. Even so, the association between satisfaction and health-related quality of life is not straightforward. An inverse relationship cannot be ruled out [3], nor can a causal relationship. In patients with mental disorders, the relationship may be direct [30] or inverse. [6]

**Table 4: Correlation between HNA 'satisfaction' items and HRQL variables**

	EQ-Mob	EQ-SC	EQ-UA	EQ-PD	EQ-AD	EQ-VAS	SAQ Phys	SAQ AS	SAQ AF	SAQ TS	SAQ QOL	PCS	MCS	HNA SAT
Getting an appointment	-0.06	<b>-.15*</b>	-0.06	-.11	<b>-.32</b>	<b>.19</b>	.06	.09	<b>.13*</b>	<b>.31</b>	<b>.24</b>	.10	<b>.22</b>	<b>.73</b>
Enough time in GP visits	-0.03	-0.06	-0.07	-0.05	<b>-.24</b>	.12	.09	-0.03	.10	<b>.33</b>	<b>.14*</b>	.04	<b>.20</b>	<b>.70</b>
Satisfaction with referral	.03	<b>-.18</b>	.02	.01	<b>-.24</b>	.07	.04	-0.05	.04	<b>.37</b>	<b>.17*</b>	.08	<b>.22</b>	<b>.67</b>
GP service satisfaction	-0.02	-0.08	-0.06	-0.04	<b>-.14</b>	.03	.05	-0.09	.08	<b>.34</b>	.07	.00	<b>.15*</b>	<b>.65</b>
Satisfied with current treatment	<b>-.21</b>	<b>-.20</b>	<b>-.15</b>	<b>-.28</b>	<b>-.35</b>	<b>.32</b>	<b>.24</b>	<b>.15*</b>	<b>.21</b>	<b>.65</b>	<b>.43</b>	<b>.27</b>	<b>.36</b>	<b>.68</b>
Satisfied with GP advice	-.11	-.10	<b>-.14*</b>	-0.03	<b>-.22</b>	.10	.05	.00	.10	<b>.45</b>	<b>.19</b>	.10	<b>.21</b>	<b>.75</b>
Satisfied with Nurse services	-.12	-0.09	<b>-.14</b>	-0.05	<b>-.23</b>	<b>.22</b>	.06	.00	.07	<b>.42</b>	<b>.28</b>	.07	<b>.25</b>	<b>.73</b>
Satisfied with Rehabilitation services	<b>-.22</b>	<b>-.23</b>	<b>-.17*</b>	-.12	<b>-.28</b>	<b>.23</b>	<b>.22</b>	.06	.02	<b>.36</b>	<b>.29</b>	<b>.21</b>	<b>.36</b>	<b>.67</b>
Satisfied with consultant services	-0.09	-0.05	-0.10	-0.12	<b>-.16*</b>	<b>.16*</b>	.09	.06	-0.02	<b>.43</b>	<b>.22</b>	.09	<b>.24</b>	<b>.65</b>
Need easier access to health services	-0.10	<b>-.20</b>	<b>-.17</b>	-0.12	<b>-.33</b>	<b>.26</b>	<b>.20</b>	.05	.11	<b>.30</b>	<b>.22</b>	<b>.20</b>	<b>.28</b>	<b>.55</b>
GP understands needs?	-0.09	-0.07	-0.05	-0.00	<b>-.15*</b>	.07	-0.01	-0.00	.06	<b>.36</b>	<b>.19</b>	.02	<b>.19</b>	<b>.61</b>
HNA global satisfaction score	<b>-.15*</b>	<b>-.21</b>	<b>-.18</b>	<b>-.17</b>	<b>-.40</b>	<b>.28</b>	<b>.18</b>	.05	<b>.15*</b>	<b>.55</b>	<b>.35</b>	<b>.20</b>	<b>.38</b>	-

\* Correlation is significant at the .05 level and bold if  $p < 0.01$  (2-tailed).

Patient satisfaction is dependent upon a variety of personal, cultural, social, socio-economic and health-related factors, set against a background of previous exposure to, and experience of, health care services. Many of these may not be readily amenable to change, but where deficiencies are identified, such as in this study, alterations in services might well be rewarded with more satisfied patients.

## References

- DoH: **Patient and public involvement in the new NHS**. Department of Health; 1999.
- Ware J. E., Jr., Davies-Avery A and Stewart AL: **The measurement and meaning of patient satisfaction**. *Health Med Care Serv Rev* 1978, **1**:1, 3-15.
- Aharony L and Strasser S: **Patient satisfaction: what we know about and what we still need to explore**. *Med Care Rev* 1993, **50**:49-79.
- Von Essen L, Larsson G, Oberg K and Sjoden PO: **'Satisfaction with care': associations with health-related quality of life and psychosocial function among Swedish patients with endocrine gastrointestinal tumours**. *Eur J Cancer Care (Engl)* 2002, **11**:91-99.
- Fitzpatrick R: **Scope and measurement of patient satisfaction**. *Measurement of Patients' Satisfaction with Their Care* Edited by: Fitzpatrick R and Hopkins A. London, Royal College of Physicians of London; 1993:1-17.
- Guldvog B: **Can patient satisfaction improve health among patients with angina pectoris?** *Int J Qual Health Care* 1999, **11**:233-240.
- Donabedian A: **The quality of care. How can it be assessed?** *Jama* 1988, **260**:1743-1748.
- Oldridge N, Gottlieb M, Guyatt G, Jones N, Streiner D and Feeny D: **Predictors of health-related quality of life with cardiac rehabilitation after acute myocardial infarction**. *J Cardiopulm Rehabil* 1998, **18**:95-103.
- Thompson DR, Meadows KA and Lewin RJ: **Measuring quality of life in patients with coronary heart disease**. *Eur Heart J* 1998, **19**:693-695.
- Gray D and Hampton JR: **Twenty years' experience of myocardial infarction: the value of a heart attack register**. *Br J Clin Pract* 1993, **47**:292-295.
- Gray D, Keating NA, Murdock J, Skene AM and Hampton JR: **Impact of hospital thrombolysis policy on out-of-hospital response to suspected myocardial infarction**. *Lancet* 1993, **341**:654-657.
- Asadi-Lari M, Packham C and Gray D: **Unmet health needs in patients with coronary heart disease: implications and potential for improvement in caring services**. *Health Qual Life Outcomes* 2003, **1**:26.
- Asadi-Lari M and Gray D: **Is quality of life measurement likely to be a proxy for health needs assessment in patients with coronary artery disease?** *Health Qual Life Outcomes* 2003, **1**:50.
- Spertus JA, Winder JA, Dewhurst TA, Deyo RA and Fihn SD: **Monitoring the quality of life in patients with coronary artery disease**. *Am J Cardiol* 1994, **74**:1240-1244.
- Dougherty CM, Dewhurst T, Nichol WP and Spertus J: **Comparison of three quality of life instruments in stable angina pectoris: Seattle Angina Questionnaire, Short Form Health Survey (SF-36), and Quality of Life Index-Cardiac Version III**. *J Clin Epidemiol* 1998, **51**:569-575.
- Brazier JE, Harper R, Jones NM, O'Cathain A, Thomas KJ, Usherwood T and Westlake L: **Validating the SF-36 health survey questionnaire: new outcome measure for primary care**. *Brmj* 1992, **305**:160-164.
- Melville M, Asadi-Lari M, Brown N, Young T, Hampton JR and Gray D: **Quality of life assessment using the SF-12 is as reliable and sensitive as the SF-36 in distinguishing symptom severity in myocardial infarction survivors**. *Heart* 2003, **89**:1445-1446.
- Jenkinson C, Layte R, Jenkinson D, Lawrence K, Petersen S, Paice C and Stradling J: **A shorter form health survey: can the SF-12 replicate results from the SF-36 in longitudinal studies?** *J Public Health Med* 1997, **19**:179-186.
- Kind P, Dolan P, Gudex C and Williams A: **Variations in population health status: results from a United Kingdom national questionnaire survey**. *Brmj* 1998, **316**:736-741.
- van Agt HM, Essink-Bot ML, Krabbe PF and Bonsel GJ: **Test-retest reliability of health state valuations collected with the Euro-Qol questionnaire**. *Soc Sci Med* 1994, **39**:1537-1544.
- Johnson JA and Coons SJ: **Comparison of the EQ-5D and SF-12 in an adult US sample**. *Qual Life Res* 1998, **7**:155-166.

22. **Standard occupational classification / Office for National Statistics.** London, Stationery Office; 2000.
23. Munro HB: **Statistical methods for health care research.** 4thth edition. Philadelphia, Lippincott Williams & Wilkins; 2001.
24. Nabati L, Shea N, McBride L, Gavin C and Bauer MS: **Adaptation of a simple patient satisfaction instrument to mental health: psychometric properties.** *Psychiatry Res* 1998, **77**:51-56.
25. Bredart A, Razavi D, Robertson C, Batel-Copel L, Larsson G, Lichosik D, Meyza J, Schraub S, von Essen L and de Haes JC: **A comprehensive assessment of satisfaction with care: preliminary psychometric analysis in French, Polish, Swedish and Italian oncology patients.** *Patient Educ Couns* 2001, **43**:243-252.
26. Larson CO, Nelson EC, Gustafson D and Batalden PB: **The relationship between meeting patients' information needs and their satisfaction with hospital care and general health status outcomes.** *Int J Qual Health Care* 1996, **8**:447-456.
27. Pascoe GC: **Patient satisfaction in primary health care: a literature review and analysis.** *Eval Program Plann* 1983, **6**:185-210.
28. Chin MH and Goldman L: **Gender differences in 1-year survival and quality of life among patients admitted with congestive heart failure.** *Med Care* 1998, **36**:1033-1046.
29. Riedinger MS, Dracup KA, Brecht ML, Padilla G, Sarna L and Ganz PA: **Quality of life in patients with heart failure: do gender differences exist?** *Heart Lung* 2001, **30**:105-116.
30. Wiersma D and van Busschbach J: **Are needs and satisfaction of care associated with quality of life? An epidemiological survey among the severely mentally ill in the Netherlands.** *Eur Arch Psychiatry Clin Neurosci* 2001, **251**:239-246.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

