

EFFECT OF ASPIRIN, CLOPIDOGREL AND DIPYRIDAMOLE ON SOLUBLE MARKERS OF VASCULAR FUNCTION IN NORMAL VOLUNTEERS AND PATIENTS WITH PRIOR ISCHAEMIC STROKE

Lian Zhao,^{1,2} Laura Gray,^{1,2} Jo Leonardi-Bee,^{1,2} Chris S Weaver,^{1,2} Stan Heptinstall,² Philip M W Bath^{1,2}

Institutes of Neuroscience¹ & Clinical Research,² University of Nottingham, Nottingham UK

Correspondence: Professor Philip Bath
Division of Stroke Medicine
University of Nottingham
Queen's Medical Centre
Nottingham NG7 2UH

Tel: 0115 970 9348

Fax: 0115 875 4506

Email: philip.bath@nottingham.ac.uk

Keywords: aspirin, clopidogrel, dipyridamole, leucocyte, platelet, stroke

Although the mechanisms of action by which aspirin, clopidogrel and dipyridamole inhibit platelets are well characterised, their effects on soluble modulators of thrombosis, inflammation, and endothelial function have yet to be assessed systematically. In this investigation aspirin (A), clopidogrel (C), and dipyridamole (D) were administered singly and in combination (A, C, D, AC, AD, CD, ACD) in random order for two weeks (without washout) to 11 healthy subjects and 11 patients with previous ischaemic stroke. At the end of each treatment period plasma cyclic guanosine monophosphate (cGMP), monocyte chemoattractant peptide-1 (MCP-1), nitric oxide metabolites (NOx), plasminogen activator inhibitor-1 (PAI-1) and von Willebrand factor (vWf); and serum C-reactive protein (CRP) and platelet derived growth factor (PDGF); were measured blinded to treatment. Dipyridamole reduced plasma vWf levels (%) in both volunteers, -10.0 (5.0), and patients, -10.1 (4.3) ($p < 0.05$). Dipyridamole also lowered CRP (mg/L) in patients, -0.96 (0.47), but not volunteers. Clopidogrel reduced PAI-1 (ng/mL) in volunteers, -5.30 (2.20) ($p < 0.05$), and patients, -3.61 (2.75) (non-significant trend). Aspirin lowered PDGF (ng/mL) in volunteers, -3.46 (1.55) ($p < 0.05$), but not patients. Triple antiplatelet therapy was superior to dual and mono therapy in reducing vWf levels. In conclusion, antiplatelet agents have non-platelet-related effects on soluble modulators of thrombosis, inflammation, and endothelial function. In particular, dipyridamole reduces plasma vWf and clopidogrel lowers plasma PAI-1 levels. These effects may explain, in part, their roles in preventing atherothrombogenesis.

INTRODUCTION

The three antiplatelet agents, aspirin, clopidogrel and dipyridamole, are widely used for secondary stroke prevention on the basis of results from large trials and systematic reviews.^{1, 2} Whilst their antiplatelet actions are well described, their effects on other vascular cells and soluble factors are less well defined.

Platelets, leucocytes, and endothelial cells contribute to the pathogenesis of vascular disease³ through direct cell-cell interactions and the release of atherothrombogenic factors. Platelets release several cytokines including platelet-derived growth factor (PDGF), platelet factor 4 and CD40 ligand, each of which has been linked to the inflammation and coagulation pathways.⁴⁻⁶ Similarly, monocytes and macrophages generate cytokines including macrophage chemoattractant protein-1 (MCP-1), a chemokine important during atherogenesis,⁷ and interleukin-6 (IL-6),⁸ a procoagulant and stimulator of hepatic synthesis of C-reactive protein (CRP); CRP amplifies inflammatory and procoagulant responses.^{9,10} IL-6 also increases endothelial production of the fibrinolytic inhibitor, plasminogen activator inhibitor type 1 (PAI-1).^{3, 11} Endothelial cells also release von Willebrand factor (vWf), which facilitates platelet and thrombus adhesion to endothelium,¹² and nitric oxide (NO), a potent vasorelaxant, antiplatelet and antileucocyte agent¹³ whose effects are mediated by the second messenger cyclic guanosine monophosphate (cGMP).¹⁴

We assessed the effects of aspirin, clopidogrel and dipyridamole on a panel of the above soluble markers of vascular function in normal volunteers and patients with a prior history of ischaemic stroke. We also assessed the effect of combining the three antiplatelet agents on the soluble markers since we have shown previously that three drugs may be superior to one or two in inhibiting platelet function *in vitro*.¹⁵

METHODS

Design

Two randomised, outcome-blinded, multiway, crossover trials of antiplatelet therapy were performed in normal volunteers and patients with prior ischaemic stroke. The study design has been published previously¹⁶ in a report of the effect of aspirin, clopidogrel and dipyridamole on the function of circulating platelets, phagocytes and their conjugates.

Subjects

The study protocol was approved by the local Research Ethics Committee. All subjects gave written informed consent and the trials were performed according to the Declaration of Helsinki and the principles of Good Clinical Practice. Eleven healthy volunteers without any history of vascular disease were recruited from hospital or university staff or their friends. Eleven stable patients with a prior history of ischaemic stroke (on clinical and neuroimaging criteria) within 5 years were recruited from the stroke service at Nottingham City Hospital;¹⁶ each patient was receiving aspirin for vascular prophylaxis. Subjects were excluded if they had a history of cerebral haemorrhage, gastrointestinal bleeding, peptic ulcer, anaemia, thrombocytopenia, any severe concomitant medical condition, or hypersensitivity or intolerance to any of the study drugs; were taking anticoagulation or a non-steroidal anti-inflammatory drug; had a blood pressure > 180/110 mmHg; or were pregnant or lactating women.

Interventions

All subjects received two-week periods of open-label aspirin (A, 75mg daily), clopidogrel (C, 75mg daily) or modified release dipyridamole (D, 200mg twice daily). The drugs were given either singly (A, C, or D), in pairs (AC, AD, or CD) or all three together (ACD) in random order.¹⁶ Normal volunteers, but not patients, were also studied off all therapy. No washout periods were used since the length of each treatment phase (14 days) exceeded the lifespan of platelets (~10 days); further, it would have been inappropriate to deprive patients of antiplatelet therapy during such periods. Randomisation was performed using a computer and the treatment codes held by the hospital's Pharmacy to ensure concealment of allocation. Blood samples were collected at the end of each two-week treatment phase. Adverse events are reported previously.¹⁶ Subjects who withdrew from treatment were replaced to ensure

that complete data were present for each person; laboratory data from withdrawing subjects are not included in the statistical analyses.

Laboratory study

Serum and plasma collected into 3.8% sodium citrate (1:9 volume) or 4.0 mmol/L EDTA were centrifuged (1,500g for 15min) within 1 hour of venepuncture. Aliquots were frozen at -80°C and then assayed in batches using commercially available ELISA kits. CRP (high sensitivity kit, Kalon Biological, Hants, UK) and PDGF (R&D System, Abington, UK) were measured in serum; cGMP (R&D System), vWf (Corgenix, Cambridgeshire, UK), MCP-1 (R&D System) and PAI-1 (Technoclone TC, Surrey, UK) were analysed in citrated plasma.

Plasma (citrated) nitric oxide (NO_x) levels were determined from measurement of its stable end products, nitrate and nitrite, by a chemiluminescence (Sievers 280 Nitric Oxide Analyser, Analytix Limited, County Durham, UK) method.¹⁷ This involved injecting 10 μL of deproteinised plasma samples into vanadium chloride.¹⁸ The nitrate and nitrite present in samples were reduced to NO, and quantified from a calibration curve based on prepared concentrations of sodium nitrate.

Statistical analysis

The primary analysis compared the presence and absence of each drug on soluble markers, i.e. aspirin versus no aspirin, clopidogrel versus no clopidogrel, and dipyridamole versus no dipyridamole. Secondary analyses compared triple therapy (ACD) versus all other drugs, as before.¹⁶ Analyses were performed using the SAS statistical package.

RESULTS

Demographic and clinical information for the subjects are given in table 1. The stroke patients are representative of our clinical stroke service and were significantly older and had more risk factors (more hypertension and smokers, higher systolic blood pressure, and elevated total cholesterol) than the volunteers (all $p < 0.05$).

When considering the individual effects of the three antiplatelet agents on soluble markers of vascular function, three observations were made. First, dipyridamole significantly reduced plasma vWf levels in both volunteers and stroke patients (table 2). Second, dipyridamole significantly reduced serum CRP in patients but not volunteers (where a non-significant increase was seen). Third, clopidogrel reduced plasma PAI-1 concentrations in volunteers (significant) and patients (non-significant trend). Fourth, aspirin significantly reduced serum PDGF levels in volunteers but not patients. The three antiplatelet agents did not appear to alter MCP-1, NOx or cGMP levels in either subject group.

Patients had significantly lower levels of plasma NOx (difference 0.37, SD 0.17, $p < 0.001$) and higher levels of plasma vWf (difference -43.1, SD 39.5, $p = 0.019$) than volunteers (table 3); no differences in the other biomarkers were apparent. The use of triple antiplatelet therapy was not superior to mono or dual therapy with the exception that the combination of all three agents was associated with significantly lower levels of plasma vWf than dual aspirin and clopidogrel in both volunteers and patients (table 3), a finding compatible with the above observation that dipyridamole lowers vWf levels.

DISCUSSION

These studies show that antiplatelet agents may alter soluble markers of vascular function. In particular, dipyridamole significantly reduced plasma vWf levels in both volunteers and patients with previous stroke, an effect seen previously in an independent study involving patients with ischaemic heart disease.¹⁹

Studies *in vitro* and *ex vivo* have found that dipyridamole significantly reduces platelet adhesion to vascular sub-endothelium although the mechanism remains unclear.^{20, 21} Both vWf and P-selectin are stored in Weibel-Palade body in the endothelium.^{22, 23} Pro-inflammatory mediators activate cells causing Weibel-Palade bodies to fuse with the cell membrane resulting in the release of vWf and surface exposure of P-selectin. vWf and P-selectin are the primary molecules involved with platelet interaction with endothelium.^{23, 24} In the present study we provide evidence that the inhibitory effect of dipyridamole on platelet adhesion may be mediated in part through inhibition of vWf release from endothelial cells.

Dipyridamole also appeared to lower serum CRP levels, a previously unreported finding. However, this finding was only present in patients with prior cerebrovascular disease and could simply reflect chance. Several studies reported that dipyridamole potentiates NO-induced vasodilator activity through inhibiting phosphodiesterase V activity and thereby enhancing intracellular cGMP levels.²⁵⁻²⁷ However, we did not observe changes in plasma NOx and cGMP levels with dipyridamole.

Clopidogrel showed an inhibitory effect on plasma PAI-1 levels (significant in volunteers, non-significant in patients) but not on other biomarkers. The reduction of PAI-1 has been observed in two earlier studies^{28, 29} and may be achieved partially through inhibition of platelet activity.^{30, 31} Previous studies reported that clopidogrel lowers CRP levels in patients with acute vascular disease.³²⁻³⁴ However, we did not confirm this observation, perhaps because CRP is an acute phase protein and is elevated significantly in acute rather than chronic vascular disease.

Aspirin reduced serum PDGF levels but only in volunteers; this finding has not been reported before and might reflect chance. A reduction in CRP with aspirin was found in an earlier study of patients with hypertension³⁵ but not in studies involving healthy subjects.^{36, 37} Apart from differences in subjects, this positive study involved prolonged

treatment over 18 weeks whereas aspirin was given for shorter periods (<1 month) in the above two neutral studies and ours.

We found, in secondary analyses, that plasma NOx levels were lower, and vWf levels higher, in patients with cerebrovascular disease than in volunteers. Both factors are produced by endothelial cells and the results support previous finding that endothelial dysfunction is present in patients with vascular disease. For example, patients with hypertension, hypercholesterolaemia, diabetes mellitus and previous clinical vascular events have lower NOx and elevated vWf levels.^{38, 39} Similarly, acute stroke is associated with low NOx and high vWf levels.^{18, 40}

Although our trials were small, we used a crossover design (thereby allowing each subject to act as their own control) which considerably increases statistical power as compared with a parallel group design. Additionally, we assessed the effects of antiplatelet agents in two different sets of subjects which increases the validity of the findings. Importantly, measurements were made by one person who was blinded to treatment.

In summary, dipyridamole reduced plasma vWf levels, a measure of endothelial dysfunction, in both normal subjects and patients with previous ischaemic stroke. Similarly, clopidogrel lowered plasma PAI-1 concentrations in both sets of subjects (significantly so in volunteers). The findings suggest that these agents have additional effects on non-platelet aspects of atherothrombogenesis, mechanisms which may explain, in part, their roles in preventing recurrent vascular events such as stroke. The effects of these agents on other vascular risk markers such as fibrinogen and P-selectin remains to be ascertained.⁴⁰

ACKNOWLEDGEMENTS

We thank the volunteers and patients who took part in the study. PB is Stroke Association Professor of Stroke Medicine; the Division of Stroke Medicine receives core funding from The Stroke Association.

REFERENCES

1. Antithrombotic Trialists Collaboration. Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. *BMJ* 2002; 324: 71-86.
2. Leonardi-Bee J, Bath P M, Bousser M G, Davalos A, Diener H-C, Guiraud-Chaumeil B, Sivenius J, Yatsu F, Dewey M E, and Dipyridamole in Stroke Collaboration (DISC). Dipyridamole for preventing recurrent ischemic stroke and other vascular events: a meta-analysis of individual patient data from randomized controlled trials. *Stroke* 2005; 36: 162-8.
3. Libby P and Simon D I. Inflammation and Thrombosis: The Clot Thickens. *Circulation* 2001; 103: 1718-1720.
4. Heldin C H and Westermark B. Mechanism of action and in vivo role of platelet-derived growth factor. *Physiol Rev* 1999; 79: 1283-316.
5. Marcum J A, McKenney J B, and Rosenberg R D. Acceleration of thrombin antithrombin complex formation in rat hindquartes via heparin-like molecules bound to the endothelium. *J Clin Invest* 1984; 74: 341.
6. Aukrust P, Damas J K, and Solum N O. Soluble CD40 ligand and platelets: self-perpetuating pathogenic loop in thrombosis and inflammation? *J Am Coll Cardiol* 2004; 43: 2326-8.
7. Reape T J and Groot P H. Chemokines and atherosclerosis. *Atherosclerosis* 1999; 147: 213-25.
8. Willerson J T and Ridker P M. Inflammation as a cardiovascular risk factor. *Circulation* 2004; 109: II2-10.
9. Pasceri V, Willerson J T, and Yeh E T H. Direct Proinflammatory Effect of C-Reactive Protein on Human Endothelial Cells. *Circulation* 2000; 102: 2165-2168.
10. Willerson J T. Systemic and local inflammation in patients with unstable atherosclerotic plaques. *Prog Cardiovasc Dis* 2002; 44: 469-78.
11. Devaraj S, Xu D Y, and Jialal I. C-Reactive Protein Increases Plasminogen Activator Inhibitor-1 Expression and Activity in Human Aortic Endothelial Cells: Implications for the Metabolic Syndrome and Atherothrombosis. *Circulation* 2003; 107: 398-404.
12. Ruggeri Z M. Role of von Willebrand factor in platelet thrombus formation. *Ann Med* 2000; 32 Suppl 1: 2-9.
13. Palmer R M, Ferrige A G, and Moncada S. Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature* 1987; 327: 524-6.

14. Rapoport R M, Draznin M B, and Murad F. Endothelium-dependent relaxation in rat aorta may be mediated through cyclic GMP-dependent protein phosphorylation. *Nature* 1983; 306: 174-6.
15. Zhao L, Bath P, and Heptinstall S. Effects of combining three different antiplatelet agents on platelets and leukocytes in whole blood in vitro. *Br J Pharmacol* 2001; 134: 353-358.
16. Zhao L, Fletcher S, Weaver C, Leonardi-Bee J, May J, Fox S, and Willmot M. Effects of aspirin, clopidogrel and dipyridamole administered singly and in combination on platelet and leucocyte function in normal volunteers and patients with prior ischaemic stroke. *Thromb Haemost* 2005; 93: 527-34.
17. Menon N, Wolf A, Zehetgruber M, and Bing R. An improved chemiluminescence assay suggests non nitric oxide mediated action of lysophosphatidylcholine and acetylcholine. *Proc Soc Exp Biol Med* 1989; 191: 316-319.
18. Rashid P A, Whitehurst A, Lawson N, and Bath P M W. Plasma nitric oxide (nitrate/nitrite) levels in acute stroke and their relationship with severity and outcome. *J Stroke Cerebrovasc Dis* 2003; 12: 82-87.
19. Sano T, Motomiya T, and Yamazaki H. Platelet release reaction in vivo in patients with ischaemic heart disease after isometric exercise and its prevention with dipyridamole. *Thromb Haemost* 1980; 42: 1589-97.
20. Lauri D, Zanetti A, and Dejana E. Effects of dipyridamole and low-dose aspirin therapy on platelet adhesion to vascular subendothelium. *Am J Cardiol* 1986; 58: 1261-1264.
21. Eisert W G. Near-field amplification of antithrombotic effects of dipyridamole through vessel walls cells. *Neurology* 2001; 57 (suppl 2): S20-S23.
22. Wagner D D, Olmsted J B, and Marder V J. Immunolocalization of von Willebrand protein in Weibel-Palade bodies of human endothelial cells. *J Cell Biol* 1982; 95: 355-60.
23. Bonfanti R, Furie B C, Furie B, and Wagner D D. PADGEM (GMP140) is a component of Weibel-Palade bodies of human endothelial cells. *Blood* 1989; 73: 1109-12.
24. Andre P, Denis C V, Ware J, Saffaripour S, Hynes R O, Ruggeri Z M, and Wagner D D. Platelets adhere to and translocate on von Willebrand factor presented by endothelium in stimulated veins. *Blood* 2000; 96: 3322-8.
25. Bult H, Fret H R, Jordaens F H, and Herman A G. Dipyridamole potentiates the anti-aggregating and vasodilator activity of nitric oxide. *Eur J Pharmacol* 1991; 199: 1-8.
26. Aktas B, Utz A, Hoenig-Liedl P, Walter U, and Geiger J. Dipyridamole enhances NO/cGMP-mediated vasodilator-stimulated phosphoprotein phosphorylation and signaling in human platelets: in vitro and in vivo/ex vivo studies. *Stroke* 2003; 34: 764-9.

27. Moreno L, Losada B, Cogolludo A L, Lodi F, Lugnier C, Villamor E, Moro M, Tamargo J, and Perez-Vizcaino F. Postnatal Maturation of Phosphodiesterase 5 (PDE5) in Piglet Pulmonary Arteries: Activity, Expression, Effects of PDE5 Inhibitors, and Role of the Nitric Oxide/Cyclic GMP Pathway. *Pediatr Res* 2004;
28. Tohgi H, Takahashi H, Chiba K, and Tamura K. Coagulation-fibrinolysis system in poststroke patients receiving antiplatelet medication. *Stroke* 1993; 24: 801-4.
29. Peters H, Eisenberg R, Daig U, Liefeldt L, Westenfeld R, Gaedeke J, Kramer S, and Neumayer H H. Platelet inhibition limits TGF-beta overexpression and matrix expansion after induction of anti-thy1 glomerulonephritis. *Kidney Int* 2004; 65: 2238-48.
30. Konkle B A, Schick P K, He X, Liu R J, and Mazur E M. Plasminogen activator inhibitor-1 mRNA is expressed in platelets and megakaryocytes and the megakaryoblastic cell line CHRF-288. *Arterioscler Thromb* 1993; 13: 669-74.
31. Mousa S A, Bozarth J, Forsythe M, Tsao P, Pease L, and Reilly T M. Role of platelet GpIIb/IIIa receptors in the modulation of platelet plasminogen activator inhibitors type-1 (PAI-1) release. *Life Sci* 1994; 54: 1155-62.
32. Cha J K, Jeong M H, Lee K M, Bae H R, Lim Y J, Park K W, and Cheon S M. Changes in platelet P-selectin and in plasma C-reactive protein in acute atherosclerotic ischemic stroke treated with a loading dose of clopidogrel. *J Thromb Thrombolysis* 2002; 14: 145-50.
33. Woodward M, Lowe G D, Francis L M, Rumley A, and Cobbe S M. A randomized comparison of the effects of aspirin and clopidogrel on thrombotic risk factors and C-reactive protein following myocardial infarction: the CADET trial. *J Thromb Haemost* 2004; 2: 1934-40.
34. Vivekananthan D P, Bhatt D L, Chew D P, Zidar F J, Chan A W, Moliterno D J, Ellis S G, and Topol E J. Effect of clopidogrel pretreatment on periprocedural rise in C-reactive protein after percutaneous coronary intervention. *Am J Cardiol* 2004; 94: 358-60.
35. Monobe H, Yamanari H, Nakamura K, and Ohe T. Effects of low-dose aspirin on endothelial function in hypertensive patients. *Clin Cardiol* 2001; 24: 705-9.
36. Feng D, Tracy R P, Lipinska I, Murillo J, McKenna C, and Tofler G H. Effect of short-term aspirin use on C-reactive protein. *J Thromb Thrombolysis* 2000; 9: 37-41.
37. Feldman M, Jialal I, Devaraj S, and Cryer B. Effects of low-dose aspirin on serum C-reactive protein and thromboxane B2 concentrations: a placebo-controlled study using a highly sensitive C-reactive protein assay. *J Am Coll Cardiol* 2001; 37: 2036-41.
38. Correa R C and Alfieri A B. Plasmatic nitric oxide, but not von Willebrand Factor, is an early marker of endothelial damage, in type 1 diabetes mellitus patients without microvascular complications. *J Diabetes Complications* 2003; 17: 264-8.

39. Blann A D, Dobrotova M, Kubisz P, and McCollum C N. von Willebrand factor, soluble P-selectin, tissue plasminogen activator and plasminogen activator inhibitor in atherosclerosis. *Thromb Haemost* 1995; 74: 626-630.
40. Bath P M W, Blann A, Smith N, and Butterworth R J. Von Willebrand factor, p-selectin and fibrinogen levels in patients with acute ischaemic and haemorrhagic stroke, and their relationship with stroke sub-type and functional outcome. *Platelets* 1998; 9: 155-159.

TABLE 1

Baseline characteristics of subjects. Mean (standard deviation) or frequency (%)

	Normal volunteers	Previous ischaemic stroke	P
Number	11	11	
Mean age (year)	39 (8)	62 (10)	<0.0001
Gender, male (%)	8 (73)	6 (55)	0.38
Hypertension (%)	0 (0)	8 (73)	0.0004
Diabetes mellitus (%)	0 (0)	0 (0)	1.0
Smoking (%)			0.03
Never	9 (82)	3 (27)	
Past	1 (9)	5 (45)	
Current	1 (9)	3 (27)	
Blood pressure (mmHg)			
Systolic	130 (14)	150 (20)	0.01
Diastolic	85 (4)	86 (8)	0.71
Platelet count ($\times 10^9$.l)	257 (35)	282 (71)	0.31
Total cholesterol (mmol.l ⁻¹)	5.0 (0.7)	5.9 (0.8)	0.01
LDL-C (mmol.l ⁻¹)	3.4 (0.6)	3.8 (0.8)	0.20
HDL-C (mmol.l ⁻¹)	1.0 (0.2)	1.4 (0.4)	0.01
Triglycerides (mmol.l ⁻¹)	1.3 (0.5)	2.2 (1.5)	0.07

LDL-C: low-density lipoprotein cholesterol; HDL-C: high-density lipoprotein cholesterol

TABLE 2

Effect of aspirin (A), clopidogrel (C) and dipyridamole (D) on soluble markers; mean difference (standard deviation); p<0.05 in bold

	cGMP (nM)	CRP (mg/L)	MCP-1 (pg/mL)	NOx (μmol/L)	PAI-1 (ng/mL)	PDGF (ng/mL)	vWf (%)
Volunteers							
A vs no A	0.30 (0.43)	0.52 (1.05)	9.59 (15.97)	-0.06 (0.05)	0.73 (2.20)	-3.46 (1.55)	3.48 (4.95)
C vs no C	-0.27 (0.43)	-0.64 (1.05)	4.45 (15.97)	0.02 (0.05)	-5.30 (2.20)	-1.89 (1.55)	3.16 (4.95)
D vs no D	-0.55 (0.43)	0.23 (1.05)	1.21 (15.97)	-0.001 (0.05)	-0.10 (2.20)	1.70 (1.55)	-10.01 (4.95)
Patients							
A vs no A	0.39 (0.57)	0.06 (0.47)	9.02 (10.50)	-0.04 (0.06)	-3.77 (2.75)	0.44 (1.98)	2.90 (4.34)
C vs no C	0.59 (0.57)	0.05 (0.47)	16.97 (10.50)	0.04 (0.06)	-3.61 (2.75)	0.72 (1.98)	1.88 (4.34)
D vs no D	-0.28 (0.57)	-0.96 (0.47)	-0.64 (10.50)	-0.03 (0.06)	0.42 (2.75)	-0.71 (1.98)	-10.11 (4.34)

cGMP: cyclic guanylate monophosphate; CRP: c-reactive protein; MCP-1: macrophage chemoattractant peptide; NOx: nitric oxide metabolites; PAI-1: plasminogen activator inhibitor-1; PDGF: platelet derived growth factor; vWf: von Willebrand factor

TABLE 3

Effect of aspirin (A), clopidogrel (C) and dipyridamole (D) on soluble markers in normal volunteers and patients with previous ischaemic stroke: mean (standard deviation), *p<0.05 ACD versus other therapies, \$p<0.05 volunteers versus patients

	A	C	D	AC	AD	CD	ACD
Volunteers							
cGMP (nmol/L)	2.9 (3.4)	1.6 (1.3)	2.5 (2.1)	3.2 (2.8)	1.8 (2.0)	2.1 (2.5)	1.7 (1.7)
CRP (mg/L)	3.7 (6.3)	2.5 (2.8)	4.0 (6.5)	2.3 (2.0)	2.3 (2.8)	1.5 (1.0)	4.4 (6.9)
MCP-1 (pg/mL)	242 (123)	211 (81)	240 (115)	260 (137)	249 (102)	247 (107)	242 (90)
NOx (μmol/L)	0.72 (0.17)	0.95 (0.42)	0.83 (0.28)	0.83 (0.31)	0.91 (0.24)	0.81 (0.25)	0.77 (0.19)
PAI-1 (ng/mL)	32.1 (17.9)	25.2 (19.1)	29.7 (21.6)	23.4 (14.8)	27.8 (15.6)	24.4 (11.5)	25.3 (13.9)
PDGF (ng/mL)	21.5 (11.3)	23.8 (14.8)	25.6 (13.1)	20.8 (17.0)	24.4 (18.0)	23.9 (17.1)	18.8 (12.7)
vWf (%)	47.3 (46.4)	56.3 (35.0)	42.2 (18.4)	62.2 (37.7)*	53.7 (29.6)	44.2 (31.8)	40.9 (18.8)
Patients							
cGMP (nmol/L)	2.6 (3.4)	2.6 (2.9)	1.3 (1.9)	2.0 (3.2)	1.9 (2.4)	2.2 (2.5)	3.2 (2.8)
CRP (mg/L)	4.3 (4.6)	4.3 (3.1)	2.9 (2.6)	3.6 (2.2)	3.3 (2.6)	3.3 (2.9)	2.9 (2.3)
MCP-1 (pg/ml)	181 (49)	202 (54)	175 (49)	205 (43)	202 (61)	195 (42)	210 (49)
NOx (μmol/L)	0.35 (0.16) ^{\$}	0.38 (0.25)	0.34 (0.21)	0.36 (0.29)	0.28 (0.18)	0.38 (0.41)	0.34 (0.27)
PAI-1 (ng/mL)	36.7 (19.6)	34.3 (14.6)	37.8 (20.2)	29.5 (19.8)	32.8 (20.6)	35.5 (16.6)	29.5 (13.3)
PDGF (ng/mL)	22.8 (17.8)	21.7 (18.1)	22.0 (15.9)	22.7 (18.5)	19.9 (11.3)	21.5 (19.1)	23.3 (13.3)
vWf (%)	90.4 (31.0) ^{\$}	89.9 (38.4)	81.6 (32.3)	96.7 (41.1)*	84.5 (36.8)	83.2 (46.2)	79.7 (31.0)

cGMP: cyclic guanylate monophosphate; CRP: c-reactive protein; MCP-1: macrophage chemoattractant peptide; NOx: nitric oxide metabolites; PAI-1: plasminogen activator inhibitor-1; PDGF: platelet derived growth factor; vWf: von Willebrand factor