

# 1 **Impacts of changing climate and agronomic factors on fusarium ear blight** 2 **of wheat in the UK**

3  
4  
5  
6  
7  
8 *Jonathan S. WEST<sup>a\*</sup>, Sarah HOLDGATE<sup>a†</sup>, James A. TOWNSEND<sup>a</sup>, Julia B HALDER<sup>ad</sup>,*  
9  
10 *Simon G. EDWARDS<sup>b</sup>, Philip JENNINGS<sup>c</sup> and Bruce D. L. FITT<sup>a</sup>*

11  
12  
13  
14  
15 <sup>a</sup> Rothamsted Research, Harpenden, AL5 2JQ, UK; <sup>b</sup> Harper Adams University College,  
16  
17 Newport, TF10 8NB, UK; <sup>c</sup> The Food and Environment Research Agency, Sand Hutton,  
18  
19 York YO41 1LZ, UK; <sup>d</sup> Imperial College, London; <sup>†</sup> current address: RAGT Seeds Ltd.,  
20  
21 Grange Road, Ickleton, Saffron Walden, CB10 1TA, UK  
22  
23  
24

25  
26  
27  
28 \*E-mail: [jon.west@bbsrc.ac.uk](mailto:jon.west@bbsrc.ac.uk)  
29  
30  
31

32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

Climate change will have direct impacts on fusarium ear blight (FEB) in wheat crops, since weather factors greatly affect epidemics, the relative proportions of species of ear blight pathogens responsible and the production of deoxynivalenol (DON) toxin by two *Fusarium* species, *F. graminearum* and *F. culmorum*. Many established weather-based prediction models do not accurately predict FEB severity in the UK. One weather-based model developed with UK data suggests a slight increase in FEB severity under climate change. However, severity of the disease is likely to increase further due to increased cropping of grain maize, since maize debris is a potent source of inoculum of *F. graminearum*. Further research on forecasting, management options to reduce mycotoxin production and breeding for resistant varieties is a high priority for the UK. Management must also consider factors such as tillage regime, wheat cultivar (flowering time and disease resistance) and fungicide use, which also influence the severity of FEB and related toxin production.

26

27 *Keywords:* food security, mycotoxins, risk prediction, wheat scab, fusarium head blight,  
28 deoxynivalenol (DON)

29

30

## 31 **Introduction**

32

33 Climate change can increase the range and severity of plant disease epidemics (Garrett *et al.*,  
34 2006). Such increases can threaten global food security if they affect staple food crops in  
35 agricultural ecosystems especially in the developing world (Chakraborty *et al.*, 2000;  
36 Anderson *et al.*, 2004; Garrett *et al.*, 2006; Schmidhuber & Tubiello, 2007). Crop disease  
37 epidemics cause instability in food supply, which can lead to famine, conflicts and mass-  
38 movement of people to more favoured areas (Stern, 2007). Climate change can directly  
39 affect plant pathogens by providing a climate that is more or less favourable to the pathogen  
40 (for infection, colonisation, reproduction or dispersal). Several successive years of  
41 favourable climate can potentially cause a build-up of inoculum, leading to epidemics that are  
42 much more severe than when a single favourable year occurs. Furthermore, climate change  
43 can indirectly affect crop diseases, for example by provoking adaptation strategies that  
44 involve changes to crop rotations to include new crops that are additional hosts to particular  
45 pathogens.

46 The weather in the UK is relatively variable compared to that of continental European  
47 locations, with large differences in monthly rainfall typically occurring during a year despite  
48 the long term average monthly rainfall at any given location being almost the same each  
49 month. Figure 1 shows the annual mean temperature in different regions of the UK, which  
50 varied more with location than with year over the period 1914-2004. UK Climate Impacts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51 Programme (UKCIP; [www.ukcip.org.uk/](http://www.ukcip.org.uk/)) projections of future weather vary depending on  
52 which of many climate change scenarios are used but the general consensus is that the UK is  
53 predicted to have a warmer climate (e.g. +2 °C in winter to +4 °C in summer), with slightly  
54 wetter winters and drier summers (Semenov, 2009; Fig. 2). The UK is also predicted to  
55 experience much more intense weather events. These are the best estimates of future climate  
56 change available, although some alternative scenarios have been suggested.

57

58 Figures 1 and 2 here.

59

60 The UKCIP projected weather would advance the date of onset of wheat anthesis (by  
61 approximately 2 weeks by 2050) and maturity for harvest (by 3 weeks) (Semenov, 2009;  
62 Madgwick *et al.*, 2011). ‘Mediterranean-type’ wheat cultivars, which respond to different  
63 environmental cues determining the time of flowering, typically flower 2 weeks earlier than  
64 current UK cultivars. Adoption of this kind of cultivar in the UK has been proposed as an  
65 adaptation strategy to avoid heat stress at flowering could advance the time of flowering by at  
66 least another week to mid-May in southern England.

67 One serious disease of small grain cereals under the influence of climate change is  
68 fusarium ear blight (FEB), also known as fusarium head blight or scab, caused by a complex  
69 of many different species in the genera *Fusarium* and *Microdochium* (Xu & Nicholson,  
70 2009). Of these, two species, *Fusarium graminearum* (Schwabe) (teleomorph, *Gibberella*  
71 *zeae*) and *Fusarium culmorum* (Smith) Sacc. (no known teleomorph) are of concern in the  
72 UK because they produce a range of toxins that can contaminate grain, particularly  
73 deoxynivalenol (DON) (Fernandez & Chen, 2005; Parry *et al.*, 1995). Due to health  
74 concerns, EU legislation limits DON levels at 1250 µg kg<sup>-1</sup> in unprocessed wheat, with lower  
75 limits for various processed foods. DON concentrations of up to 600 mg kg<sup>-1</sup> in grain caused

76 by natural infections have been reported (Sinha & Savard, 1997), hence two or three heavily-  
77 infected grains per 1000 grains can make a batch close to the rejection limit. In the UK, in  
78 most years, few grain samples exceed DON thresholds in the UK [e.g. 0-5% of loads have  
79 exceeded thresholds since 2001, except in 2008 (10.2%) (<http://www.hgca.com/>)].  
80 Furthermore, the toxin zearalenone is also produced by *F. graminearum* and *F. culmorum*,  
81 while toxins such as HT2 and T2 are produced by other species of *Fusarium* and there are  
82 indications that strains or species that are favoured by warmer climates are more likely to  
83 produce some of these toxins (Jestoi *et al.*, 2009).

84 Problems associated with FEB are three-fold. Firstly, diseased ears that senesce  
85 prematurely (Fig. 3a) leads to shrivelling of infected grain (Fig. 3d) and substantial yield  
86 losses, ranging from 5-75% (Lin *et al.*, 2004; Parry *et al.*, 1995). There is also often an  
87 associated stem-base infection, fusarium foot-rot, caused by the same pathogens (Bateman *et*  
88 *al.*, 2007; Fig. 3c) that can contribute to these yield losses and act as a source of inoculum for  
89 ear infection. Secondly, the bread-making quality of infected wheat grains is also reduced  
90 due to their decreased protein and starch content (Parry *et al.*, 1995). Severely infected  
91 batches of grain may be rejected by millers on grounds of poor quality. The third effect is the  
92 contamination of grain by a range of mycotoxins, which is of great concern for safety of  
93 human food and animal feed (Wei *et al.*, 2005; Kostelanska *et al.*, 2009). In commercial  
94 practice, the lightest grains, which can be heavily infected, are usually removed at combining.  
95 However, despite post-harvest methods, such as optical sorting (Champeil *et al.*, 2004a) to  
96 reduce the impact of infected grain, FEB and contamination by mycotoxins remain major  
97 concerns.

98  
99 Figure 3 here\*\*\*\*\*  
100

101 DON is produced primarily during infection of the ear by *F. culmorum* and *F.*  
102 *graminearum* when water availability is high, which in the UK is usually before harvest,  
103 rather than in storage as farmers store grain at <15% (Gilbert & Tekauz, 2000; Jennings *et*  
104 *al.*, 2004a;b; Hope *et al.*, 2005; Chakraborty & Newton, 2011). A delay to harvest, caused by  
105 wet weather can substantially increase DON production (Anon 2009b). DON is thought to  
106 play a role in virulence (Jansen *et al.*, 2005; Proctor *et al.*, 1995; Maier *et al.*, 2006) and  
107 possibly (along with other trichothecenes) in competition against other fungi.

108 FEB disease occurs throughout the wheat growing regions of the world, with  
109 epidemics recorded in major cereal growing areas (McMullen *et al.*, 1997; Parry *et al.*, 1995).  
110 The two main DON-producing species, *F. graminearum* and *F. culmorum*, have slightly  
111 different temperature optima for growth, 24-28°C for *F. graminearum* and 20-25°C for *F.*  
112 *culmorum* (Doohan *et al.*, 2003; Brennan *et al.*, 2003). This small difference may explain  
113 why *F. graminearum* predominates in regions with relatively hot summers, such as the USA,  
114 Canada, Australia and parts of continental Europe, whereas *F. culmorum* is found in cooler  
115 maritime regions such as north western Europe (Moss, 2002; Parry *et al.*, 1995). However, in  
116 the UK, there appears to be no trend associated with mean temperature for years when *F.*  
117 *graminearum* has predominated over *F. culmorum* and vice versa (Anon. 2010). In the UK,  
118 these two species interact with other species responsible for FEB, principally *F. poae*, *F.*  
119 *avenaceum*, *Microdochium nivale* and *M. majus*, and there are other interacting species  
120 worldwide, many of which produce mycotoxins (Xu & Nicholson, 2009). Disease surveys in  
121 the UK have shown that the incidence of fusarium ear blight in the UK has been sporadic but  
122 has increased over the past decade (Fig. 4). Additionally, the incidence of *F. graminearum*  
123 has started to increase substantially since 2002 (www.cropmonitor.co.uk; Jennings *et al.*,  
124 2004b). During the 2006/2007 season, there was an unusually high incidence of FEB in the  
125 UK, with the principal causal agent being *M. nivale* (Fig. 4). A wet summer in 2008 was

126 associated with a relatively severe FEB epidemic and associated DON contamination, with  
127 more *F. graminearum* than *F. culmorum*. Before then, FEB had not been considered to pose  
128 a serious risk of DON contamination in the UK, particularly if good farming practices were  
129 followed. This review investigates the potential direct and indirect impacts of climate change  
130 on fusarium ear blight and subsequent mycotoxin contamination in the UK, particularly  
131 impacts of altered farming practices.

132

133 Figure 4 here.\*\*\*\*\*

134

135

### 136 **Disease-cycle of FEB**

137 *F. graminearum* and *F. culmorum* survive the inter-crop period on infected seed and crop  
138 debris in the field (Bateman *et al.*, 2007). For both *F. graminearum* and *F. culmorum*,  
139 asexually-produced macroconidia (Fig. 3b) formed in sporodochia are dispersed by rain  
140 splash, to infect florets at anthesis either directly from inoculum sources on the ground or via  
141 infections on intermediate leaf layers, that lead to secondary sporulation (Beyer *et al.*, 2005;  
142 Parry *et al.*, 1995; Paul *et al.*, 2004; Schmale *et al.*, 2005). Additionally, *F. graminearum*  
143 develops a sexual stage when conditions are warm and humid, forming blue or black  
144 ascospore-bearing perithecia on the surface of colonised debris such as chaff, grain and leaf  
145 litter, giving a ‘scabbed’ appearance (Doohan *et al.*, 2003; Sutton *et al.*, 1980; Goswami &  
146 Kistler, 2004; Kang & Buchenauer, 2000; Parry *et al.*, 1995). These discharge ascospores  
147 into the air in the spring for long-distance dispersal when the air temperature is decreasing  
148 and relative humidity (RH) increasing (i.e. typically in the evening) but not when RH is  
149 extremely high (Doohan *et al.*, 2003). Maldonado-Ramirez *et al.* (2005) assessed  
150 concentrations of viable *F. graminearum* propagules in air by impaction onto selective agar

151 plates on remotely operated pilotless drone aircraft. They found that airborne inoculum (of *F.*  
152 *graminearum*/*G. zea*) was abundant and well dispersed in air above New York state  
153 throughout the wheat flowering period (May/June) in each of four years. There was a  
154 homogeneous spatial pattern of propagule deposition detected by samplers located over a 1  
155 km scale (Schmale *et al.*, 2006).

156 Infection occurs when spores are deposited on the flowering wheat ear during periods  
157 of wetness or high humidity, with ideal conditions 25°C and 100% relative humidity for 24  
158 hours post inoculation (Abramson *et al.*, 1987; Parry *et al.*, 1995). Wheat plants are most  
159 susceptible during anthesis but infection can occur up to the soft dough stage (Windels,  
160 2000). Mycelium is able to infect florets in humid conditions by growing along the surface  
161 of spikelets and particularly on the anthers as they are extruded from the wheat ear during  
162 flowering. The initial symptoms of FEB are pale brown water soaked lesions on infected  
163 spikelets and at this stage there is often a substantial area of asymptomatic infection ahead of  
164 the visible disease symptoms (Brown *et al.*, 2010). Visible symptoms vary according to the  
165 species and to some extent the toxins produced by the infecting pathogen. In most cases the  
166 diseased areas of discolouration expand, causing spikelets to turn prematurely light brown.  
167 Bleaching that may occur on florets above the infection point (Fig 3a); is associated with  
168 DON production in some species (Maier *et al.*, 2006). Individual florets may become sterile,  
169 leading to poorly filled, shrivelled grain. Infected grains may have a chalky, floury interior  
170 and may show salmon-pink patches of conidia (Fig. 3d) along the edges of the glume and the  
171 base of the spikelet and the peduncle beneath the infected ear darkens (Ruckenbauer *et al.*,  
172 2001, [www.scabusa.org](http://www.scabusa.org)).

173

174

175 **Factors affecting severity of FEB and DON production**

176 **Weather factors**

177 Weather has been investigated as a factor affecting FEB in a number of studies. Xu *et al.*  
178 (2008) suggested that *Fusarium poae* was associated with relatively dry and warm  
179 conditions, *F. graminearum* with warm, humid conditions, *F. avenaceum* and *F. culmorum*  
180 with cool, wet or humid conditions, while two *Microdochium* species were associated with  
181 cool to moderate temperatures and frequent rain showers. Some forecasting models suggest  
182 that occurrence of rain and high humidity approximately one week before anthesis increases  
183 FEB risk and this is thought to be because this stimulates sporulation (de Wolf *et al.*, 2003;  
184 Hooker *et al.*, 2002; Moschini *et al.*, 2001). In addition, rain and, or high humidity at  
185 anthesis has been highly correlated with FEB incidence (Bateman *et al.*, 2007; de Wolf *et al.*,  
186 2003; Moschini *et al.*, 2001) but not with DON production (Hooker *et al.*, 2002). Warm (i.e.  
187 15°C to 30°C), wet weather at anthesis not only promotes infection but also encourages  
188 vegetative spread of mycelium to more florets (Parry *et al.*, 1995; deWolf *et al.*, 2003).  
189 However, in continental European locations, where hot conditions can occur, concentrations  
190 of DON decreased with increase in days exceeding 32°C (Hooker *et al.*, 2002). Weather  
191 factors (rain) contributing to a late harvest were also associated with increased DON  
192 contamination (Eiblmeier *et al.*, 2007).

193 A recent study that investigated impacts of climate change on Fusarium ear blight in  
194 the UK combined a crop growth model and a weather-based disease model with simulated  
195 future climate data (Madgwick *et al.*, 2011; Figure 5). The incidence of fusarium ear blight  
196 was related to rainfall during anthesis and temperature during the preceding 6 weeks. It was  
197 projected that, with climate change, wheat anthesis dates will be approximately two weeks  
198 earlier than at present. As a result, the rain-related risk of infection at anthesis did not  
199 decrease, as would have been predicted if anthesis had remained in mid-June (rainfall is  
200 projected to be almost unchanged in May but substantially reduced in June). Due to wetter



201 and warmer conditions in spring, the model predicted a slight increase in severity of fusarium  
202 ear blight epidemics by the 2050s, particularly in southern England (Madgwick *et al.*, 2011).  
203 This predicted slight increase reflected purely the weather-related risk and did not include  
204 effects of other indirect factors discussed in the following sections.

205

206

207 Fig 5 near here\*\*\*\*\*

208

209 In the long-term, climate appears to influence the predominant *Fusarium* species  
210 occurring in a given location. Bottalico (1998) suggests that differences in the predominant  
211 mycotoxin-producing *Fusarium* species present in cereals between northern and southern  
212 Europe are primarily due to differences in their survival on different substrates and timing of  
213 spore release relative to wheat anthesis. Furthermore, warm dry weather from autumn to  
214 early spring in the UK appears to increase risk by increasing inoculum build-up (Anon 2007).  
215 In northern Europe, climate change may be a factor responsible for a recent shift from *F.*  
216 *culmorum* to *F. graminearum* as the main cause of FEB (Waalwijk *et al.*, 2008). A similar  
217 change has been observed in maize, where the predominant *Fusarium* species was *F.*  
218 *graminearum* but increasingly the predominant species are others with favoured by warmer  
219 conditions, such as the *Gibberella fujikuroi* complex: *F. verticillioides*, *F. proliferatum*  
220 and/or *F. subglutinans*, which produce several mycotoxins (Waalwijk *et al.* 2008). However,  
221 changes to the climate of the UK and northern Europe have been only very slight over the  
222 past 30 years (Fig. 1). Although greater changes are predicted for the future, it seems that  
223 other factors, such as tillage, maize cultivation, fungicide use and cultivar resistance appear  
224 most likely to have contributed to the change in prevalence of *F. graminearum* and *F.*  
225 *culmorum* reported since 2002 (Anon 2010).

226

227

## 228 **Inoculum availability and crop cultivation**

229 In the UK, the incidence and severity of FEB epidemics on wheat varies from year to year but  
230 there has been a trend for the disease to increase in importance over the last decade  
231 (www.cropmonitor.co.uk; Fig. 4). In the UK, FEB is associated with a range of species in the  
232 genera *Fusarium* and *Microdochium*. Severe epidemics of FEB occur when there is a  
233 coincidence of a large amount of inoculum, suitable weather for infection and host plants at a  
234 susceptible growth stage, i.e. at flowering. The amount of inoculum is enhanced by increased  
235 cultivation of cereal hosts, especially maize, and by increased use of direct drilling or  
236 minimal tillage (Bateman *et al.* 2007; Eiblmeier *et al.*, 2007), which provides the pathogen  
237 with more plant residues on the soil surface on which to over-winter and sporulate (Teich &  
238 Hamilton, 1985; Windels, 2000; Champeil *et al.*, 2004b; Šíp *et al.*, 2007). Weather  
239 influences inoculum availability; perithecia of *G. zaeae* were reported to develop on debris and  
240 grains (of maize or wheat) on the soil surface (requiring light for development) at  
241 temperatures between 15-30°C (Gilbert & Fernando, 2004; Trail *et al.*, 2002), while *F.*  
242 *culmorum* conidia are also produced on debris on the soil surface (Bateman *et al.*, 2007).  
243 Conidia (both macro- and microconidia) are also produced on fusarium foot (stem-base) rots  
244 or from superficial infections on leaves (Anon 2010). These factors, other climatic factors  
245 and geology (soil type) affect each of the FEB-causing species and their interaction with the  
246 host in different ways; this then affects the relative amounts of inoculum of each species.

247 Due to different responses of the ear blight fungi to weather (see previous section), in  
248 any season the coincidence of spore release and wheat flowering may favour one species over  
249 another. This fits with the principle of competitive exclusion (Gause, 1934), which states  
250 that two species occupying the same niche cannot coexist indefinitely, while those occupying

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

251 different niches can coexist. DON contamination of grain is most severe in wheat crops that  
252 are preceded by grain maize (Eiblmeier *et al.*, 2007). This is thought to be because the large  
253 mass of infected crop debris that remains on the soil surface after harvesting of grain maize  
254 has a substantial capacity for production of wind-dispersed ascospores of *G. zeae* (*F.*  
255 *graminearum*) and persists in soil for a long time (Beck & Lepschy, 2000; Bateman *et al.*,  
256 2007). Since airborne inoculum of *F. graminearum* was found by Maldonado-Ramirez *et al.*  
257 (2005) to be dispersed regionally in the planetary boundary layer, it follows that regions with  
258 intense wheat and, or maize cultivation are likely to have greater concentrations of airborne  
259 inoculum. In the UK, grain maize is restricted to Southern England and the greatest density  
260 of forage maize crops is in areas of livestock production, especially the southwest where  
261 there are more cattle (Figs 6a and 6b). Climate change is predicted to cause the summer  
262 months to become warm enough for grain maize to be economically viable over a larger area  
263 of the UK (Kenny & Harrison, 1992). Provided that suitable spring weather promotes  
264 sporulation, additional maize cropping has potential to cause a substantial increase in FEB on  
265 UK wheat due to increased inoculum availability.

266  
267 Figure 6 here.

### 270 **Crop and crop-protection factors**

271 In addition to maize cultivation and minimal tillage, other agronomic and crop-protection  
272 factors influence the amount of FEB and the predominant species occurring. Commercially  
273 available fungicides can reduce the incidence of FEB by 50-70% when applied at the correct  
274 time, during flowering (Jennings 2002; Jennings *et al.*, 2000). However fungicide  
275 applications at a sub-optimal time may result in little effect (Jennings *et al.*, 2000). D'Mello

276 *et al.* (1998) found increases in DON and other mycotoxins occurred after sub-lethal  
277 exposures to MBC and triazole fungicides *in vitro*, while others suggest that some strobilurin  
278 (QoI) fungicides may increase DON production (Eiblmeier *et al.*, 2007; Blandino & Reyneri,  
279 2009; Jennings *et al.*, 2000; Edwards *et al.*, 2001; Simpson *et al.*, 2001). Even when  
280 fungicides are applied to protect most wheat ears at the correct time, ears of late-developing  
281 tillers are likely to be missed. These late tillers also tend to be shorter and are therefore in a  
282 more humid microclimate within the crop canopy, which increases the probability of  
283 infection. Furthermore, recent European legislation (EC No 1107/2009 repeal of directive  
284 91/414/EEC) will mean that fewer fungicides are available to control FEB in the future.

285 Cultivar resistance affects the severity of FEB and toxin production and is classified  
286 into five main types: type I is resistance to initial infection, type II is resistance to the spread  
287 of the pathogen after the initial infection, types III - V refer to the resistance to DON  
288 accumulation and tolerance of the disease (reviewed by Champeil *et al.*, 2004a; Buerstmayr  
289 *et al.*, 2009). One of the best known sources of resistance is the Chinese variety Sumai 3,  
290 which shows excellent type II resistance (Bai & Shaner, 2004) and a much lower level of  
291 DON accumulation in the harvested grain. Numerous other sources of resistance to FEB  
292 have also been identified and the underlying QTL identified (reviewed by Buerstmayr *et al.*,  
293 2009). Generally the difference in disease severity due to resistance that is currently  
294 available in UK varieties is relatively small with none completely resistant (Anon 2009a).  
295 Further research is in progress to improve and understand the different mechanisms of host  
296 resistance, which have great potential in the integrated management of FEB.

297 Cultural practices and chemical treatments that affect canopy density and crop height  
298 may affect spore dispersal and deposition on wheat ears (particularly for rain-splashed  
299 conidia) and the presence of certain weeds may influence the amount of inoculum and the  
300 predominant FEB species since several common weeds are hosts of *F. culmorum* and *F.*

1 301 *graminearum* (Champeil *et al.*, 2004a). Additionally, Grewal *et al.* (1996) reported that zinc  
2 302 deficiency in soil reduced resistance to foot rot (crown rot) caused by *F. graminearum* in  
3  
4 303 wheat, which may increase inoculum availability for FEB. Other mineral deficiencies may  
5  
6  
7 304 prolong flowering in wheat due to poor pollination success and this could also increase  
8  
9 305 susceptibility to FEB.

11 306

14 307

### 17 308 **Integrated risk assessment**

19 309 A mycotoxin risk assessment tool that is available for UK growers integrates some of these  
20  
21 310 different factors affecting fusarium mycotoxin risk. It can assist early decision-making about  
22  
23  
24 311 crop management, such as tillage or fungicide regime, and allow farmers to indicate the  
25  
26 312 relative fusarium risk for a field according to region of production, previous crop, cultivation  
27  
28  
29 313 method, cultivar resistance, anthesis-applied fungicide (application time known as T3 in the  
30  
31 314 UK), rainfall at flowering and pre-harvest rainfall (Anon 2009b). Acceptance of risk  
32  
33  
34 315 assessments by industries such as millers could reduce the need for mycotoxin testing in most  
35  
36 316 cases, i.e. where risk is predicted to be small.

39 317

41 318

### 44 319 **Conclusions: impacts of climate change on FEB**

46 320 Disease surveys in the UK have shown that *F. graminearum* is becoming more common on  
47  
48  
49 321 wheat ears than *F. culmorum*, with the incidence of *F. graminearum* in the UK exceeding that  
50  
51 322 of *F. culmorum* since 2002 (Anon 2010; Jennings *et al.*, 2004b; Gosman *et al.*, 2007).  
52  
53  
54 323 However, such changes in prevalence of *Fusarium* species have occurred too recently to  
55  
56 324 conclude that they are due to climate change (see Fig. 1). Although it is clear that weather  
57  
58  
59 325 influences epidemics of different *Fusarium* species in different ways, many changes in

60

61

62

63

64

65

1 326 farming practices, such as tillage regime, maize cultivation and fungicide use, have also  
2  
3 327 occurred recently. It is likely that climate change will have direct impacts on the prevalence  
4  
5 328 and severity of FEB and associated DON production in wheat crops as weather is one of the  
6  
7 329 main factors affecting the severity of epidemics and the relative proportions of different ear  
8  
9 330 blight pathogens responsible. The UK is in a unique situation due to its very variable climate  
10  
11 331 and complex of several interacting species of FEB pathogens. There are indications that the  
12  
13 332 predominant species have changed recently in association with a range of cultural changes  
14  
15 333 and possibly weather-related factors. It is unclear whether *F. graminearum* and *F. culmorum*  
16  
17 334 and other FEB fungi will respond to changes in climate in terms of inoculum production  
18  
19 335 exactly to coincide with the advancement in wheat flowering date predicted as a result of  
20  
21 336 UKCIP climate projections (Semenov, 2009).  
22  
23  
24  
25

26 337 Predicted drier summers may, despite the continued presence of FEB, result in  
27  
28 338 reduced DON contamination, because there will be a decrease in pre-harvest rainfall (from  
29  
30 339 GS 87 or hard-dough stage to harvest). Preharvest rainfall can increase DON production  
31  
32 340 substantially (Anon 2009b). However, this conclusion is based on a long term predicted  
33  
34 341 rainfall trend, while in practice there will be considerable variation in summer rainfall from  
35  
36 342 one year to another in the UK.  
37  
38  
39  
40

41 343 Furthermore, an indirect effect of climate change will be enhanced grain maize  
42  
43 344 cropping, which will substantially increase the capacity for inoculum production. It seems  
44  
45 345 likely that one reason why the occurrence of *F. graminearum* is much greater in North  
46  
47 346 America and continental Europe than in the UK is that the density of grain maize is much  
48  
49 347 greater in these areas and that maize debris provides a potent source of *F. graminearum*  
50  
51 348 ascospores for infecting wheat crops. The culture-based assessment method used by Schmale  
52  
53 349 *et al.* (2006) and Maldonado-Ramirez *et al.* (2005) did not identify the airborne inoculum  
54  
55 350 source, which could be ascospores, conidia or hyphal fragments, but it did show the presence  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 352  
4  
5 353  
6  
7 354  
8  
9  
10 355  
11  
12 356  
13  
14  
15  
16  
17 358  
18  
19 359  
20  
21  
22 360  
23  
24 361  
25  
26  
27 362  
28  
29 363  
30  
31 364  
32  
33  
34 365  
35  
36 366  
37  
38  
39 367  
40  
41 368  
42  
43

of inoculum dispersed evenly over the entire region. In contrast to the situation in North America, a lack of a homogeneous regional air-spora of *F. graminearum* (*G. zeae*) due to sparse occurrence of infected debris is thought to explain why weather-based disease models developed in other parts of the world frequently overestimate observed disease in the UK (Madgwick *et al.*, 2011). If inoculum availability is heterogeneous in the UK, then disease forecasting schemes that include a component of inoculum detection will be important.

14 357  
15  
16  
17 358  
18  
19 359  
20  
21  
22 360  
23  
24 361  
25  
26  
27 362  
28  
29 363  
30  
31 364  
32  
33  
34 365  
35  
36 366  
37  
38  
39 367  
40  
41 368  
42  
43

The impact of climate change on FEB is likely to have implications for breeders, growers and policy makers. Strategies to breed for resistance and to develop methods to reduce inoculum (crop debris), combined with effective crop protection products will continue to be needed to maintain a safe supply of grain. Forecasting schemes to target control options (fungicides applied at anthesis) to high risk seasons and, or locations would avoid unnecessary fungicide applications in low-risk sites. Further research to understand crop exposure to pathogen inoculum could allow a climate- or direct inoculum-based disease forecast to be issued in time to guide spray applications at anthesis. Disease forecasting and further research to improve crop protection solutions for growers must aim to mitigate the increased risk of FEB predicted under climate change.

#### 44 369 **Acknowledgements**

45  
46 370  
47  
48  
49 371  
50  
51 372  
52  
53 373  
54  
55  
56 374  
57  
58 375  
59  
60  
61  
62  
63  
64  
65

We thank the Department for Environment, Food and Rural Affairs (Defra Project FO 0302 and LK09111) and the UK Home-Grown Cereals Authority (AHDB-HGCA RD-2008-3573 and RD-2007-339) for funding this work and the HGCA for Fig. 3c, Martin Urban (Rothamsted Research) for Fig 3b. Rothamsted Research receives grant-aided support from the UK Biotechnology and Biological Sciences Research Council (BBSRC) for the Bioenergy and Climate Change ISPG.

376

377

## 378 **References**

379 Abramson D, Clear RM, Nowicki TW, 1987. *Fusarium* species and trichothecene

380 mycotoxins in suspect samples of 1985 Manitoba wheat. *Canadian Journal of Plant*

381 *Science* **67**: 611-620.

382 Anon. 2007. Guidelines to minimise risk of *Fusarium* mycotoxins in cereals. HGCA London.

383 Anon. 2009a. HGCA recommended list 2009/10 for cereals and oilseeds. AHDB-HGCA

384 (<http://www.hgca.com>).

385 Anon. 2009b. HGCA topic sheet 104: Managing *Fusarium* mycotoxin risk in wheat –

386 changes for harvest 2009, AHDB-HGCA,

387 [http://www.hgca.com/document.aspx?fn=load&media\\_id=5320&publicationId=6154](http://www.hgca.com/document.aspx?fn=load&media_id=5320&publicationId=6154)

388 Anon. 2010.

389 [https://secure.fera.defra.gov.uk/cropmonitor/wwheat/encyclopaedia/fusariumEB/inc\\_200](https://secure.fera.defra.gov.uk/cropmonitor/wwheat/encyclopaedia/fusariumEB/inc_200)

390 9.cfm (accessed March 2010).

391 Anderson PK, Cunningham AA, Patel NG, Morales FJ, Epstein PR, Daszak P, 2004.

392 Emerging infectious diseases of plants: pathogen pollution, climate change and

393 agrotechnology drivers. *Trends in Ecology and Evolution* **19**: 535-544.

394 Bai GH, Shaner G, 2004. Management and resistance in wheat and barley to fusarium head

395 blight. *Annual Review of Phytopathology* **42**: 135-161.

396 Bateman GL, Gutteridge RJ, Gherbawy Y, Thomsett MA, Nicholson P, 2007. Infection of

397 stem bases and grains of winter wheat by *Fusarium culmorum* and *F. graminearum* and

398 effects of tillage method and maize-stalk residues. *Plant Pathology* **56**: 604–615.



- 399 Beck R, Lepschy J, 2000. Ergebnisse aus dem fusarium-monitoring 1989-1999 - einfluss der  
1  
2 400 produktionstechnischen faktoren fruchtfolge und bodenbearbeitung. *Bodenkultur und*  
3  
4 401 *Pflanzenbau* **4**: 39-47.
- 5  
6  
7 402 Beyer M, Verreet JA, Ragab WSM, 2005. Effect of relative humidity on germination of  
8  
9 403 ascospores and macroconidia of *Gibberella zae* and deoxynivalenol production.  
10  
11 404 *International Journal of Food Microbiology* **98**: 233-240.
- 12  
13  
14 405 Blandino M, Reyneri A, 2009. Effect of fungicide and foliar fertilizer application to winter  
15  
16 406 wheat at anthesis on flag leaf senescence, grain yield, flour bread-making quality and  
17  
18 407 DON contamination. *European Journal of Agronomy* **30**: 275–282.
- 19  
20  
21 408 Bottalico A, 1998. *Fusarium* diseases of cereals: Species complex and related mycotoxin  
22  
23 409 profiles, in Europe. *Journal of Plant Pathology* **80**: 85-103.
- 24  
25  
26 410 Brennan JM, Fagan B, van Maanen A, Cooke BM, Doohan FM, 2003. Studies on *in vitro*  
27  
28 411 growth and pathogenicity of European *Fusarium* fungi. *European Journal of Plant*  
29  
30 412 *Pathology* **109**: 577-587.
- 31  
32  
33 413 Brown NA, Urban M, van de Meene AML, Hammond-Kosack KE, 2010. The infection  
34  
35 414 biology of *Fusarium graminearum*: defining the pathways of spikelet to spikelet  
36  
37 415 colonisation in wheat ears. *Fungal Biology* **114**: 555-571.
- 38  
39  
40 416 Buerstmayr H, Ban T, Anderson JA, 2009. QTL mapping and marker-assisted selection for  
41  
42 417 fusarium head blight resistance in wheat: a review. *Plant Breeding* **128**: 1-26.
- 43  
44  
45 418 Chakraborty S, Newton AC, 2011. Climate change, plant diseases and food security, an  
46  
47 419 overview. *Plant Pathology*. **60**: 2-14.
- 48  
49  
50 420 Chakraborty S, Tiedemann AV, Teng PS, 2000. Climate change: potential impact on plant  
51  
52 421 diseases. *Environmental Pollution* **108**: 317-326.
- 53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 422 Champeil A, Doré T, Fourbet JF, 2004a. Fusarium head blight: epidemiological origin of the  
423 effects of cultural practices on head blight attacks and the production of mycotoxins by  
424 *Fusarium* in wheat grains. *Plant Science* **166**: 1389–1415.
- 425 Champeil A, Fourbet JF, Doré T, Rossignol L, 2004b. Influence of cropping system on  
426 fusarium head blight and mycotoxin levels in winter wheat. *Crop Protection* **23**: 531–  
427 537.
- 428 De Wolf ED, Madden LV, Lipps PE, 2003. Risk assessment models for wheat fusarium head  
429 blight epidemics based on within-season weather data. *Phytopathology* **93**: 428-435.
- 430 Doohan FM, Brennan J, Cooke BM, 2003. Influence of climatic factors on *Fusarium* species  
431 pathogenic to cereals. *European Journal of Plant Pathology* **109**: 755-768.
- 432 D'Mello JPF, Macdonald AMC, Postel D, Dijkma WTP, Dujardin A, Placinta CM, 1998.  
433 Pesticide use and mycotoxin production in *Fusarium* and *Aspergillus* phytopathogens.  
434 *European Journal of Plant Pathology* **104**: 741-751.
- 435 Edwards S G, Pirgozliev SR, Hare MC, Jenkinson P, 2001. Quantification of trichothecene  
436 producing *Fusarium* species in harvested grain by competitive PCR to determine  
437 efficacies of fungicides against fusarium head blight of winter wheat. *Applied and*  
438 *Environmental Microbiology* **67**: 1575-1580.
- 439 Eiblmeier P, von Gleissenthall JL, Lepschy J, 2007. Risk evaluation of deoxynivalenol levels  
440 in Bavarian wheat from survey data. *Journal of Plant Diseases and Protection* **114**: 69-  
441 75.
- 442 Fernandez MR, Chen Y, 2005. Pathogenicity of *Fusarium* species on different plant parts of  
443 spring wheat under controlled conditions. *Plant Disease* **89**: 164-169.
- 444 Garrett KA, Dendy SP, Frank EE, Rouse MN, Travers SE, 2006. Climate change effects on  
445 plant disease: genomes to ecosystems. *Annual Review of Phytopathology* **44**: 489-509.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 446 Gause GF, 1934. *The Struggle for Existence*. Baltimore: Williams and Wilkins. Reprinted in  
447 2003 as Dover Phoenix Editions, Dover Publications, 176pp.
- 448 Gilbert J, Tekauz A, 2000. Recent developments in research on fusarium head blight of wheat  
449 in Canada. *Canadian Journal of Plant Pathology* **22**: 1–8.
- 450 Gilbert J, Fernando WGD, 2004 Epidemiology and biological control of *Gibberella zeae* /  
451 *Fusarium graminearum*. *Canadian Journal of Plant Pathology* **26**: 464-472.
- 452 Gosman N, Bayles R, Jennings P, Kirby J, Nicholson P, 2007. Evaluation and  
453 characterization of resistance to fusarium head blight caused by *Fusarium culmorum* in  
454 UK winter wheat cultivars. *Plant Pathology* **56**: 264-276.
- 455 Goswami RS, Kistler HC, 2004. Heading for disaster: *Fusarium graminearum* on cereal  
456 crops. *Molecular Plant Pathology* **5**: 515-525.
- 457 Grewal HS, Graham RD, Rengel Z, 1996. Genotypic variation in zinc efficiency and  
458 resistance to crown rot disease (*Fusarium graminearum* Schw. Group 1) in wheat. *Plant*  
459 *and Soil* **186**: 219-226.
- 460 Hope R, Aldred D, Magan N 2005. Comparison of environmental profiles for growth and  
461 deoxynivalenol production by *Fusarium culmorum* and *F. graminearum* on wheat grain.  
462 *Letters in Applied Microbiology* **40**: 295-300.
- 463 Hooker DC, Schaafsma AW, Tamburic-Ilincic L, 2002. Using weather variables pre- and  
464 post-heading to predict deoxynivalenol content in winter wheat. *Plant Disease* **86**: 611-  
465 619.
- 466 Jansen C, von Wettstein D, Schafer W, Kogel KH, Felk A, Maier FJ, 2005. Infection patterns  
467 in barley and wheat spikes inoculated with wild-type and trichodiene synthase gene  
468 disrupted *Fusarium graminearum*. *Proceedings of the National Academy of Science of*  
469 *the United States of America* **102**: 16892–16897.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 470 Jennings P, 2002. Control of the fungus through use of fungicides, In O. E. Scholten, P.  
471 Ruckenbauer, A. Visconti, W. A. van Osenbruggen, and den Nijs A.P.M. (eds.), Food  
472 safety of Cereals: A chain-wide approach to reduce *Fusarium* mycotoxins. Final Report  
473 of EU project FAIR-CT-98-4094, p. 22-24.
- 474 Jennings P, Coates ME, Turner JA, Chandler EA, Nicholson P, 2004a. Determination of  
475 deoxynivalenol and nivalenol chemotypes of *Fusarium culmorum* isolates from England  
476 and Wales by PCR assay. *Plant Pathology* **53**: 182-190.
- 477 Jennings P, Coates ME, Walsh K, Turner JA, Nicholson P, 2004b. Determination of  
478 deoxynivalenol- and nivalenol-producing chemotypes of *Fusarium graminearum*  
479 isolated from wheat crops in England and Wales. *Plant Pathology* **53**: 643-652.
- 480 Jennings P, Turner JA, Nicholson P, 2000. Overview of fusarium ear blight in the UK - effect  
481 of fungicide treatment on disease control and mycotoxin production. Proceedings of the  
482 BCPC Conference Pests and Diseases 2000 **2**: 707-712.
- 483 Jestoi M, Kokkonen M, Uhlig S, 2009. What about the “other” *Fusarium* mycotoxins? *World*  
484 *Mycotoxin Journal* **2**: 181-192.
- 485 Kang Z, Buchenauer H, 2000. Ultrastructural and cytochemical studies on cellulose, xylan  
486 and pectin degradation in wheat spikes infected by *Fusarium culmorum*. *Journal of*  
487 *Phytopathology* **148**: 263-275.
- 488 Kenny GJ, Harrison PA, 1992. Thermal and moisture limits of grain maize in Europe: model  
489 testing and sensitivity to climate change. *Climate Research* **2**: 113-129.
- 490 Kostelanska M, Hajslova J, Zachariasova M, Malachova, A, Kalachova K, Poustka J, Fiala J,  
491 Scott PM, Berthiller F, Krska R, 2009. Occurrence of deoxynivalenol and its major  
492 conjugate, deoxynivalenol-3-glucoside, in beer and some brewing intermediates. *Journal*  
493 *of Agricultural and Food Chemistry* **57**: 3187-3194

- 494 Lin F, Kong ZX, Zhu HL, Xue SL, Wu JZ, Tian DG, Wei JB, Zhang CQ, Ma ZQ, 2004.  
1  
2  
3 495 Mapping QTL associated with resistance to fusarium head blight in the Nanda2419 x  
4  
5 496 Wangshuibai population. I. Type II resistance. *Theoretical and Applied Genetics* **109**:  
6  
7 497 1504-1511.  
8  
9  
10 498 Madgwick J, West JS, White R, Semenov M, Townsend JA, Turner JA, Fitt BDL. 2011.  
11  
12 499 Future threat; direct impact of climate change on wheat fusarium ear blight in the UK.  
13  
14 500 *European Journal of Plant Pathology* (in press).  
15  
16  
17  
18 501 Maier FJ, Miedaner T, Hadelar B, Felk A, Salomon S, Lemmens M, Kassner H, Schäfer W,  
19  
20 502 2006. Involvement of trichothecenes in fusarioses of wheat, barley and maize evaluated  
21  
22 503 by gene disruption of the trichodiene synthase (*Tri5*) gene in three field isolates of  
24  
25 504 different chemotype and virulence. *Molecular Plant Pathology* **7**: 449–461.  
26  
27 505 Maldonado-Ramirez SL, Schmale DG, Shields EJ, Bergstrom GC, 2005. The relative  
28  
29 506 abundance of viable spores of *Giberella zeae* in the planetary boundary layer suggests a  
31  
32 507 role of long-distance transport in regional epidemics of fusarium head blight.  
33  
34 508 *Agricultural and Forest Meteorology* **132**: 20-27.  
36  
37 509 McMullen M, Jones R, Gallenberg D, 1997. Scab of wheat and barley: A re-emerging disease  
38  
39 510 of devastating impact. *Plant Disease* **81**: 1340-1348.  
41  
42 511 Moschini RC, Pioli R, Carmona M, Sacchi O, 2001. Empirical predictions of wheat head  
43  
44 512 blight in the Northern Argentinean Pampas region. *Crop Science* **41**: 1541-1545.  
46  
47 513 Moss MO, 2002. Mycotoxin review - 2. Fusarium. *Mycologist* **16**: 158-161.  
48  
49 514 Parry DW, Jenkinson P, McLeod L, 1995. Fusarium ear blight (scab) in small-grain cereals -  
50  
51 515 a review. *Plant Pathology* **44**: 207-238.  
53  
54 516 Paul PA, El-Allaf SM, Lipps PE, Madden LV, 2004. Rain splash dispersal of *Gibberella zeae*  
55  
56 517 within wheat canopies in Ohio. *Phytopathology* **94**: 1342-1349.  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 518 Proctor RH, Hohn TM, McCormick SP, 1995. Reduced virulence of *Gibberella zeae* caused  
519 by disruption of a trichothecene toxin biosynthetic gene. *Molecular Plant Microbe*  
520 *Interactions* **8**: 593–601.
- 521 Ruckenbauer P, Buerstmayr H, Lemmens M, 2001. Present strategies in resistance breeding  
522 against scab (*Fusarium* spp.). *Euphytica* **119**: 121-127.
- 523 Schmale DG, Arntsen QA, Bergstrom GC, 2005. The forcible discharge distance of  
524 ascospores of *Gibberelia zeae*. *Canadian Journal of Plant Pathology* **27**: 376-382.
- 525 Schmale DG, Shields EJ, Bergstrom GC, 2006. Night time spore deposition of the fusarium  
526 head blight pathogen, *Giberella zeae*, in rotational wheat fields. *Canadian Journal of*  
527 *Plant Pathology* **28**: 100-108.
- 528 Schmidhuber J, Tubiello FN, 2007. Global food security under climate change. *Proceedings*  
529 *of the National Academy of Science of the United States of America* **104**: 19703-19708.
- 530 Semenov M, 2009. Impacts of climate change on wheat in England and Wales. *Journal of the*  
531 *Royal Society Interface* **6**: 343-350.
- 532 Simpson DR, Weston GE, Turner JA, Jennings P, Nicholson P, 2001. Differential control of  
533 head blight pathogens of wheat by fungicides and consequences for mycotoxin  
534 contamination of grain. *European Journal of Plant Pathology* **107**: 421-431.
- 535 Sinha RC, Savard ME, 1997. Concentration of deoxynivalenol in single kernels and various  
536 tissues of wheat heads. *Canadian Journal of Plant Pathology* **19**: 8-12.
- 537 Šíp V, Chrptová J, Leišová L, Sýkorová S, Kučera L, Ovesná J, 2007. Implications for  
538 fusarium head blight control from study of factors determining pathogen and DON  
539 content in grain of wheat cultivars. In: Buck HT, Nisi JE, Salomón N, eds. *Developments*  
540 *in Plant Breeding “Wheat Production in Stressed Environments”*. Springer, the  
541 Netherlands, 281–287.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 542 Stern N, 2007. *The Economics of Climate Change: The Stern Review*. Cambridge University  
543 Press, UK.
- 544 Sutton JC, Balico W, Funnel HS, 1980. Relation of weather variables to incidence of  
545 zearalenone in corn in southern Ontario. *Canadian Journal of Plant Science* **60**: 149-  
546 155.
- 547 Teich AH, Hamilton JR, 1985. Effect of cultural practices, soil phosphorus, potassium, and  
548 pH on the incidence of fusarium head blight and deoxynivalenol levels in wheat. *Applied*  
549 *and Environmental Microbiology* **49**: 1429-1431.
- 550 Trail F, Xu H, Loranger R, Gadoury D, 2002. Physiological and environmental aspects of  
551 ascospore discharge in *Gibberella zeae* (anamorph *Fusarium graminearum*). *Mycologia*  
552 **94**: 181-189.
- 553 Waalwijk C, van der Lee T, Yang L, de Vries I, Görtz A, Kema G, 2008. Are changes in the  
554 composition of the fusarium head blight complex caused by climate change? Pest and  
555 Climate Change, December 3, 2008. Abstracts.
- 556 Wei Y-M, Hou Y-C, Yan Z-H, Wu W, Zhang Z-Q, Liu D-C, Zheng Y-L, 2005.  
557 Microsatellite DNA polymorphism divergence in Chinese wheat (*Triticum aestivum* L.)  
558 landraces highly resistant to fusarium head blight. *Journal of Applied Genetics* **46**: 3-9.
- 559 Windels CE, 2000. Economic and social impacts of fusarium head blight: Changing farms  
560 and rural communities in the Northern Great Plains. *Phytopathology* **90**: 17-21.
- 561 Xu X, Nicholson P, 2009. Community ecology of fungal pathogens causing wheat head  
562 blight. *Annual Review of Phytopathology* **47**: 83-103.
- 563 Xu XM, Nicholson P, Thomsett MA, Simpson D, Cooke BM, Doohan FM, Brennan J,  
564 Monaghan S, Moretti A, Mule G, Homok L, Beki E, Tatnell J, Ritieni A, Edwards SG,  
565 2008. Relationship between the fungal complex causing fusarium head blight of wheat  
566 and environmental conditions *Phytopathology* **98**: 69-78.

567

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

568 Figure Legends:

569

570 **Fig 1 - Annual mean temperature from 1914 to 2004 for different regions of the UK,**  
571 **smoothed with a triangular kernel filter with 14 terms on either side of each target**  
572 **point. © Crown copyright 2006, the Met Office, used with permission.**

573

574 **Fig 2 - Baseline 30-year (1971-2000) monthly mean rainfall and temperature at**  
575 **Rothamsted, UK (51° 48' 0" N, 0° 21' 0" W) and projected monthly means for 2050**  
576 **according to the high emission (2050HI) climate scenario HadRM3 climate model.**

577

578 **Fig 3 - Fusarium ear blight in a winter wheat crop, cultivar Hereward, Rothamsted field**  
579 **experiment, July 2007 (A); *Fusarium graminearum* macroconidia (B); fusarium foot rot**  
580 **(C); comparison between healthy harvested grain (left) and *Fusarium* infected grain**  
581 **(right) (D).**

582

583 **Fig 4 - Incidence (% crops affected) of fusarium ear blight on winter wheat at GS 75 in**  
584 **England and Wales. Results from the CropMonitor survey of winter wheat crops (1991**  
585 **- 2008, <http://www.cropmonitor.co.uk/>)**

586

587 **Fig 5 - Maps showing the projected average fusarium ear blight incidence (% plants**  
588 **affected) generated by a fusarium ear blight model and based on advanced anthesis**  
589 **dates for three weather scenarios; baseline, 2020 high emission scenario (2020HI), and**



590 2050 high emission scenario (2050HI). The baseline scenario is based on weather from  
1  
2 591 1960-1990. The maps were produced by spatial interpolation between the 14 sites.  
3  
4  
5 592 Adapted from Madgwick *et al.* (2011).  
6

7 593  
8  
9 594  
10

11 **Fig 6 - Relationship between number of cattle and area of maize grown in counties of**  
12 **England in 2008 (A) and maize distribution in England in 2008 as a percentage of the**  
13 **total arable area in each county (B). Maize is a potent source of ascospores of**  
14 ***Gibberella zeae* (*F. graminearum*). Data ([www.defra.gov.uk](http://www.defra.gov.uk)); no data were available for**  
15 **arable areas in Scotland and Wales (shaded grey).**  
16  
17  
18  
19  
20  
21  
22  
23

24 600  
25  
26  
27 601  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Figure

[Click here to download high resolution image](#)




Figure2

[Click here to download high resolution image](#)




Figure3

[Click here to download high resolution image](#)



Figure4

[Click here to download high resolution image](#)




Figure5




Figure6  
[Click here to download high resolution image](#)

